
089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_positive_max_area_50C
Pearson r=-0.025 (p=0.892) | Spearman ρ=-0.134 (p=0.472) | Kendall τ=-0.071 (p=0.585)
Pearson log1p(activity) r=-0.095 (p=0.612)
PA6 activity (µM)
SURFMATCH score_positive_max_area
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
UMGSP1
HotPETase
FsC
NfpolyA
ThcCut1
LCC_ICCG
UMGSP3
Tfcut2
HiC
ph_NylC
TtA
ReU
MGS0156
ThcCut2
NylB
Cut190
Trypsin
MsmegA
AoC
Ooh
chymotrypsin
Papain
Subtilisin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.001\nRMSE = 89.19 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.20 (p=0.672)
Spearman ρ=0.14 (p=0.758)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.46 (p=0.178)
Spearman ρ=-0.13 (p=0.712)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.00 (p=0.997)
Spearman ρ=0.11 (p=0.78)