
108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_negative_max_area_50C
Pearson r=-0.194 (p=0.295) | Spearman ρ=-0.124 (p=0.507) | Kendall τ=-0.091 (p=0.493)
Pearson log1p(activity) r=-0.131 (p=0.483)
PA6 activity (µM)
SURFMATCH score_negative_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.038\nRMSE = 87.52 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.16 (p=0.733)
Spearman ρ=-0.17 (p=0.711)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.20 (p=0.583)
Spearman ρ=-0.34 (p=0.331)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.19 (p=0.625)
Spearman ρ=0.32 (p=0.394)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
GatA
nc_NylC
UMGSP1
NfpolyA
HotPETase
FsC
Tfcut2
ThcCut1
LCC_ICCG
TtA
UMGSP3
Trypsin
HiC
ph_NylC
ReU
ThcCut2
MGS0156
NylB
MsmegA
AoC
Ooh
Subtilisin
Cut190
Papain
chymotrypsin