
108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_positive_total_area_50C
Pearson r=0.425 (p=0.0172) | Spearman ρ=0.421 (p=0.0182) | Kendall τ=0.302 (p=0.0186)
Pearson log1p(activity) r=0.415 (p=0.0201)
PA6 activity (µM)
SURFMATCH score_positive_total_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.180\nRMSE = 80.76 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.26 (p=0.571)
Spearman ρ=-0.18 (p=0.696)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.19 (p=0.6)
Spearman ρ=-0.20 (p=0.578)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.55 (p=0.127)
Spearman ρ=0.61 (p=0.0826)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
PSC
UMGSP1
ThcCut1
LCC_ICCG
MGS0156
HiC
Trypsin
NylB
Papain
Ooh
Cut190
chymotrypsin
Subtilisin
ThcCut2
Tfcut2
AoC
NfpolyA
UMGSP3
ph_NylC
ReU
TtA
MsmegA