
127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_negative_max_area_50C
Pearson r=-0.279 (p=0.129) | Spearman ρ=-0.200 (p=0.28) | Kendall τ=-0.146 (p=0.26)
Pearson log1p(activity) r=-0.220 (p=0.234)
PA6 activity (µM)
SURFMATCH score_negative_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.078\nRMSE = 85.68 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.14 (p=0.762)
Spearman ρ=0.41 (p=0.357)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.03 (p=0.938)
Spearman ρ=0.08 (p=0.834)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.17 (p=0.669)
Spearman ρ=-0.06 (p=0.88)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
NfpolyA
ThcCut1
FsC
UMGSP1
HotPETase
ThcCut2
UMGSP3
LCC_ICCG
HiC
TtA
ph_NylC
Trypsin
Tfcut2
ReU
MGS0156
Cut190
Subtilisin
AoC
MsmegA
Ooh
NylB
Papain
chymotrypsin