
127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_negative_total_area_50C
Pearson r=0.413 (p=0.0208) | Spearman ρ=0.229 (p=0.215) | Kendall τ=0.164 (p=0.201)
Pearson log1p(activity) r=0.274 (p=0.136)
PA6 activity (µM)
SURFMATCH score_negative_total_area
0
50
100
150
200
250
300
140
160
180
200
220
240
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
FsC
UMGSP1
NfpolyA
UMGSP3
LCC_ICCG
HiC
ThcCut1
ThcCut2
Tfcut2
AoC
Trypsin
MGS0156
Subtilisin
Cut190
NylB
Ooh
TtA
ph_NylC
ReU
Papain
MsmegA
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.171\nRMSE = 81.24 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.01 (p=0.986)
Spearman ρ=-0.13 (p=0.788)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.76 (p=0.011)
Spearman ρ=-0.72 (p=0.0182)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.35 (p=0.355)
Spearman ρ=0.37 (p=0.325)