
146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_global_score_50C
Pearson r=0.368 (p=0.0419) | Spearman ρ=0.337 (p=0.0638) | Kendall τ=0.225 (p=0.0792)
Pearson log1p(activity) r=0.385 (p=0.0325)
PA6 activity (µM)
SURFMATCH global score
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.135\nRMSE = 82.97 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.71 (p=0.0741)
Spearman ρ=0.63 (p=0.129)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.29 (p=0.416)
Spearman ρ=0.19 (p=0.59)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.31 (p=0.417)
Spearman ρ=0.18 (p=0.65)
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
nc_NylC
GatA
UMGSP2
HotPETase
NfpolyA
UMGSP1
FsC
UMGSP3
MGS0156
MsmegA
ThcCut1
LCC_ICCG
HiC
Tfcut2
ph_NylC
TtA
NylB
ThcCut2
Cut190
AoC
Trypsin
Papain
chymotrypsin
PleU
Ooh
Subtilisin