
001 | orientation selected from FORMULA global_score
FORMULA_negative_max_area_50C
Pearson r=-0.299 (p=0.103) | Spearman ρ=-0.096 (p=0.606) | Kendall τ=-0.086 (p=0.515)
Pearson log1p(activity) r=-0.144 (p=0.441)
PA6 activity (µM)
SURFMATCH score_negative_max_area
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
NfpolyA
FsC
UMGSP1
LCC_ICCG
UMGSP3
ThcCut1
Tfcut2
ph_NylC
MsmegA
ThcCut2
TtA
HiC
NylB
Ooh
Cut190
ReU
MGS0156
AoC
Papain
chymotrypsin
Subtilisin
Trypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=-0.02, p=0.963)
Global statistics\nn = 31\nR² = 0.089\nRMSE = 85.14 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.02 (p=0.963)
Spearman ρ=-0.04 (p=0.938)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.11 (p=0.761)
Spearman ρ=0.24 (p=0.511)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.05 (p=0.907)
Spearman ρ=0.09 (p=0.811)