
012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_global_score_50C
Pearson r=-0.019 (p=0.918) | Spearman ρ=0.094 (p=0.617) | Kendall τ=0.055 (p=0.67)
Pearson log1p(activity) r=0.021 (p=0.909)
PA6 activity (µM)
SURFMATCH global score
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
GatA
nc_NylC
HotPETase
NfpolyA
LCC_ICCG
FsC
ThcCut1
UMGSP3
UMGSP1
HiC
ThcCut2
chymotrypsin
Papain
TtA
ReU
Tfcut2
ph_NylC
Trypsin
Subtilisin
NylB
MGS0156
MsmegA
AoC
Cut190
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=-0.01, p=0.986)
Global statistics\nn = 31\nR² = 0.000\nRMSE = 89.20 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.01 (p=0.986)
Spearman ρ=-0.18 (p=0.699)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.04 (p=0.91)
Spearman ρ=-0.12 (p=0.751)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.42 (p=0.261)
Spearman ρ=-0.11 (p=0.78)