
031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_negative_max_area_50C
Pearson r=-0.234 (p=0.206) | Spearman ρ=-0.216 (p=0.243) | Kendall τ=-0.151 (p=0.245)
Pearson log1p(activity) r=-0.248 (p=0.179)
PA6 activity (µM)
SURFMATCH score_negative_max_area
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
UMGSP1
NfpolyA
FsC
HotPETase
ThcCut1
UMGSP3
LCC_ICCG
ph_NylC
Tfcut2
TtA
ReU
MGS0156
ThcCut2
HiC
AoC
NylB
Papain
Subtilisin
MsmegA
Trypsin
Ooh
Cut190
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=0.66, p=0.105)
Global statistics\nn = 31\nR² = 0.055\nRMSE = 86.75 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.66 (p=0.105)
Spearman ρ=0.72 (p=0.0691)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.57 (p=0.0854)
Spearman ρ=-0.60 (p=0.0653)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.51 (p=0.156)
Spearman ρ=-0.44 (p=0.24)