
069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_positive_max_area_50C
Pearson r=-0.042 (p=0.823) | Spearman ρ=-0.102 (p=0.585) | Kendall τ=-0.058 (p=0.656)
Pearson log1p(activity) r=-0.077 (p=0.682)
PA6 activity (µM)
SURFMATCH score_positive_max_area
0
50
100
150
200
250
300
10
15
20
25
30
35
40
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
UMGSP1
LCC_ICCG
UMGSP3
ThcCut2
NylB
Ooh
AoC
MsmegA
chymotrypsin
Trypsin
Tfcut2
ThcCut1
Cut190
Papain
TtA
MGS0156
ph_NylC
Subtilisin
NfpolyA
ReU
HiC
FsC
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=0.06, p=0.906)
Global statistics\nn = 31\nR² = 0.002\nRMSE = 89.14 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.06 (p=0.906)
Spearman ρ=0.01 (p=0.985)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.24 (p=0.499)
Spearman ρ=-0.17 (p=0.646)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.08 (p=0.831)
Spearman ρ=-0.07 (p=0.854)