078 | orientation selected from PEARSON global_score
PEARSON_negative_total_area_50C
Pearson r=0.557 (p=0.00112) | Spearman ρ=0.420 (p=0.0188) | Kendall τ=0.299 (p=0.0204)
Pearson log1p(activity) r=0.457 (p=0.00983)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG1 NylC-like regression (r=0.14, p=0.771)

Global statistics\nn = 31\nR² = 0.311\nRMSE = 74.06 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.14 (p=0.771)
  Spearman ρ=-0.13 (p=0.788)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.51 (p=0.129)
  Spearman ρ=-0.55 (p=0.101)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.69 (p=0.0379)
  Spearman ρ=0.61 (p=0.0843)