
078 | orientation selected from PEARSON global_score
PEARSON_positive_max_area_50C
Pearson r=-0.215 (p=0.246) | Spearman ρ=-0.257 (p=0.162) | Kendall τ=-0.185 (p=0.156)
Pearson log1p(activity) r=-0.144 (p=0.44)
PA6 activity (µM)
SURFMATCH score_positive_max_area
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
UMGSP1
FsC
NfpolyA
LCC_ICCG
ThcCut1
UMGSP3
ThcCut2
Tfcut2
TtA
ph_NylC
HiC
ReU
NylB
AoC
Trypsin
Cut190
MsmegA
MGS0156
Papain
Subtilisin
Ooh
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=-0.39, p=0.385)
Global statistics\nn = 31\nR² = 0.046\nRMSE = 87.14 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.39 (p=0.385)
Spearman ρ=-0.18 (p=0.694)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.08 (p=0.819)
Spearman ρ=0.17 (p=0.634)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.31 (p=0.415)
Spearman ρ=-0.39 (p=0.304)