
089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_negative_total_area_50C
Pearson r=0.450 (p=0.0112) | Spearman ρ=0.282 (p=0.124) | Kendall τ=0.221 (p=0.0851)
Pearson log1p(activity) r=0.287 (p=0.117)
PA6 activity (µM)
SURFMATCH score_negative_total_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=-0.18, p=0.706)
Global statistics\nn = 31\nR² = 0.202\nRMSE = 79.69 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.18 (p=0.706)
Spearman ρ=-0.32 (p=0.478)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.35 (p=0.324)
Spearman ρ=-0.41 (p=0.241)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.07 (p=0.852)
Spearman ρ=0.07 (p=0.854)
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
FsC
UMGSP1
NfpolyA
UMGSP3
ThcCut1
LCC_ICCG
HiC
Tfcut2
ThcCut2
AoC
chymotrypsin
Trypsin
MGS0156
Cut190
NylB
Papain
Subtilisin
TtA
Ooh
MsmegA
ph_NylC
ReU