
089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_positive_total_area_50C
Pearson r=0.474 (p=0.00704) | Spearman ρ=0.413 (p=0.021) | Kendall τ=0.315 (p=0.0141)
Pearson log1p(activity) r=0.419 (p=0.0189)
PA6 activity (µM)
SURFMATCH score_positive_total_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=0.60, p=0.156)
Global statistics\nn = 31\nR² = 0.225\nRMSE = 78.55 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.60 (p=0.156)
Spearman ρ=0.59 (p=0.159)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.32 (p=0.37)
Spearman ρ=-0.32 (p=0.374)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.45 (p=0.224)
Spearman ρ=0.45 (p=0.22)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
UMGSP1
NfpolyA
FsC
ThcCut1
LCC_ICCG
Tfcut2
ThcCut2
MGS0156
AoC
Trypsin
Papain
Cut190
Subtilisin
chymotrypsin
HiC
NylB
ReU
ph_NylC
TtA
UMGSP3
Ooh
MsmegA