
108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_positive_max_area_50C
Pearson r=-0.221 (p=0.232) | Spearman ρ=-0.163 (p=0.381) | Kendall τ=-0.133 (p=0.305)
Pearson log1p(activity) r=-0.255 (p=0.166)
PA6 activity (µM)
SURFMATCH score_positive_max_area
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
UMGSP1
ThcCut1
FsC
NfpolyA
LCC_ICCG
UMGSP3
TtA
ThcCut2
Tfcut2
HiC
MGS0156
AoC
ph_NylC
Cut190
Trypsin
chymotrypsin
Ooh
ReU
NylB
Subtilisin
MsmegA
Papain
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=-0.40, p=0.377)
Global statistics\nn = 31\nR² = 0.049\nRMSE = 87.01 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.40 (p=0.377)
Spearman ρ=-0.43 (p=0.339)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.02 (p=0.946)
Spearman ρ=0.15 (p=0.671)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.03 (p=0.934)
Spearman ρ=0.24 (p=0.542)