
127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_positive_max_area_50C
Pearson r=-0.138 (p=0.459) | Spearman ρ=0.054 (p=0.772) | Kendall τ=0.031 (p=0.811)
Pearson log1p(activity) r=0.017 (p=0.926)
PA6 activity (µM)
SURFMATCH score_positive_max_area
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
NfpolyA
FsC
UMGSP1
LCC_ICCG
ThcCut1
TtA
UMGSP3
Tfcut2
HiC
ReU
MsmegA
ph_NylC
Ooh
MGS0156
ThcCut2
Trypsin
Papain
Cut190
NylB
AoC
Subtilisin
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=-0.41, p=0.355)
Global statistics\nn = 31\nR² = 0.019\nRMSE = 88.36 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.41 (p=0.355)
Spearman ρ=-0.07 (p=0.877)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.56 (p=0.0893)
Spearman ρ=0.46 (p=0.181)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.01 (p=0.985)
Spearman ρ=0.28 (p=0.461)