
146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_negative_max_area_50C
Pearson r=-0.225 (p=0.225) | Spearman ρ=-0.096 (p=0.608) | Kendall τ=-0.056 (p=0.669)
Pearson log1p(activity) r=-0.209 (p=0.26)
PA6 activity (µM)
SURFMATCH score_negative_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=0.51, p=0.241)
Global statistics\nn = 31\nR² = 0.050\nRMSE = 86.94 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.51 (p=0.241)
Spearman ρ=0.39 (p=0.386)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.27 (p=0.456)
Spearman ρ=0.59 (p=0.071)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.38 (p=0.318)
Spearman ρ=-0.40 (p=0.282)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
NfpolyA
FsC
UMGSP3
TtA
ThcCut2
HiC
ph_NylC
NylB
MGS0156
ReU
Ooh
MsmegA
Subtilisin
ThcCut1
UMGSP1
LCC_ICCG
Tfcut2
Trypsin
Cut190
AoC
chymotrypsin
Papain