
146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_negative_total_area_50C
Pearson r=0.411 (p=0.0217) | Spearman ρ=0.315 (p=0.0845) | Kendall τ=0.239 (p=0.0631)
Pearson log1p(activity) r=0.355 (p=0.0501)
PA6 activity (µM)
SURFMATCH score_negative_total_area
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
FsC
UMGSP1
HotPETase
NfpolyA
ThcCut1
LCC_ICCG
UMGSP3
Tfcut2
HiC
ThcCut2
AoC
Trypsin
Cut190
Subtilisin
MGS0156
NylB
Papain
chymotrypsin
TtA
Ooh
MsmegA
ph_NylC
ReU
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=-0.72, p=0.0681)
Global statistics\nn = 31\nR² = 0.169\nRMSE = 81.34 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.72 (p=0.0681)
Spearman ρ=-0.74 (p=0.0579)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.28 (p=0.442)
Spearman ρ=-0.40 (p=0.253)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.77 (p=0.0151)
Spearman ρ=0.62 (p=0.0738)