
146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_positive_max_area_50C
Pearson r=-0.018 (p=0.924) | Spearman ρ=-0.065 (p=0.73) | Kendall τ=-0.047 (p=0.72)
Pearson log1p(activity) r=0.006 (p=0.973)
PA6 activity (µM)
SURFMATCH score_positive_max_area
0
50
100
150
200
250
300
10
15
20
25
30
35
40
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
nc_NylC
GatA
UMGSP2
HotPETase
UMGSP1
FsC
LCC_ICCG
ThcCut1
NfpolyA
UMGSP3
HiC
Tfcut2
ph_NylC
Trypsin
ThcCut2
NylB
TtA
MsmegA
MGS0156
ReU
Papain
AoC
Cut190
Ooh
chymotrypsin
Subtilisin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=0.23, p=0.617)
Global statistics\nn = 31\nR² = 0.000\nRMSE = 89.20 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.23 (p=0.617)
Spearman ρ=0.20 (p=0.667)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.25 (p=0.482)
Spearman ρ=-0.15 (p=0.674)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.19 (p=0.62)
Spearman ρ=-0.17 (p=0.666)