001 | orientation selected from FORMULA global_score
FORMULA_positive_max_area_50C
Pearson r=-0.143 (p=0.442) | Spearman ρ=-0.261 (p=0.156) | Kendall τ=-0.188 (p=0.146)
Pearson log1p(activity) r=-0.140 (p=0.452)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.30, p=0.402)

Global statistics\nn = 31\nR² = 0.021\nRMSE = 88.30 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.19 (p=0.681)
  Spearman ρ=-0.16 (p=0.728)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.30 (p=0.402)
  Spearman ρ=-0.08 (p=0.834)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.22 (p=0.561)
  Spearman ρ=-0.14 (p=0.714)