012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_negative_max_area_50C
Pearson r=-0.228 (p=0.217) | Spearman ρ=-0.099 (p=0.596) | Kendall τ=-0.061 (p=0.644)
Pearson log1p(activity) r=-0.120 (p=0.52)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.12, p=0.75)

Global statistics\nn = 31\nR² = 0.052\nRMSE = 86.86 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.46 (p=0.301)
  Spearman ρ=-0.19 (p=0.682)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.12 (p=0.75)
  Spearman ρ=-0.26 (p=0.46)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.17 (p=0.659)
  Spearman ρ=0.12 (p=0.761)