031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_global_score_50C
Pearson r=-0.294 (p=0.109) | Spearman ρ=-0.145 (p=0.438) | Kendall τ=-0.111 (p=0.385)
Pearson log1p(activity) r=-0.126 (p=0.499)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.31, p=0.378)

Global statistics\nn = 31\nR² = 0.086\nRMSE = 85.28 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
Pearson r=-0.24 (p=0.599)
Spearman ρ=0.02 (p=0.969)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.31 (p=0.378)
Spearman ρ=0.26 (p=0.476)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.62 (p=0.0734)
Spearman ρ=0.55 (p=0.121)