031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_negative_max_area_50C
Pearson r=-0.234 (p=0.206) | Spearman ρ=-0.216 (p=0.243) | Kendall τ=-0.151 (p=0.245)
Pearson log1p(activity) r=-0.248 (p=0.179)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.57, p=0.0854)

Global statistics\nn = 31\nR² = 0.055\nRMSE = 86.75 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.66 (p=0.105)
  Spearman ρ=0.72 (p=0.0691)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.57 (p=0.0854)
  Spearman ρ=-0.60 (p=0.0653)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.51 (p=0.156)
  Spearman ρ=-0.44 (p=0.24)