031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_positive_total_area_50C
Pearson r=0.509 (p=0.00343) | Spearman ρ=0.416 (p=0.0201) | Kendall τ=0.300 (p=0.0195)
Pearson log1p(activity) r=0.461 (p=0.00912)




Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.10, p=0.78)

Global statistics\nn = 31\nR² = 0.259\nRMSE = 76.78 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.19 (p=0.676)
  Spearman ρ=0.11 (p=0.818)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.10 (p=0.78)
  Spearman ρ=-0.02 (p=0.947)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.31 (p=0.417)
  Spearman ρ=0.19 (p=0.625)