050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_positive_max_area_50C
Pearson r=-0.065 (p=0.726) | Spearman ρ=0.076 (p=0.683) | Kendall τ=0.038 (p=0.771)
Pearson log1p(activity) r=0.041 (p=0.825)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.49, p=0.15)

Global statistics\nn = 31\nR² = 0.004\nRMSE = 89.02 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.03 (p=0.953)
  Spearman ρ=0.20 (p=0.67)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.49 (p=0.15)
  Spearman ρ=0.45 (p=0.193)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.08 (p=0.84)
  Spearman ρ=0.27 (p=0.475)