069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_negative_max_area_50C
Pearson r=-0.135 (p=0.468) | Spearman ρ=0.093 (p=0.62) | Kendall τ=0.060 (p=0.645)
Pearson log1p(activity) r=0.075 (p=0.687)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.15, p=0.68)

Global statistics\nn = 31\nR² = 0.018\nRMSE = 88.39 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.45 (p=0.316)
  Spearman ρ=-0.50 (p=0.253)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.15 (p=0.68)
  Spearman ρ=0.26 (p=0.466)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.18 (p=0.65)
  Spearman ρ=0.33 (p=0.389)