089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_negative_max_area_50C
Pearson r=-0.190 (p=0.305) | Spearman ρ=-0.174 (p=0.348) | Kendall τ=-0.108 (p=0.411)
Pearson log1p(activity) r=-0.237 (p=0.199)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.17, p=0.638)

Global statistics\nn = 31\nR² = 0.036\nRMSE = 87.59 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.07 (p=0.883)
  Spearman ρ=0.10 (p=0.83)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.17 (p=0.638)
  Spearman ρ=0.17 (p=0.63)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.38 (p=0.311)
  Spearman ρ=-0.18 (p=0.635)