089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_positive_total_area_50C
Pearson r=0.474 (p=0.00704) | Spearman ρ=0.413 (p=0.021) | Kendall τ=0.315 (p=0.0141)
Pearson log1p(activity) r=0.419 (p=0.0189)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.32, p=0.37)

Global statistics\nn = 31\nR² = 0.225\nRMSE = 78.55 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.60 (p=0.156)
  Spearman ρ=0.59 (p=0.159)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.32 (p=0.37)
  Spearman ρ=-0.32 (p=0.374)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.45 (p=0.224)
  Spearman ρ=0.45 (p=0.22)