127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_positive_max_area_50C
Pearson r=-0.138 (p=0.459) | Spearman ρ=0.054 (p=0.772) | Kendall τ=0.031 (p=0.811)
Pearson log1p(activity) r=0.017 (p=0.926)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.56, p=0.0893)

Global statistics\nn = 31\nR² = 0.019\nRMSE = 88.36 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.41 (p=0.355)
  Spearman ρ=-0.07 (p=0.877)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.56 (p=0.0893)
  Spearman ρ=0.46 (p=0.181)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.01 (p=0.985)
  Spearman ρ=0.28 (p=0.461)