146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_negative_max_area_50C
Pearson r=-0.225 (p=0.225) | Spearman ρ=-0.096 (p=0.608) | Kendall τ=-0.056 (p=0.669)
Pearson log1p(activity) r=-0.209 (p=0.26)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.27, p=0.456)

Global statistics\nn = 31\nR² = 0.050\nRMSE = 86.94 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.51 (p=0.241)
  Spearman ρ=0.39 (p=0.386)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.27 (p=0.456)
  Spearman ρ=0.59 (p=0.071)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.38 (p=0.318)
  Spearman ρ=-0.40 (p=0.282)