146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_positive_max_area_50C
Pearson r=-0.018 (p=0.924) | Spearman ρ=-0.065 (p=0.73) | Kendall τ=-0.047 (p=0.72)
Pearson log1p(activity) r=0.006 (p=0.973)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.25, p=0.482)

Global statistics\nn = 31\nR² = 0.000\nRMSE = 89.20 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.23 (p=0.617)
  Spearman ρ=0.20 (p=0.667)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.25 (p=0.482)
  Spearman ρ=-0.15 (p=0.674)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.19 (p=0.62)
  Spearman ρ=-0.17 (p=0.666)