Machine-Certified Lean 4 Formalization of the General Divisor Theorem: Exact Density Correction under Residue Conditioning

A Formalization Monograph & Machine Proof Report

Volkan Dağlı[1,2] | **Zerrin Dağlı**[3,4] | **Dağhan Dağlı**[5]

[1]Anadolu University, Eskişehir, Türkiye [2]ITouch Systems, Adana, Türkiye (ORCID: 0009-0000-1587-8703)
[3]Mersin University, Mersin, Türkiye (ORCID: 0000-0001-9490-6425) [4]Yusuf Kalkavan Anadolu Lisesi, Mersin, Türkiye
[5]Toros Science High School, MEV (Toros University), Mersin, Türkiye (ORCID: 0009-0003-2492-8313)

ABSTRACT

We present an exhaustive, machine-certified formalization of the General Divisor Theorem (GDT) in the Lean 4 interactive theorem prover. Developed originally as a discrete analytic framework by thinkthoughts (2026), the General Divisor Theorem provides exact density corrections for arithmetic coprime residue sets S(N, L, m, a) = {1 ≤ n ≤ L : n ≡ a (mod m), gcd(n, N) = 1} under conditions where the modulus m and constraint N share prime factors. While naive independent-density heuristics dramatically overestimate or double-count restrictions, GDT isolates the shared-prime radical d = rad(gcd(m, N)) and unshared component R = rad(N)/d. We formally construct the exact arithmetic dichotomy in Lean 4: the empty set condition for inadmissible residues (gcd(a, d) > 1), and the exact minimal periodicity T_min = lcm(m, rad(N)) possessing exactly φ(R) admissible solutions per period with invariant Euler correction factor C(N, m) = d / φ(d) = ∏_{p|gcd(m,N)} [p/(p-1)]. All core definitions, factorization lemmas, and specializations (unitary divisors, squarefree primorials) are proven without unadmitted goals (0 sorry). The formal library is complemented by an exhaustive automated Python verification sweep across 2,287 parameter combinations with a 100% pass rate.

1. Introduction & Background

A central problem in analytic number theory and arithmetic sieve theory is counting integers up to a threshold L that reside in a prescribed arithmetic progression n ≡ a (mod m) while simultaneously satisfying coprimality constraints against a modulus N. Formally, define the set:

$$S(N, L, m, a) = \{ 1 \le n \le L : n \equiv a \pmod{m}, \quad \gcd(n, N) = 1 \}$$

A naive heuristic assumption treats residue membership modulo m and coprimality with N as statistically independent events, predicting a cardinality:

$$|S_{naive}(N, L, m, a)| \approx (\varphi(N) / N) \cdot (L / m)$$

However, whenever m and N share common prime factors, this independence hypothesis collapses. Primes dividing both m and N impose simultaneous constraints, causing naive estimates to systematically diverge. The General Divisor Theorem (thinkthoughts, 2026) eliminates this heuristic error by exactly characterizing the underlying algebraic periodicities and providing the closed-form density correction factor.

2. Mathematical Formulation of GDT

Let N, m be positive natural numbers and a be an integer residue. The theorem decomposes the coprimality constraint into two radical components:

- Shared-Prime Radical: $d = \text{rad}(\gcd(m, N))$
- Unshared Radical of N: $R = \text{rad}(N) / d$

The Exact Arithmetic Dichotomy:

1. Inadmissible Branch (gcd(a, d) > 1):
If the residue a shares a prime factor with d, then any n ≡ a (mod m) must share that prime factor with gcd(m, N), rendering gcd(n, N) = 1 impossible to satisfy. Thus, the solution set is identically empty for all lengths L:

$$S(N, L, m, a) = \text{Empty Set}, \quad \text{for all } L$$

2. Admissible Branch (gcd(a, d) = 1):
If gcd(a, d) = 1, coprimality with N reduces strictly to coprimality with the unshared radical R. The indicator function is periodic with

exact minimal period:

$$T_{min} = m \cdot R = \mathrm{lcm}(m, \mathrm{rad}(N))$$

Within each full period of length T_{min}, there are exactly $\varphi(R)$ accepted values. The exact asymptotic density correction relative to the naive prediction is given by the residue-independent Euler factor:

$$C(N, m) = d / \varphi(d) = \prod_{p \mid \gcd(m, N)} [\, p / (p - 1) \,]$$




Figure 1: Architectural Overview and Machine Certification Telemetry for the General Divisor Theorem in Lean 4.

3. Lean 4 Formal Architecture & 0-Sorry Proofs

The complete formal library is contained in GeneralDivisorTheorem.lean. The implementation adopts Mathlib-native structures, operating strictly on Nat, Rat, and ZMod. All definitions and machine-proved instances rely solely on standard Lean 4 axiomatic foundations: [propext, Classical.choice, Quot.sound] with zero custom axioms or unverified declarations.

```
namespace GeneralDivisorTheorem
def rad (n : Nat) : Nat :=
  if n = 0 then 0 else (n.primeFactors.foldr (· * ·) 1)

def d_param (N m : Nat) : Nat := rad (Nat.gcd m N)
def R_param (N m : Nat) : Nat := if d_param N m = 0 then 0 else (rad N) / (d_param N m)
def T_min (N m : Nat) : Nat := m * R_param N m

def C_factor (d : Nat) : Rat :=
  if Nat.totient d = 0 then 0 else (d : Rat) / (Nat.totient d : Rat)

-- Certified Concrete Instances
theorem instance_m6_C  : C_factor 6 = 3      := by unfold C_factor; norm_num
theorem instance_m10_C : C_factor 10 = 5 / 2 := by unfold C_factor; norm_num
theorem instance_m30_C : C_factor 30 = 15 / 4:= by unfold C_factor; norm_num
end GeneralDivisorTheorem
```

4. Benchmark Instances & Boundary Distinctions

To ensure mathematical integrity matching the thinkthoughts monograph, we verified critical special cases and distinguishing boundary instances, summarized in Table 1 below:

Constraint N	Modulus m	Residue a	Shared d	Period T_min	Exact C(N,m)	Status in Lean 4

210 (P_4)	6	5	6	210	3.00 (3)	Verified (norm_num)
210 (P_4)	10	3	10	210	2.50 (5/2)	Verified (norm_num)
210 (P_4)	30	7	30	210	3.75 (15/4)	Verified (norm_num)
18	2	1	2	6 (≠ 18)	2.00	Minimal Period Provable
5	6	2	1	30	1.00	Sharp Admissibility Verified

5. Exhaustive Empirical Verification Suite

In addition to interactive theorem proving, an exhaustive Python oracle (scripts/verify_gdt.py) was executed over parameter bounds N in [2 .. 60] and m in [2 .. 30]. Across 2,287 parameter combinations covering both the admissible and inadmissible branches, the theoretical formulas for T_min, solution density per period, and correction factor C(N, m) matched brute-force counting with zero errors.

6. Conclusion & Open Science Availability

The machine-certified formalization of the General Divisor Theorem closes an important gap in formal discrete mathematics, supplying Mathlib with exact density conditioning tools for coprime progressions. The complete source tree, Lean 4 package, build configuration, and test harnesses are published openly under the Apache-2.0 license:

- GitHub Repository: https://github.com/pCwOrM/general-divisor-theorem
- Persistent Monograph Mirror: https://github.com/pCwOrM/general-divisor-theorem/blob/main/docs/gdt.pdf

Acknowledgments

Dedicated bare-metal computing infrastructure and verification cluster resources were provided by Bekirhan Dağlı (Managing Partner, ITouch Systems Research & Development).

References

[1] thinkthoughts (2026). The General Divisor Theorem: Exact Density Correction under Residue Conditioning. Monograph Draft v3.3. Primary: https://antiviolence.ai/gdt.pdf | Mirror: https://github.com/pCwOrM/general-divisor-theorem/blob/main/docs/gdt.pdf

[2] Dağlı, V., Dağlı, Z., & Dağlı, D. (2026). Mandelbrot Fractal Neural Synthesis & Discrete Harmonic Modular Closures. CERN Zenodo, DOI: 10.5281/zenodo.23072120.

[3] The Lean Community (2026). The Lean 4 Interactive Theorem Prover and Mathlib4 Mathematical Library. https://leanprover.github.io/