Automated Reasoning and In-Worker Verification for High-Rank Elliptic Curves: The 4-Stage Werr Algebraic Sieve, Conductor Certification, and Isogeny Graph Transport on the NSF ICARM Leaderboard

Volkan Dağlı[1,2] **Zerrin Dağlı**[3,4] **Dağhan Dağlı**[5]

[1]Anadolu University, Eskişehir, Türkiye
[2]ITouch Systems Research & Development, Adana, Türkiye
[3]Mersin University, Mersin, Türkiye
[4]Yusuf Kalkavan Anadolu Lisesi, Mersin, Türkiye
[5]Toros Science High School, MEV (Toros University), Mersin, Türkiye

ORCID: 0009-0000-1587-8703 · 0000-0001-9490-6425 · 0009-0003-2492-8313
Correspondence: ask@answerr.me

October 2026

Abstract

The computational verification of high-rank elliptic curves over $\mathbb{Q}$ has historically been bottlenecked by the difficulty of certifying global arithmetic invariants, most notably the arithmetic conductor N. In modern automated reasoning platforms such as the NSF Institute for Computer-Aided Reasoning in Mathematics (ICARM) Elliptic Curve Rank Leaderboard (supported by U.S. National Science Foundation Grant DMS-2425401), candidate curves are subjected to strict client-side and in-worker WebAssembly PARI/GP 2.17.2 kernel verification. While geometric non-singularity, rational point membership, torsion subgroup structures, and Néron–Tate height regulator determinants are verified efficiently, computing the global conductor via Tate's algorithm (`ellglobalred`) requires the complete prime factorization of the minimal discriminant Δ. For state-of-the-art curves specializing high-rank parametric families, $|\Delta|$ routinely spans 50 to 110 decimal digits, leaving many landmark curves across the literature with uncertified conductors (marked as unknown or unranked).

In this paper, we present an automated certificate-generation and isogeny graph exploration framework that completely resolves this bottleneck. Our methodology unites two computational engines derived from our formally verified Lean 4 mathematical corpus: (1) the *4-Stage Werr Multi-Layer Algebraic Sieve*, which couples algebraic invariant-root peeling ($\gcd(a_4, a_6, \Delta, c_4, c_6)$ and 2-torsion monic cubic divisors $\Delta = d_0 \cdot s^3$), primorial radical wheel contraction via the General Divisor Theorem ($T_{\min} = \mathrm{lcm}(m, \mathrm{rad}(N))$), 24-dimensional Niemeier lattice A_8^3 isotropic glue code compression, and 14-modulus orbit-stabilizer quadratic residue coset filtering (161,352.41× candidate reduction prior to `isqrt`) with harmonic multiplier Fermat conditioning, word-packed $\mathbb{F}_2$ MPQS, and carrier-parity ECM; and (2) systematic *Vélu Isogeny Graph Transport*, traversing rational n-isogeny orbits ($n \in \{2, 3, 4, 6\}$) to extract minimal Weierstrass models with strictly smaller discriminants, Faltings heights, and naive heights while preserving Mordell–Weil rank and conductor.

Deploying this pipeline on the NSF ICARM leaderboard, we established **230 newly certified curves** (`#4308`–`#4537`) with **100% certified conductor factorizations** (230/230, eliminating all missing conductor certificates), securing **30 #1 Gold records**, **80 Top-3 podium finishes**, and **418 Top-10 placements** across 43 active leaderboards. Landmark results include the "Grand Slam" curve `#4315` ($T = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}, \mathrm{Rank} \geq 9$), which captures first place in all four competition metrics simultaneously, the complete factorization of a 110-digit discriminant into 21 distinct bad primes for the Rank ≥ 12 curve `#4383`, and full conductor certification for the Elkies Rank ≥ 19 curve `#4323`. All datasets, source scripts, and verification receipts are permanently archived at CERN Zenodo (DOI: 10.5281/zenodo.23436282).

Keywords: Automated Reasoning, Proof Certificates, Elliptic Curves, Conductor Certification, Tate's Algorithm, Vélu Isogenies, Algebraic Sieve, Symbolic Computation, Computer-Aided Mathematics.

2020 Mathematics Subject Classification: 11G05 (Primary); 11Y05, 11Y16, 14H52, 68T15, 68W30.

ACM Computing Classification (CCS 2012): Theory of computation → Logic and verification; Automated reasoning; Computing methodologies → Symbolic and algebraic manipulation.

1 Introduction and Computer-Aided Reasoning in Arithmetic Geometry

The interface between automated deduction, formal proof verification, and computational number theory has undergone a structural transformation over the past decade [1, 2, 10]. Historically, computational records in arithmetic geometry—such as tables of elliptic curves over $\mathbb{Q}$ with high Mordell–Weil rank or prescribed torsion subgroups—were compiled as static text repositories maintained by individual researchers. While invaluable to the mathematical community, static compendia rely on implicit trust in unbundled off-line scripts and often omit computationally expensive secondary invariants.

A major step toward interactive, machine-certified mathematical benchmarking is the **NSF Institute for Computer-Aided Reasoning in Mathematics (ICARM)**, established under U.S. National Science Foundation Grant DMS-2425401. As part of its automated reasoning infrastructure, ICARM operates the official **Elliptic Curve Rank Leaderboard** (`https://elliptic-rank.icarm.cloud`), architected and maintained by David Renshaw and Seewoo Lee [11]. Rather than accepting unverified user claims, the ICARM platform enforces a zero-trust, certificate-driven verification protocol executed inside an isolated WebAssembly (WASM) worker running the computer algebra kernel **PARI/GP 2.17.2** [13].

1.1 The In-Worker WebAssembly Verification Protocol

When a researcher submits a candidate elliptic curve $E/\mathbb{Q}$ specified by integer Weierstrass coefficients $\mathbf{a} = [a_1, a_2, a_3, a_4, a_6]$, a list of rational point coordinates $\mathcal{P} = \{P_1, \dots, P_r\}$, and an optional list of prime factors $\mathcal{S} = \{p_1, \dots, p_k\}$, the ICARM verification kernel executes a six-stage formal decision procedure:

1. **Geometric Non-Singularity Check:** The kernel evaluates the Weierstrass invariants $b_2, b_4, b_6, b_8, c_4, c_6$ and the discriminant $\Delta(\mathbf{a})$. It verifies that $\Delta(\mathbf{a}) \neq 0$, guaranteeing that E defines a smooth projective curve of genus 1 over $\mathbb{Q}$.
2. **Canonical Model Key Generation:** Using LMFDB-compatible invariant reduction, the kernel computes the canonical isomorphism class key $(c_4 : c_6 : \Delta)$ to detect isomorphic duplicate submissions across different Weierstrass coordinates.
3. **Exact Rational Point Membership:** For every asserted generator $P_j = (x_j, y_j) \in \mathbb{Q}^2$, the kernel verifies exact equality in $\mathbb{Q}$:
$$y_j^2 + a_1 x_j y_j + a_3 y_j = x_j^3 + a_2 x_j^2 + a_4 x_j + a_6. \tag{1}$$
4. **Torsion Subgroup Certification:** Applying Lutz–Nagell bounds and division polynomial root analysis via `elltors(E)`, the kernel determines the exact rational torsion subgroup $E(\mathbb{Q})_{\text{tors}} \cong T$ among the 15 admissible groups classified by Mazur's Torsion Theorem [12].
5. **Mordell–Weil Rank Certification via Regulator Positivity:** The kernel computes the canonical Néron–Tate height pairing matrix
$$R(\mathcal{P}) = (\langle P_i, P_j \rangle)_{1 \le i,j \le r}, \quad \langle P_i, P_j \rangle = \hat{h}(P_i + P_j) - \hat{h}(P_i) - \hat{h}(P_j), \tag{2}$$
using `ellbil(E, P_i, P_j)` at high precision, and extracts the maximum $\mathbb{Z}$-linearly independent subset of $\mathcal{P}$ whose Gram determinant $\det R > 0$ is strictly bounded away from zero. This provides a machine-certified lower bound $\operatorname{rank}(E(\mathbb{Q})) \ge r$.
6. **Global Minimal Reduction and Conductor Certification ($\log N$):** Finally, the kernel invokes Tate's global reduction algorithm [16] via `ellglobalred(E)` to transform E to its global minimal integral Weierstrass equation, compute the minimal discriminant $|\Delta_{\min}|$, the Faltings height $h_F(E)$, the naive height $h_{\text{naive}}(E)$, and the global arithmetic conductor N.

1.2 The Conductor Bottleneck on High-Rank Leaderboards

Across the ICARM leaderboard, curves within each (T, r) category—where $T = E(\mathbb{Q})_{\text{tors}}$ is the torsion subgroup and r is the certified Mordell–Weil rank lower bound—are ranked across four primary minimality metrics:

$$\text{(i) Conductor } N, \quad \text{(ii) Minimal Discriminant } |\Delta|, \quad \text{(iii) Faltings Height } h_F(E), \quad \text{(iv) Naive Height } h_{\text{naive}}(E). \tag{3}$$

While $|\Delta|$, $h_F(E)$, and $h_{\text{naive}}(E)$ can be evaluated directly from the Weierstrass coefficients once small local scaling factors at $p \in \{2, 3\}$ are removed, computing the conductor $N = \prod_{p|\Delta} p^{f_p}$ requires knowing the *complete prime factorization* of the discriminant Δ.

In high-rank elliptic curve constructions pioneered by Noam Elkies [9], Andrej Dujella [8], Michael Rubinstein [14], and Zev Klagsbrun, curves of rank $r \in [7, 29]$ are obtained by specializing multi-parameter elliptic surfaces over $\mathbb{Q}(t_1, \ldots, t_m)$ at rational parameters of large height. Because the discriminant Δ is a polynomial of high degree in the specialization parameters, the resulting integer $|\Delta|$ routinely spans **50 to 110 decimal digits** (165 to 365 bits). When submitted without a pre-computed prime factorization certificate $\mathcal{S}$, the in-worker PARI/GP kernel cannot factor $|\Delta|$ within the interactive execution budget, leaving the conductor field marked as `?` (unfactored/unknown) and disqualifying the curve from the Conductor N leaderboard.

Furthermore, many high-rank curves in the historical literature are recorded in a single Weierstrass coordinate system tied to their fibration parameterization, without exploring the finite graph of $\mathbb{Q}$-isogenous models. Because rational isogenies preserve both the Mordell–Weil rank r and the conductor N while altering $|\Delta|$, $h_F(E)$, and $h_{\text{naive}}(E)$, uninspected vertices of the rational isogeny graph often conceal strictly superior minimal models.

1.3 Summary of Contributions

In this work, we bridge automated proof-certificate generation and computational arithmetic geometry through three primary contributions:

1. **Formalization of Conductor Certification via the 4-Stage Werr Algebraic Sieve:** We synthesize four formally verified mathematical pillars from our Lean 4 research corpus—the General Divisor Theorem (GDT v3.0) [4], the 24-dimensional Niemeier lattice A_8^3 glue code [5], GAP Phase 1–2 orbit-stabilizer coset enumeration [6], and Schönhage bipartite tensor/word-packed $\mathbb{F}_2$ algebra [7]—with algebraic invariant-root peeling to factor all residual composite discriminants up to 110 decimal digits.
2. **Systematic Vélu Isogeny Graph Transport Engine:** We implement an exact rational isogeny graph explorer covering cyclic and non-cyclic subgroups of orders $n \in \{2, 3, 4, 5, 6, 7, 8, 9, 10, 12\}$, transporting rational Mordell–Weil generators $P_j \mapsto \phi(P_j)$ and performing u-scaling to discover global minimal models with substantially reduced $|\Delta|$, h_F, and h_{naive}.
3. **Large-Scale Public Verification on the NSF ICARM Leaderboard:** We submitted and certified **230 distinct Top-10 curves** (`#4308` through `#4537`) under contributor profile `user/114` (https://elliptic-rank.icarm.cloud/user/114), achieving **100.0% conductor certification** (230/230 with zero unknown conductors), **30 #1 Gold records**, **80 Top-3 podiums**, and **418 Top-10 metric placements** across 43 active leaderboards.

2 Conductor Determination as Formal Proof Certificate Checking

Let $E/\mathbb{Q}$ be an elliptic curve defined by a generalized Weierstrass equation with integer coefficients:

$$E: \quad y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6, \quad a_1, a_2, a_3, a_4, a_6 \in \mathbb{Z}. \tag{4}$$

The standard Tate quantities b_2, b_4, b_6, b_8 and covariants c_4, c_6, Δ are given by:

$$b_2 = a_1^2 + 4a_2, \qquad b_4 = 2a_4 + a_1a_3, \qquad b_6 = a_3^2 + 4a_6, \tag{5}$$

$$b_8 = a_1^2a_6 + 4a_2a_6 - a_1a_3a_4 + a_2a_3^2 - a_4^2, \tag{6}$$

$$c_4 = b_2^2 - 24b_4, \qquad c_6 = -b_2^3 + 36b_2b_4 - 216b_6, \tag{7}$$

$$\Delta = -b_2^2b_8 - 8b_4^3 - 27b_6^2 + 9b_2b_4b_6 = \frac{c_4^3 - c_6^2}{1728}. \tag{8}$$

An admissible change of Weierstrass coordinates over $\mathbb{Q}$ takes the form $(x, y) = (u^2x' + r, u^3y' + u^2sx' + t)$ with $u \in \mathbb{Q}^\times$ and $r, s, t \in \mathbb{Q}$, under which the invariants transform as:

$$c_4' = u^{-4}c_4, \qquad c_6' = u^{-6}c_6, \qquad \Delta' = u^{-12}\Delta. \tag{9}$$

An integral model (4) is *globally minimal* over $\mathbb{Z}$ if $v_p(\Delta)$ is minimized simultaneously at every prime p. Because $\mathbb{Z}$ has class number 1, every elliptic curve $E/\mathbb{Q}$ admits a unique normalized global minimal model satisfying $a_1, a_3 \in \{0, 1\}$ and $a_2 \in \{-1, 0, 1\}$.

Definition 2.1 (Arithmetic Conductor and Ogg–Saito Formula)**.** Let $E/\mathbb{Q}$ be a globally minimal Weierstrass model with minimal discriminant $\Delta_{\min}$. The *arithmetic conductor* of E is the positive integer

$$N(E) = \prod_{p \mid \Delta_{\min}} p^{f_p}, \tag{10}$$

where the local exponent $f_p \in \mathbb{Z}_{\geq 0}$ is given by the Ogg–Saito formula:

$$f_p = v_p(\Delta_{\min}) - m_p + 1, \tag{11}$$

with m_p denoting the number of irreducible components (counted without multiplicity) of the special fiber $\mathcal{E}_p$ of the Néron minimal model of E over $\mathbb{Z}_p$. Specifically:

- $f_p = 0$ if and only if $p \nmid \Delta_{\min}$ (E has good reduction at p, Kodaira type I_0);
- $f_p = 1$ if and only if $p \parallel c_4$ fails and E has multiplicative reduction at p (Kodaira type I_n, $n = v_p(\Delta_{\min}) \geq 1$);
- $f_p = 2 + \delta_p \geq 2$ if E has additive (potentially good or potentially multiplicative) reduction at p, where the wild Swan conductor $\delta_p = 0$ for all primes $p \geq 5$, while $0 \leq \delta_3 \leq 3$ (so $f_3 \leq 5$) and $0 \leq \delta_2 \leq 6$ (so $f_2 \leq 8$).

Proposition 2.2 (Certificate Verification Complexity of Tate's Algorithm)**.** *Let $E/\mathbb{Q}$ be specified by an integral Weierstrass tuple $\mathbf{a} = [a_1, a_2, a_3, a_4, a_6] \in \mathbb{Z}^5$ with discriminant $\Delta \neq 0$. Suppose an external prover supplies a prime factorization certificate*

$$\mathcal{S} = \{p_1, p_2, \ldots, p_k\} \quad \textit{such that} \quad |\Delta| = \prod_{i=1}^{k} p_i^{e_i}, \quad e_i = v_{p_i}(\Delta) \geq 1. \tag{12}$$

Then a verification kernel can certify the primality of each p_i, execute Kraus–Laska–Connell local minimization at every p_i with $e_i \geq 12$, run Tate's algorithm [16] at all bad primes, and output the exact global minimal model $E_{\min}$ and certified conductor $N(E)$ in deterministic time polynomial in $\log |\Delta|$.

Proof. Verification of the identity $|\Delta| = \prod_{i=1}^{k} p_i^{e_i}$ requires at most $\sum_{i=1}^{k} e_i \leq \log_2 |\Delta|$ multi-precision divisions of bit-length $B = \lceil \log_2 |\Delta| \rceil$, costing $O(B^2)$ bit operations. Primality verification of each $p_i \leq |\Delta|$ via deterministic APRCL or ECPP certificates runs in time $O((\log p_i)^{4+\varepsilon})$. For each prime $p_i \in \mathcal{S}$, Tate's algorithm [3,16] executes at most $\lfloor e_i/12 \rfloor + 1$ local loop iterations. Each iteration computes p_i-adic valuations of the invariants and finds roots over $\mathbb{F}_{p_i}$ of auxiliary polynomials of degree at most 3:

$$P_1(T) = T^2 + a_1 T - a_2 \pmod{p_i}, \tag{13}$$

$$P_2(Y) = Y^2 + a_3 Y - a_6 \pmod{p_i}, \tag{14}$$

$$P_3(X) = X^3 + a_{2,1} X^2 + a_{4,2} X + a_{6,3} \pmod{p_i}, \quad a_{j,m} = p_i^{-m} a_j. \tag{15}$$

Root-finding for degree ≤ 3 polynomials over $\mathbb{F}_{p_i}$ requires $O(\log^2 p_i)$ field operations via Cantor–Zassenhaus. Since $k \leq \log_2 |\Delta|$ and $\sum_{i=1}^{k} \log p_i \leq \log |\Delta|$, the entire global reduction and conductor certification procedure terminates in $O(\mathrm{poly}(\log |\Delta|))$ time. □

Remark 2.3 (Automated Reasoning Architecture: Separating Search from Verification). Proposition 2.2 exemplifies the foundational design principle of modern computer-aided reasoning (cf. `cs.LO`): *separating untrusted, computationally intensive certificate search from lightweight, deterministic kernel verification.* In the ICARM architecture, the submission payload includes the field `"primes": ["p_1", ..., "p_k"]`. The off-line prover (our 4-Stage Werr Algebraic Sieve) expends arbitrary bare-metal compute to factor 70–110 digit discriminants; the in-worker PARI/GP 2.17.2 engine then ingests $\mathcal{S}$ via `addprimes(S)`, verifies primality and trial-division completeness against Δ, and certifies $\log N$ in under 2 seconds.

3 The 4-Stage Werr Multi-Layer Algebraic Sieve

To factor massive discriminants $|\Delta|$ up to 110 decimal digits and construct complete prime certificates $\mathcal{S}$ across all candidate curves, we engineered the **4-Stage Werr Multi-Layer Algebraic Sieve**. Rather than treating Δ as an unstructured black-box integer, our pipeline exploits both the *algebraic structure of Weierstrass invariants* and a *four-pillar formally verified pre-sieve hierarchy* developed in our Lean 4 open-science corpus.

3.1 Architectural Lineage: The Four Formally Verified Pillars

Every mathematical layer of the pre-sieve and solver pipeline is grounded in an independently machine-certified Lean 4 repository (`0 axioms`, `0 sorry`). In accordance with strict scientific transparency, Table 1 details the exact provenance and quantitative contribution of each pillar.

Table 1: Formal Verification Lineage of the 4-Pillar Werr Algebraic Sieve Architecture

Pillar	Archival Deposit (DOI)	Lean 4 Status	Algebraic Mechanism & Sieve Contribution
Pillar I (Layer 1)	`general-divisor-theorem` 10.5281/zenodo.23425590 [4]	81 Theorems (`0 sorry`) *Spec:* D. Hawkley	**Radical Wheel Contraction** (5.85×)**:** Contracts primorial wheel periods via $T_{\min} = \mathrm{lcm}(m, \mathrm{rad}(N))$, prunes inadmissible progressions ($\gcd(\|a\|, d) > 1 \Rightarrow \emptyset$), and counts survivors in $O(1)$ blocks ($C(qT_{\min} + s) = q\varphi(R) + C(s)$).
Pillar II (Layer 2)	`niemeier-a8-moonshine` 10.5281/zenodo.23324067 [5]	Certified (`0 sorry`)	**Isotropic Glue Compression** (11.69× **Cumul.):** Projects candidate triples onto the 24D Niemeier lattice A_8^3 glue code over $(\mathbb{Z}/9\mathbb{Z})^3$ generated by $g = (1, 1, 4)$ with $\|g\|^2 = 4 \in 2\mathbb{Z}$, compressing $729 \to 27$ states (27.0×) and halving survivors via $(\mathbb{Z}/9\mathbb{Z})^\times$ unit pairing.
Pillar III (Layers 3–4)	`gap-lean4-port` (v0.2.0) 10.5281/zenodo.23045504 [6]	35 Theorems (`0 sorry`) Phases 1 & 2	**Orbit-Stabilizer QR Super-Wheel** (161,352.41×)**:** Applies certified Sims stabilizer chains (`GAP-0190`) and orbit-stabilizer coset enumeration (`GAP-0247`) across 14 coprime moduli, leaving **62** survivors per 10^7 window (99.99938% pruned before `isqrt`).
Pillar IV (Solver)	`integer-mult-bounds` 10.5281/zenodo.23268580 [7]	Certified (`0 sorry`)	**Word-Packed $\mathbb{F}_2$ & Carrier Parity Cascade:** 64-bit word-packed MSB-pivot $\mathbb{F}_2$ Gaussian elimination for MPQS coupled with Schönhage bipartite alternating carrier parity flips (`-377,890` wires) in parallel batched Pollard–Brent ρ and ECM.

3.2 Stage 1: Algebraic Invariant-Root Peeling and Cubic Divisors

Before invoking any general-purpose factorization routine on Δ, Stage 1 exploits algebraic gcd relations among the Weierstrass invariants and division polynomials:

1. **Covariant GCD Cascade:** From the syzygy $1728\Delta = c_4^3 - c_6^2$, any prime power p^e dividing $\gcd(c_4, c_6)$ automatically divides Δ with multiplicity at least $2e$. More generally, Stage 1 computes
$$g_1 = \gcd(|c_4|, |c_6|), \quad g_2 = \gcd(|a_4|, |a_6|), \quad g_3 = \gcd(|b_2|, |b_4|, |b_6|), \quad g_4 = \gcd(|\Delta|, |c_4 \pm c_6|), \tag{16}$$
splitting Δ into smaller algebraic factors prior to sieving.
2. **Monic Cubic Peeling for 2-Torsion Curves:** When $E(\mathbb{Q})[2] \neq \{O\}$, the 2-division polynomial factors over $\mathbb{Q}$, decomposing Δ into lower-degree algebraic components:

Proposition 3.1 (Algebraic Discriminant Decomposition via Rational 2-Torsion)**.** *Let $E/\mathbb{Q}$ be an integral Weierstrass model possessing at least one rational 2-torsion point $P_0 = (x_0, y_0) \in E(\mathbb{Q})[2] \setminus \{O\}$. Then $X_0 = 4x_0 \in \mathbb{Z}$ is an integer root of the monic cubic polynomial*
$$f(X) = X^3 + B_2X^2 + B_4X + B_6 = 0, \quad B_2 = b_2, \quad B_4 = 8b_4, \quad B_6 = 16b_6 \in \mathbb{Z}. \tag{17}$$
Factoring $f(X) = (X - X_0)(X^2 + \alpha X + \beta)$ with $\alpha = B_2 + X_0 \in \mathbb{Z}$ and $\beta = B_4 + \alpha X_0 \in \mathbb{Z}$ decomposes the elliptic discriminant $\Delta(E)$ algebraically into the product:
$$\Delta(E) = 2^{-12} \cdot (\alpha^2 - 4\beta) \cdot (3X_0^2 + 2B_2X_0 + B_4)^2. \tag{18}$$
In particular, if $E(\mathbb{Q})[2] \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ with three integer roots X_0, X_1, X_2 of $f(X)$, then $\Delta(E)$ splits completely into the square of the pairwise root differences:
$$\Delta(E) = 2^{-12} \cdot (X_0 - X_1)^2(X_0 - X_2)^2(X_1 - X_2)^2. \tag{19}$$

Proof. A point $P_0 = (x_0, y_0)$ satisfies $2P_0 = O$ if and only if $2y_0 + a_1x_0 + a_3 = 0$. Completing the square in the Weierstrass equation (4) yields $(2y + a_1x + a_3)^2 = 4x^3 + b_2x^2 + 2b_4x + b_6$. Multiplying by 16 and substituting $X = 4x$ produces the monic integer polynomial $f(X) = X^3 + B_2X^2 + B_4X + B_6$. By Gauss's Lemma, any rational root of $f(X)$ is an integer $X_0 \in \mathbb{Z}$. The discriminant of $f(X)$ is $\mathrm{Disc}(f) = 2^{12}\Delta(E)$. For a factored polynomial $f(X) = (X - X_0)q(X)$, the resultant identity $\mathrm{Disc}(f) = \mathrm{Disc}(q) \cdot q(X_0)^2 = (\alpha^2 - 4\beta)(f'(X_0))^2$ yields (18) and (19). □

Crucially, finding the integer roots $X_0 \in \mathbb{Z}$ of $f(X)$ requires *zero integer factorization*: since $f(X)$ is a cubic with at most two local extrema, Stage 1 computes the exact bit-length root bound $M = 2^{\max(\mathrm{bits}(B_2), \mathrm{bits}(B_4)/2, \mathrm{bits}(B_6)/3)+4}$ and locates all integer roots in $O(\log |B_6|)$ arithmetic operations via exact integer bisection/Newton search. For curves with torsion $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ (such as the 110-digit discriminant curve `#4383`), identity (19) immediately reduces the problem of factoring a 110-digit integer $|\Delta|$ to factoring three integers $|X_i - X_j|$ of at most 18–20 digits each!

3.3 Stage 2: Coprime Wheel Contraction and 14-Modulus QR Super-Wheel

After peeling small primes up to 10^6 and algebraic root factors from Stage 1, any remaining composite cofactor C enters Stage 2. In congruence-of-squares sieving ($a^2 - kC = b^2$), the computational bottleneck is evaluating the multi-precision integer square root $\texttt{isqrt}(a^2 - kC)$. By composing Pillars I, II, and III, non-square candidates are eliminated via single-cycle 64-bit register bitmasks before `isqrt` is ever invoked:

1. **Level L1 (Pillar I — GDT v3.0 Radical Contraction):** Using the foundational General Divisor Theorem specification of Hawkley and its Lean 4 formalization [4], the primorial modulus $M_8 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17 \cdot 19 = 9{,}699{,}690$ is contracted via $T_{\min} = \mathrm{lcm}(m, \mathrm{rad}(C))$. Inadmissible residue progressions satisfying $\gcd(\|a\|, d) > 1$ are pruned identically, and exact survivor offsets are stepped in $O(1)$ blocks via Theorem 18 ($C(qT_{\min} + s) = q\varphi(R) + C(s)$), eliminating 82.89766% of candidates (5.85× speedup).
2. **Level L2 (Pillar II — Niemeier A_8^3 Glue Compression):** Residue triples mod 9 are mapped into the ternary glue code $G_{\text{glue}} \subset (A_8^*/A_8)^3 \cong (\mathbb{Z}/9\mathbb{Z})^3$ of the 24-dimensional Niemeier lattice $N(A_8^3)$ [5]. Compression to the 27 isotropic glue words coupled with $(\mathbb{Z}/9\mathbb{Z})^\times$ unit pairing halves the surviving candidate stream (11.69× cumulative speedup).
3. **Levels L3 & L4 (Pillar III — 14-Modulus Orbit-Stabilizer QR Super-Wheel):** Applying our Lean 4 formalization of Sims stabilizer chains (`GAP-0190`) and orbit-stabilizer coset enumeration (`GAP-0247`) [6], we construct a 64-bit bitmask lookup table over 14 mutually coprime moduli:

$$\mathcal{M}_{14} = \{64, 27, 25, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43\}. \tag{20}$$

For each $m \in \mathcal{M}_{14}$, a 64-bit word $W_m \in \mathbb{Z}/2^{64}\mathbb{Z}$ encodes the admissible cosets $a \bmod m$ for which $(a^2 - kC) \bmod m$ is a quadratic residue in $\mathbb{Z}/m\mathbb{Z}$. A candidate a survives if and only if

$$\bigwedge_{m \in \mathcal{M}_{14}} ((W_m \gg (a \bmod m)) \,\&\, 1) = 1. \tag{21}$$

Table 2 records the exact ablation profile across a contiguous window of $W = 10{,}000{,}000$ candidates above $\lfloor\sqrt{\text{RSA-270}}\rfloor$. Layering Pillars I–III reduces 10^7 candidates to just **62** `isqrt` evaluations, achieving a verified 161,352.41× **reduction** (99.99938% pruned).

Table 2: 5-Level Pre-Sieve Ablation on an 895-Bit Window ($W = 10{,}000{,}000$ Candidates)

Level	Pillar & Sieve Layer Mechanism	Survivors	Pruned (%)	Cumul. Speedup	Marginal Leap
L0	Baseline Parity Sieve (mod 2)	5,000,000	50.00000%	2.00×	2.00×
L1	Pillar I: GDT v3.0 Radical Contraction (M_8)	1,710,234	82.89766%	5.85×	+2.92×
L2	Pillar II: Niemeier A_8^3 Isotropic Glue Code	855,117	91.44883%	11.69×	+2.00×
L3	Pillar III: 11-Modulus QR Coset Quotient ($p \le 31$)	760	99.99240%	13,157.89×	+1,125.15×
L4	Pillar III: 14-Modulus 64-Bit QR Super-Wheel	**62**	**99.99938%**	**161,352.41×**	+12.26×

3.4 Stage 3: Multi-Regime Solver Cascade and Benchmark Validation

Cofactors surviving pre-sieving are dispatched to a three-regime solver cascade powered by Pillar IV [7]:

- **Regime A (`VolkanBridge-Harmonic-Fermat`):** For near-square or ratio-correlated cofactors $C = p \cdot q$, classical Fermat ($k = 1$) stalls whenever $|p - q|$ is large. By conditioning a harmonic multiplier $k \in \{1, 2, \ldots, K_{\max}\}$ in $a^2 - 4kC = b^2$ gated by the 14-modulus QR Super-Wheel, the effective distance collapses to $|kp - q|$. As shown in Table 3, for a 117-bit asymmetric semiprime with $q \approx 10p$ ($p = 10^{17} + 3, q = 10^{18} + 3$), classical Fermat requires $\sim 4.5 \times 10^{17}$ steps, whereas harmonic conditioning at $k = 10$ collapses $|10p - q| = 27$ and resolves the factors in **0.35 ms** (577 checks).
- **Regime B (`Schonhage-WordPacked-F2-QS`):** For uncorrelated balanced cofactors up to 95 bits, polynomial relations are sieved over the GDT-admissible factor base $\mathcal{B} = \{p \le B_{\max} : (\frac{C}{p}) = 1\}$ using Tonelli–Shanks root lifting. Smooth exponent vectors modulo 2 are packed into 64-bit machine words, and linear dependencies in the $\mathbb{F}_2$-nullspace are extracted via bitwise MSB-pivot Gaussian elimination.
- **Regime C (`Schonhage-Carrier-Brent-Rho` & Parallel ECM):** For large unbalanced cofactors up to 110 decimal digits containing 15–35 digit primes, Stage 3 runs parallel batched Pollard–Brent ρ and Lenstra Elliptic Curve Method (ECM) workers. Upon cycle-horizon stagnation, workers inject Schönhage bipartite alternating carrier parity flips [7] coupled with $(\mathbb{Z}/9\mathbb{Z})^\times$ polynomial shifts.

Table 3: 10/10 Multi-Regime Semiprime Factorization Benchmark (40-Bit to 257-Bit)

Benchmark Target	Bits	Arithmetic Regime	Solver Triggered	Time (ms)	Status
`Semiprime-40bit-Bal`	40	Near-square balanced	`VolkanBridge-Harmonic-Fermat(k=1)`	0.01 ms	**PASS**
`Semiprime-66bit-Bal`	67	Near-square balanced	`VolkanBridge-Harmonic-Fermat(k=1)`	0.01 ms	**PASS**
`Semiprime-80bit-Bal`	80	Near-square balanced	`VolkanBridge-Harmonic-Fermat(k=1)`	0.01 ms	**PASS**
`Semiprime-100bit-Bal`	107	Near-square balanced	`VolkanBridge-Harmonic-Fermat(k=1)`	0.00 ms	**PASS**
`Semiprime-117bit-Asym10x`	117	10 : 1 Asymmetric ($10^{17} \times 10^{18}$)	`VolkanBridge-Harmonic-Fermat(k=10)`	**0.35 ms**	**PASS**
`Semiprime-61bit-Uncorr`	61	Uncorrelated general semiprime	`Schonhage-WordPacked-F2-QS(266 rels)`	**233.62 ms**	**PASS**
`Semiprime-128bit-Bal`	129	128-bit near-square block	`VolkanBridge-Harmonic-Fermat(k=1)`	0.02 ms	**PASS**
`Semiprime-160bit-Bal`	161	160-bit near-square block	`VolkanBridge-Harmonic-Fermat(k=1)`	0.01 ms	**PASS**
`Semiprime-201bit-Bal`	201	201-bit near-square block	`VolkanBridge-Harmonic-Fermat(k=1)`	0.01 ms	**PASS**
`Semiprime-257bit-Asym`	257	Extreme asymmetric (33b × 225b)	`Schonhage-Carrier-Brent-Rho(27902)`	**1,857.36 ms**	**PASS**

3.5 Scope and Complexity Boundary: Pre-Sieve Speedup vs. Asymptotic GNFS

In accordance with rigorous computational number theory standards, we explicitly distinguish between *constant-factor pre-sieve pruning* and *asymptotic sub-exponential complexity*. Combining Pillar I (GDT v3.0), Pillar II (Niemeier A_8^3), and Pillar III (14-modulus QR Super-Wheel) achieves a verified 161,352.41× constant-factor reduction in square-root evaluations, while Stage 1 algebraic root peeling and Stage 3 ECM/MPQS completely resolve the highly structured, multi-factor 50–110 digit discriminants of high-rank elliptic curves. Conversely, for unstructured cryptographic semiprimes of scale RSA-270 (895 bits) where two 447-bit primes $p, q \approx 10^{135}$ are chosen uniformly at random without algebraic root structure or small prime factors, a $1.61 \times 10^5\times$ linear pre-sieve speedup cannot bridge a 10^{135} search interval; such targets remain governed by the General Number Field Sieve (GNFS) with heuristic complexity $L_N[1/3, (64/9)^{1/3}] \approx 2^{108}$ operations.

4 Vélu Rational Isogeny Graph Transport and Minimal Model Search

While the 4-Stage Werr Algebraic Sieve resolves the conductor certification bottleneck ($\log N$), our second engine exploits the geometry of rational isogeny graphs to discover strictly smaller minimal Weierstrass models.

4.1 Invariance of Rank and Conductor Across $\mathbb{Q}$-Isogeny Orbits

Let $E/\mathbb{Q}$ and $E'/\mathbb{Q}$ be elliptic curves connected by a non-constant $\mathbb{Q}$-rational isogeny $\phi : E \to E'$ of degree $n = |\ker \phi|$. Let $\hat{\phi} : E' \to E$ denote the dual isogeny satisfying $\hat{\phi} \circ \phi = [n]_E$ and $\phi \circ \hat{\phi} = [n]_{E'}$.

Theorem 4.1 (Invariance of Rank, Conductor, and L-Function under $\mathbb{Q}$-Isogeny)**.** *Let $\phi : E \to E'$ be a $\mathbb{Q}$-rational isogeny between elliptic curves over $\mathbb{Q}$. Then:*

1. ***Mordell–Weil Rank Invariance:*** *ϕ induces a $\mathbb{Q}$-linear isomorphism $E(\mathbb{Q}) \otimes_{\mathbb{Z}} \mathbb{Q} \xrightarrow{\sim} E'(\mathbb{Q}) \otimes_{\mathbb{Z}} \mathbb{Q}$, so* $\operatorname{rank}(E'(\mathbb{Q})) = \operatorname{rank}(E(\mathbb{Q}))$.
2. ***Conductor and Bad Prime Invariance:*** *By the Néron–Ogg–Shafarevich criterion and Tate's isogeny theorem, the ℓ-adic Galois representations $V_\ell(E) \cong V_\ell(E')$ are isomorphic as $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$-modules. Consequently, E and E' have the exact same set of bad primes $\mathcal{S}$, the same global arithmetic conductor $N(E') = N(E)$, and identical Hasse–Weil L-functions $L(E', s) = L(E, s)$.*

Crucially, however, the *minimal discriminant* $|\Delta(E')|$, the *Faltings height* $h_F(E')$, and the *naive height* $h_{\text{naive}}(E') = \log \max(4|a_4|^3, 27a_6^2)$ are **not** invariant under isogeny! For a prime ℓ-isogeny $\phi : E \to E'$, the minimal discriminants at primes p of multiplicative reduction (I_m) change by powers of ℓ: specifically, a fiber of type I_m can transition to $\mathrm{I}_{m\cdot\ell}$ or $\mathrm{I}_{m/\ell}$, altering $|\Delta_{\min}|$ by $\ell^{\pm 12}$ or $\ell^{\pm 6}$ at multiple primes simultaneously, while the canonical height regulator changes by rational powers of ℓ and the Faltings height shifts via the Parshin–Faltings degree formula.

4.2 Explicit Vélu Quotient Equations and Generator Point Transport

Let $G \subset E(\mathbb{Q})_{\text{tors}}$ be a finite $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$-stable subgroup of order n. Following Vélu [17], we partition the non-zero elements $G \setminus \{O\} = G_2 \sqcup G_+ \sqcup G_-$, where $G_2 = \{Q \in G \setminus \{O\} : 2Q = O\}$ consists of the 2-torsion points in G, and each inverse pair $\{Q, -Q\}$ of order > 2 has one representative in G_+ and one in G_-. Setting

$S = G_2 \sqcup G_+$, for each $Q = (x_Q, y_Q) \in S$ we compute in exact rational arithmetic (`fractions.Fraction`):

$$g_Q^x = 3x_Q^2 + 2a_2x_Q + a_4 - a_1y_Q, \qquad g_Q^y = -2y_Q - a_1x_Q - a_3, \tag{22}$$

$$t_Q = \begin{cases} g_Q^x & \text{if } Q \in G_2, \\ 2g_Q^x - a_1g_Q^y & \text{if } Q \in G_+, \end{cases} \qquad u_Q = (g_Q^y)^2, \qquad w_Q = u_Q + x_Q t_Q. \tag{23}$$

Summing $t = \sum_{Q\in S} t_Q$ and $w = \sum_{Q\in S} w_Q$, the quotient curve $E' = E/G$ is given by the Weierstrass model $\mathbf{A} = [A_1, A_2, A_3, A_4, A_6]$:

$$A_1 = a_1, \quad A_2 = a_2, \quad A_3 = a_3, \quad A_4 = a_4 - 5t, \quad A_6 = a_6 - b_2t - 7w. \tag{24}$$

Simultaneously, for every independent Mordell–Weil generator $P_j = (x_j, y_j) \in \mathcal{P}$ on $E(\mathbb{Q})$, our engine transports P_j across the isogeny via the exact rational map $\phi(P_j) = (X_j, Y_j) \in E'(\mathbb{Q})$:

$$X_j = x_j + \sum_{Q\in G\setminus\{O\}} \left(x(P_j + Q) - x(Q)\right), \qquad Y_j = y_j + \sum_{Q\in G\setminus\{O\}} \left(y(P_j + Q) - y(Q)\right). \tag{25}$$

When the coordinates of $Q \in G \setminus \{O\}$ have rational denominators (for example, $x_Q = X_0/4$ from a root of the monic cubic (17)), the raw Vélu coefficients $A_4, A_6 \in \mathbb{Q}$ may contain powers of 2 or 3 in their denominators. Our engine automatically finds the minimal scaling factor $u \in \mathbb{Q}^\times$ such that

$$\tilde{A}_1 = uA_1, \quad \tilde{A}_2 = u^2A_2, \quad \tilde{A}_3 = u^3A_3, \quad \tilde{A}_4 = u^4A_4, \quad \tilde{A}_6 = u^6A_6 \in \mathbb{Z} \tag{26}$$

forms a globally minimal integral model, and scales every transported generator accordingly via $(\tilde{X}_j, \tilde{Y}_j) = (u^2X_j, u^3Y_j)$.

4.3 Torsion Subgroup Transitions Along Isogeny Graphs

A remarkable feature of Vélu graph exploration is that an isogeny $\phi : E \to E/G$ can either *preserve* the torsion structure $E'(\mathbb{Q})_{\text{tors}} \cong E(\mathbb{Q})_{\text{tors}}$ (improving records within the same leaderboard table) or *transition* between distinct torsion subgroups (populating adjacent leaderboard tables at identical high rank r):

- $\mathbb{Z}/4\mathbb{Z} \longleftrightarrow \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ via quotienting by the unique rational subgroup of order 2;
- $\mathbb{Z}/4\mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ when the dual 2-isogeny pulls back full rational 2-torsion alongside a rational 4-cycle;
- $\mathbb{Z}/2\mathbb{Z} \longleftrightarrow \mathbb{Z}/2\mathbb{Z}$ and $\mathbb{Z}/6\mathbb{Z} \longleftrightarrow \mathbb{Z}/6\mathbb{Z}$ via rational 2-isogenies and 3-isogenies;
- $\mathbb{Z}/3\mathbb{Z} \longrightarrow$ Trivial [] (or $\mathbb{Z}/3\mathbb{Z} \longleftrightarrow \mathbb{Z}/3\mathbb{Z}$ when the kernel of the dual 3-isogeny splits over $\mathbb{Q}$).

5 Empirical Leaderboard Results on NSF ICARM

Executing the combined 4-Stage Werr Algebraic Sieve and Vélu Isogeny Graph Engine across the live ICARM database and historical parametric curve families, we submitted and certified **230 distinct Top-10 curves** (`#4308` through `#4537`) on `https://elliptic-rank.icarm.cloud` under contributor profile **Volkan Dagli** (`user/114`).

5.1 Global Certification Metrics and Torsion–Rank Distribution

Table 4 summarizes the overall audit of our 230 accepted curves verified by the ICARM in-worker PARI/GP 2.17.2 engine, and Table 5 gives the exact distribution of these 230 curves across all 8 torsion subgroups and 13 rank thresholds ($r = 4$ to $r = 19$).

5.2 Complete Breakdown of #1 Gold Records (30 First-Place Placements)

Table 6 presents the flagship curves that achieved `#1` (First Place / Gold Record) across the four official leaderboard metrics: Conductor (N), Minimal Discriminant ($|\Delta|$), Faltings Height (h_F), and Naive Height (h_{naive}).

Table 4: Official Verification and Leaderboard Audit Summary on NSF ICARM (`user/114`)

Leaderboard Metric / Verification Attribute	**Certified Value**
Total Curves Accepted in Top-10 (`#4308`–`#4537`)	**230 curves** (100.0% `status: created`)
Certified Conductor Factorizations ($\log N$)	**230 / 230** (100.0% **factored, 0 unknown ?**)
Active (T, Rank) Leaderboards with Top-10 Presence	**43 distinct leaderboards**
Total **#1 Gold** (First-Place World Record) Placements	**30 First-Place Records**
Total **Top-3** (Podium: #1, #2, #3) Placements	**80 Podium Finishes**
Total **Top-10** Placements Across All 4 Metrics	**418 Top-10 Placements**
Mordell–Weil Rank Range of Certified Curves	**r $\geq$ 4 up to r $\geq$ 19**
Largest Minimal Discriminant Completely Factored	**110 decimal digits** (21 distinct prime factors)

Table 5: Distribution of the 230 Certified Curves Across Torsion Subgroups and Rank Bounds

Torsion T	**ICARM Key**	**r=4**	**5**	**6**	**7**	**8**	**9**	**10**	**11**	**12**	**13**	**14**	**15**	**19**	**Total**
Trivial $\{O\}$	`[]`	–	–	19	19	16	14	2	8	12	12	1	4	1	**108**
$\mathbb{Z}/3\mathbb{Z}$	`[3]`	–	9	8	9	2	–	5	2	–	–	–	–	–	**35**
$\mathbb{Z}/2\mathbb{Z}$	`[2]`	–	–	–	5	6	5	3	3	–	–	–	–	–	**22**
$\mathbb{Z}/2\mathbb{Z}$ (low-r)	`[2]`	2	3	–	–	–	–	–	–	–	–	–	–	–	**5**
$\mathbb{Z}/5\mathbb{Z}$	`[5]`	12	7	–	1	–	–	–	–	–	–	–	–	–	**20**
$\mathbb{Z}/4\mathbb{Z}$	`[4]`	6	6	3	1	1	–	–	–	–	–	–	–	–	**17**
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$	`[2, 2]`	–	4	7	1	2	1	–	–	1	–	–	–	–	**16**
$\mathbb{Z}/6\mathbb{Z}$	`[6]`	–	1	1	1	1	–	–	–	–	–	–	–	–	**4**
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$	`[4, 2]`	1	1	1	–	–	–	–	–	–	–	–	–	–	**3**
Total Curves		**21**	**31**	**39**	**37**	**28**	**20**	**10**	**13**	**13**	**12**	**1**	**4**	**1**	**230**

Table 6: Summary of #1 Gold Records Established on the NSF ICARM Leaderboard

Torsion T	**Rank**	**Curve ID**	**#1 Gold Record Metrics**	**Additional Podium Finishes**
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$	≥ 9	**#4315**	**#1 Conductor (N), #1 $\lvert\Delta\rvert$, #1 Faltings, #1 Naive**	*Grand Slam: 4 out of 4 #1 records simultaneously*
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$	≥ 8	**#4314**	**#1 Naive Height**	#2 $\lvert\Delta\rvert$, #2 Faltings, #3 Conductor N
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$	≥ 7	**#4313**	**#1 $\lvert\Delta\rvert$, #1 Faltings, #1 Naive**	#6 Conductor N
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$	≥ 6	**#4320**	**#1 $\lvert\Delta\rvert$, #1 Faltings, #1 Naive**	#5 Conductor N
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$	≥ 5	**#4322**	**#1 $\lvert\Delta\rvert$, #1 Faltings, #1 Naive**	#2 Conductor N
$\mathbb{Z}/4\mathbb{Z}$	≥ 8	**#4319**	**#1 $\lvert\Delta\rvert$, #1 Faltings, #1 Naive**	#3 Conductor N
$\mathbb{Z}/4\mathbb{Z}$	≥ 7	**#4318**	**#1 $\lvert\Delta\rvert$, #1 Faltings, #1 Naive**	#3 Conductor N
$\mathbb{Z}/5\mathbb{Z}$	≥ 7	**#4321**	**#1 $\lvert\Delta\rvert$, #1 Faltings, #1 Naive**	#2 Conductor N
$\mathbb{Z}/5\mathbb{Z}$	≥ 5	**#4380**	**#1 Discriminant $\lvert\Delta\rvert$**	#2 $\lvert\Delta\rvert$ (`#4381`), #3 Faltings
$\mathbb{Z}/2\mathbb{Z}$	≥ 9	**#4312**	**#1 Faltings, #1 Naive Height**	#3 Discriminant $\lvert\Delta\rvert$
$\mathbb{Z}/2\mathbb{Z}$	≥ 7	**#4311**	**#1 Naive Height**	#2 $\lvert\Delta\rvert$, #2 Faltings (`#4311`), #3 $\lvert\Delta\rvert$ (`#4511`)
$\mathbb{Z}/3\mathbb{Z}$	≥ 8	**#4317**	**#1 Naive Height**	#2 Faltings (`#4317`), #3 $\lvert\Delta\rvert$ (`#4324`)
$\mathbb{Z}/3\mathbb{Z}$	≥ 7	**#4316**	**#1 Faltings, #1 Naive Height**	#2 $\lvert\Delta\rvert$ (`#4327`), #2 Naive (`#4330`)

5.3 Detailed Case Studies of Landmark Curves

5.3.1 Case Study 1: The "Grand Slam" Curve (#4315, $T = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, Rank ≥ 9)

Curve #4315 (https://elliptic-rank.icarm.cloud/curve/4315), originating from the Layr-Labs / Yukon Elliptic Rank Challenge (2x2-9.ec) and certified with its complete conductor factorization via our 4-Stage Werr Algebraic Sieve, is defined by the minimal Weierstrass equation:

$$E_{4315}: \quad y^2 = x^3 - x^2 - 567553808246049x + 2655925051270455158049. \tag{27}$$

Its certified arithmetic invariants are:

- **Mordell–Weil Rank & Regulator:** $\mathrm{rank}(E_{4315}(\mathbb{Q})) \geq 9$, with regulator $R \approx 16095.67$.
- **Minimal Discriminant (46 digits):** $\Delta = 1172828613149539200258814761002361732009367961 6$.
- **Certified Bad Primes (10 primes):** $\mathcal{S}_{4315} = \{2, 3, 5, 29, 31, 37, 73, 109, 149, 151\}$.
- **Certified Conductor (22 digits):** $N = 1049969472658866185280$ ($\log N \approx 48.404$).
- **Heights:** Faltings height $h_F = 11.233$, Naive height $h_{\text{naive}} = 50.258$.
- **Leaderboard Outcome:** Captures a **Grand Slam**—ranking **#1 in Conductor** N, **#1 in Discriminant** $|\Delta|$, **#1 in Faltings Height**, and **#1 in Naive Height** simultaneously among all known curves with $T = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and rank ≥ 9.

5.3.2 Case Study 2: 110-Digit Discriminant Factorization (#4383, $T = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, Rank ≥ 12)

Curve #4383 (https://elliptic-rank.icarm.cloud/curve/4383), specialized in Dujella's historical tables for $T = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ at rank ≥ 12, is given by the minimal model:

$$\begin{aligned} E_{4383}: \quad y^2 = x^3 - x^2 &- 6773066279657575296262579351595404 33x \\ &+ 1193016672543994534753764971576784359254622489500827 37. \end{aligned} \tag{28}$$

Its minimal discriminant Δ spans **110 decimal digits** (363 bits):

$$\begin{aligned} \Delta_{4383} = 1185513245992202784880227490300308437643763218996215765 \\ \times 10^{55} + 3924667473642622724854325059908145184220980000000000000. \end{aligned}$$

Using Proposition 3.1 (Stage 1 monic cubic root peeling) followed by Stage 2–3 sieving, our pipeline factored Δ_{4383} into **21 distinct prime factors**:

$$\begin{aligned} \mathcal{S}_{4383} = \{2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 41, 61, 71, 101, \\ 241, 317, 1019, 2087, 14009, 283115478426179\}, \end{aligned} \tag{29}$$

certifying its 48-digit conductor $N = 109459202330992507705092784715165810164461886730$ ($\log N \approx 108.316$) and securing #2 across all four metrics on the $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, Rank ≥ 12 leaderboard.

5.3.3 Case Study 3: Elkies Rank ≥ 19 Conductor Certification (#4323, Trivial Torsion)

Curve #4323 (https://elliptic-rank.icarm.cloud/curve/4323), specialized from Elkies' high-rank fibration families and cataloged in Rubinstein's tables (family_trivial.tsv.gz:114), achieves certified Mordell–Weil rank $\mathrm{rank}(E(\mathbb{Q})) \geq 19$ with equation:

$$E_{4323}: \quad y^2 = x^3 - 7394310748527678442369 75x + 331314067119134775943540728349251250. \tag{30}$$

Its 75-digit negative discriminant $\Delta_{4323} = -126092004224607\ldots400000000$ resisted unassisted in-worker factorization due to two large prime factors of 22 and 26 digits. The Werr Algebraic Sieve resolved all **9 bad primes**:

$$\mathcal{S}_{4323} = \{2, 3, 5, 7, 17, 47, 1597, \underbrace{5579416217636942995817}_{\text{22-digit prime}}, \underbrace{80758784220964527622731963}_{\text{26-digit prime}}\}, \tag{31}$$

certifying the 62-digit conductor N_{4323} ($\log N \approx 140.91$) and placing #4323 inside the overall Top-10 across all elliptic curves of rank ≥ 19.

6 Intellectual Attribution and Scientific Lineage

In open science and computational mathematics, new algorithmic syntheses build directly upon decades of foundational theoretical and computational discoveries:

1. **Original Discoverers of High-Rank Parametric Families:** Full credit for the underlying elliptic surface fibrations, Mordell–Weil lattice specializations, and p-adic point searches belongs to their original discoverers: **Prof. Noam D. Elkies** (Harvard University) [9], **Prof. Andrej Dujella** (University of Zagreb) [8], **Prof. Michael Rubinstein** (University of Waterloo) [14], **Dr. Zev Klagsbrun**, and the contributors of the **Layr-Labs / Yukon Elliptic Rank Challenge**, alongside the foundational methodologies of J. Silverman [15], J. Cremona [3], K. Eriksson, and P. Brumer.
2. **Foundational GDT Specification (Pillar I):** We thank **Dan Hawkley** (`@thinkthoughts`) for the foundational specification of the General Divisor Theorem [4], which forms Layer 1 of our multi-layer pre-sieve.
3. **NSF ICARM & Leaderboard Maintainers:** We gratefully acknowledge the **NSF Institute for Computer-Aided Reasoning in Mathematics (ICARM)**, supported under **U.S. National Science Foundation Grant DMS-2425401**, and thank **David Renshaw** (`@dwrensha`) and **Seewoo Lee** (`@seewoo5`) for building and maintaining the interactive PARI/GP 2.17.2 WebAssembly verification platform.
4. **Computing Infrastructure:** Dedicated bare-metal verification clusters and high-memory compute nodes were provided by **Bekirhan Dağlı** (Managing Partner, ITouch Systems Research & Development).

7 Conclusion and Future Work

By uniting the 4-Stage Werr Multi-Layer Algebraic Sieve with systematic Vélu Isogeny Graph Transport, we resolved the conductor certification bottleneck across 230 high-rank elliptic curves over $\mathbb{Q}$ on the NSF ICARM leaderboard, achieving 100% certified conductor factorizations, 30 #1 Gold records, 80 podiums, and 418 Top-10 placements. Future work will extend this certificate-driven methodology to genus-2 Jacobian varieties over $\mathbb{Q}$ and formalize Tate's local reduction algorithm directly in Lean 4.

Data Availability and Reproducibility

All 230 certified curves, complete Weierstrass models, rational generator coordinates, certified bad prime factorizations, and standalone Python reproduction scripts are permanently available under open licenses:

- **CERN Zenodo Concept DOI:** `https://doi.org/10.5281/zenodo.23436282`
- **CERN Zenodo Release v1.0 DOI:** `https://doi.org/10.5281/zenodo.23436283`
- **GitHub Repository:** `https://github.com/pCwOrM/werr-elliptic-sieve`
- **Official ICARM Contributor Profile:** `https://elliptic-rank.icarm.cloud/user/114`

References

[1] J. Avigad, *Mathematical Logic and Formalized Mathematics*, Cambridge University Press, 2021.

[2] J. C. Blanchette, M. Haslbeck, D. Matichuk, and T. Nipkow, Mining the Archive of Formal Proofs, in *Proc. Int. Conf. on Intelligent Computer Mathematics (CICM)*, Springer, pp. 3–17, 2016.

[3] J. E. Cremona, *Algorithms for Modular Elliptic Curves*, 2nd ed., Cambridge University Press, 1997.

[4] V. Dağlı, Z. Dağlı, D. Dağlı, and D. Hawkley, *General Divisor Theorem (GDT v3.0): 81 Lean 4 Machine-Certified Theorems, Exact $O(1)$ Block Sieving, and Primorial Radical Contraction*, CERN Zenodo, DOI: 10.5281/zenodo.23425590, 2026.

[5] V. Dağlı, Z. Dağlı, and D. Dağlı, *Machine-Certified Formalization of the 24-Dimensional Niemeier Lattice A_8^3, K3 Hodge Duality and Umbral Moonshine in Lean 4*, CERN Zenodo, DOI: 10.5281/zenodo.23324067, 2026.

[6] V. Dağlı, Z. Dağlı, and D. Dağlı, *GAP-Lean4-Port (v0.2.0): Formalization of Sims Stabilizer Chains (`GAP-0190`) and Orbit-Stabilizer Coset Enumeration (`GAP-0247`) in Lean 4*, CERN Zenodo, DOI: 10.5281/zenodo.23045504, 2026.

[7] V. Dağlı, Z. Dağlı, and D. Dağlı, *Bipartite Alternating Carrier Parity Flips and Word-Packed $\mathbb{F}_2$ Linear Algebra for Integer Multiplication and Sieve Acceleration*, CERN Zenodo, DOI: 10.5281/zenodo.23268580, 2026.

[8] A. Dujella, *High rank elliptic curves with prescribed torsion group*, Online Tables, Department of Mathematics, University of Zagreb, https://web.math.pmf.unizg.hr/~dujella/tors/tors.html, accessed October 2026.

[9] N. D. Elkies, $\mathbb{Z}/2\mathbb{Z}$-torsion elliptic curves over $\mathbb{Q}$ with rank at least 19, and an elliptic curve over $\mathbb{Q}$ of rank at least 28, *Number Theory Listserver Announcements*, Harvard University, 2006.

[10] T. Hales et al., A formal proof of the Kepler conjecture, *Forum of Mathematics, Pi*, vol. 5, e2, pp. 1–29, 2017.

[11] D. Renshaw and S. Lee, *Elliptic Curve Rank Leaderboard*, NSF Institute for Computer-Aided Reasoning in Mathematics (ICARM), U.S. National Science Foundation Grant DMS-2425401, https://elliptic-rank.icarm.cloud, 2026.

[12] B. Mazur, Modular curves and the Eisenstein ideal, *Publications Mathématiques de l'IHÉS*, vol. 47, pp. 33–186, 1977.

[13] The PARI Group, *PARI/GP version 2.17.2*, Université de Bordeaux, https://pari.math.u-bordeaux.fr/, 2024.

[14] M. Rubinstein, *Elliptic curves by rank and torsion*, Computational Number Theory Database, University of Waterloo, 2024.

[15] J. H. Silverman, *The Arithmetic of Elliptic Curves*, Graduate Texts in Mathematics, vol. 106, Springer-Verlag, New York, 1986.

[16] J. Tate, Algorithm for determining the type of a singular fiber in an elliptic pencil, in *Modular Functions of One Variable IV*, Lecture Notes in Mathematics, vol. 476, Springer, pp. 33–52, 1975.

[17] J. Vélu, Isogénies entre courbes elliptiques, *Comptes Rendus de l'Académie des Sciences de Paris, Série A-B*, vol. 273, pp. A238–A241, 1971.