Sharp linear hypercontractivity for ultraspherical measures

Paata Ivanisvili

This proof was generated on October 2, 2026, by Grok Bot using Cursor's agents, guided by several prompts from Paata Ivanisvili

Abstract

The linear hypercontractive estimate of Ivanisvili, Lindenberger, Müller and Schmuckenschläger on the circle, proved in their work for exponents $6 \le p \le q$, holds for every admissible pair. For $m \ge -1$ and $q \ge p > 0$ with $p + m > 0$,

$$\|1 + rbz\|_{L^q(\nu_m)} \le \|1 + bz\|_{L^p(\nu_m)}$$

for all real b, if and only if $|r| \le \sqrt{(p+m)/(q+m)}$. At $m = -1$ the restriction $p + m > 0$ is Bonami's condition $p > 1$. The same threshold is sharp for the corresponding linear estimate on every sphere.

1 Statement

For $m > -1$ let ν_m be the ultraspherical probability measure on the unit circle,

$$d\nu_m(e^{i\theta}) = c_m \,|\sin\theta|^m \, d\theta, \qquad c_m = \frac{\Gamma(m/2+1)}{2\sqrt{\pi}\,\Gamma(m/2+1/2)},$$

the integral running over $[0, 2\pi]$, and let $\nu_{-1} = \frac{1}{2}(\delta_1 + \delta_{-1})$. The constant c_m is chosen so that ν_m is a probability measure. Indeed,

$$\int_0^{\pi/2} \sin^m\theta \, d\theta = \frac{1}{2} B\Big(\frac{m+1}{2}, \frac{1}{2}\Big) = \frac{\sqrt{\pi}\,\Gamma(m/2+1/2)}{2\,\Gamma(m/2+1)},$$

and four copies of this integral give

$$\int_0^{2\pi} |\sin\theta|^m \, d\theta = 2\sqrt{\pi}\,\frac{\Gamma(m/2+1/2)}{\Gamma(m/2+1)}.$$

The estimate under consideration is

$$\|1 + rbz\|_{L^q(\nu_m)} \le \|1 + bz\|_{L^p(\nu_m)} \qquad \text{for every } b \in \mathbb{R}. \tag{1}$$

Theorem 1. *Let $m \ge -1$ and $q \ge p > 0$ with $p + m > 0$. Then (1) holds for every real b if and only if*

$$|r| \le \sqrt{\frac{p+m}{q+m}}.$$

At $m = -1$, the hypothesis $p + m > 0$ is exactly $p > 1$.

The case $m = -1$ is Bonami's theorem, and the case $m = 0$ is Weissler's theorem on the circle. Theorem 1 removes the restriction $6 \le p \le q$ from Theorem 3.1 of Ivanisvili–Lindenberger–Müller–Schmuckenschläger, and it extends their Theorem 2.1 from the exponents $0 < p \le 2 = q$ on the sphere to every pair $q \ge p > 0$.

Theorem 2. *Let $n \geq 2$, let σ be the uniform probability measure on $\mathbb{S}^{n-1}$, and let $q \geq p > 0$. For every $x \in \mathbb{R}^n$ and every real a,*

$$\|x + ary\|_{L^q(\mathbb{S}^{n-1},\sigma)} \leq \|x + ay\|_{L^p(\mathbb{S}^{n-1},\sigma)}$$

whenever $|r| \leq \sqrt{(p+n-2)/(q+n-2)}$. The constant is sharp.

Throughout, $m = 2\lambda$ with $\lambda > -1/2$, and μ_λ denotes the law of $\operatorname{Re} z$ under ν_m. Explicitly, μ_λ is the probability measure on $[-1, 1]$ with density

$$\frac{\Gamma(\lambda+1)}{\sqrt{\pi}\,\Gamma(\lambda+1/2)}\,(1-t^2)^{\lambda-1/2}.$$

Indeed, $\cos\theta = t$ has two preimages in $[0, 2\pi)$, each with Jacobian $1/\sqrt{1-t^2}$, so the density of the push-forward is

$$2c_{2\lambda}(1-t^2)^{\lambda-1/2},$$

and $2c_{2\lambda}$ is the constant displayed above. For $c \geq 0$ write

$$F_c(t) = 1 + 2ct + c^2$$

and

$$N_\alpha(c) = \left(\int_{-1}^{1} F_c(t)^\alpha \, d\mu_\lambda(t)\right)^{1/\alpha},$$

for every $\alpha > 0$ with $\alpha + \lambda > 0$. Since ν_m is invariant under $z \mapsto -z$,

$$\|1 + bz\|_{L^{2\alpha}(\nu_m)} = N_\alpha(|b|)^{1/2}.$$

2 Necessity

Proposition 3. *If $m > -1$ and $p + m > 0$, $q + m > 0$, and if (1) holds for every small real b, then $|r| \leq \sqrt{(p+m)/(q+m)}$.*

Proof. The identity $\cos^2\theta = 1 - \sin^2\theta$ gives

$$\int (\operatorname{Re} z)^2 \, d\nu_m = 1 - \frac{c_m}{c_{m+2}}.$$

The recurrence $\Gamma(w+1) = w\,\Gamma(w)$ gives

$$\frac{c_m}{c_{m+2}} = \frac{\Gamma(m/2+1)}{\Gamma(m/2+2)} \cdot \frac{\Gamma(m/2+3/2)}{\Gamma(m/2+1/2)} = \frac{m/2+1/2}{m/2+1} = \frac{m+1}{m+2},$$

so the second moment equals $1 - (m+1)/(m+2) = 1/(m+2)$.

For $|b| < 1/2$ one has $1 + 2b\operatorname{Re} z + b^2 \geq (1-|b|)^2 > 1/4$. Taylor expansion with a uniform remainder therefore yields

$$\begin{aligned}\int (1 + 2b\operatorname{Re} z + b^2)^{p/2} \, d\nu_m &= 1 + \frac{p}{2}b^2 + \frac{p(p-2)}{2(m+2)}b^2 + O(b^3) \\ &= 1 + \frac{pb^2}{2} \cdot \frac{p+m}{m+2} + O(b^3).\end{aligned}$$

The odd contributions in b integrate to zero by the symmetry $z \mapsto -z$. Raising to the power $1/p$ gives

$$\|1 + bz\|_{L^p(\nu_m)} = 1 + \frac{b^2}{2} \cdot \frac{p+m}{m+2} + O(b^3).$$

The same expansion at exponent q, with b replaced by rb, gives

$$\|1 + rbz\|_{L^q(\nu_m)} = 1 + \frac{r^2 b^2}{2} \cdot \frac{q+m}{m+2} + O(b^3).$$

Subtract the constant term 1, divide by b^2, and let $b \to 0$. Inequality (1) forces

$$r^2 \frac{q+m}{m+2} \le \frac{p+m}{m+2}.$$

□

The same expansion at $m = -1$, where $\int (\operatorname{Re} z)^2 \, d\nu_{-1} = 1$, gives the necessity of $|r| \le \sqrt{(p-1)/(q-1)}$ for $p, q > 1$.

3 The hypergeometric representation

For $0 \le c < 1$ and every real α,

$$\int F_c^\alpha \, d\mu_\lambda = {}_2F_1(-\alpha, -\alpha - \lambda; \lambda + 1; c^2). \tag{2}$$

The identity $F_c = c^2 F_{1/c}$ gives, for every $c > 0$,

$$N_\alpha(c) = c^2 N_\alpha(1/c). \tag{3}$$

To prove (2), set $z = c^2$ and $y(z) = \int F_{\sqrt{z}}^\alpha \, d\mu_\lambda$. The function

$$Y(z) = {}_2F_1(-\alpha, -\alpha - \lambda; \lambda + 1; z)$$

is the solution of

$$z(1-z)Y'' + \big[\lambda + 1 - (1 - 2\alpha - \lambda)z\big]Y' - \alpha(\alpha + \lambda)Y = 0 \tag{4}$$

which is analytic at the origin and satisfies $Y(0) = 1$. The constant term of y is 1. The coefficient of z is computed from

$$F_c^\alpha = 1 + \alpha(b^2 + 2bt) + \alpha(\alpha - 1)2b^2 t^2 + O(b^3),$$

with $b = \sqrt{z}$. The remainder $O(b^3)$ is $O(z^{3/2})$, hence $o(z)$. Together with $\int t \, d\mu_\lambda = 0$ and

$$\int t^2 \, d\mu_\lambda = \frac{1}{2(\lambda + 1)}.$$

(The second moment is the case $m = 2\lambda$ of the computation in Proposition 3.) Integrating and using $\int t \, d\mu_\lambda = 0$ produces

$$y(z) = 1 + \alpha z + \frac{\alpha(\alpha - 1)}{\lambda + 1} \, z + o(z) = 1 + \frac{\alpha(\alpha + \lambda)}{\lambda + 1} \, z + o(z),$$

so

$$y'(0) = \frac{\alpha(\alpha+\lambda)}{\lambda+1}.$$

The derivative of ${}_2F_1(a,b;c_*;z)$ at the origin equals ab/c_*. With $a=-\alpha$, $b=-\alpha-\lambda$ and $c_*=\lambda+1$, this is the same value. The map $c \mapsto \int F_c^\alpha \, d\mu_\lambda$ is analytic for $|c|<1$, because $F_c \geq (1-|c|)^2 > 0$, and it is even by the substitution $t \mapsto -t$. An even analytic function of c is analytic in c^2, so y is analytic on $[0,1)$. The hypergeometric series is the unique analytic solution of (4) with value 1 at the origin: the constant term of the differential equation forces the coefficient of z, and each higher coefficient is then determined by the recurrence. Thus it is enough to show that y satisfies (4) on $(0,1)$.

Lemma 4. *Let $c \in \mathbb{R}$ and let $F = F_c$. On $(-1,1)$,*

$$\frac{d}{dt}\Big(F^{\alpha-1}(1-t^2)^{\lambda+1/2}\Big) = -Q\,F^{\alpha-2}(1-t^2)^{\lambda-1/2},$$

where

$$Q = (2\lambda+1)(1+c^2)\,t + 2c(\alpha+2\lambda)\,t^2 + 2c(1-\alpha).$$

Proof. Differentiate the product. With $F' = 2c$,

$$\begin{aligned}\frac{d}{dt}\Big(F^{\alpha-1}(1-t^2)^{\lambda+1/2}\Big) &= 2c(\alpha-1)F^{\alpha-2}(1-t^2)^{\lambda+1/2} - t(2\lambda+1)F^{\alpha-1}(1-t^2)^{\lambda-1/2} \\ &= F^{\alpha-2}(1-t^2)^{\lambda-1/2}\Big(2c(\alpha-1)(1-t^2) - t(2\lambda+1)F\Big).\end{aligned}$$

The expression in parentheses equals $-Q$, because

$$\begin{aligned}t(2\lambda+1)F - 2c(\alpha-1)(1-t^2) &= (2\lambda+1)(1+c^2)t + 2c(2\lambda+1)t^2 + 2c(\alpha-1)t^2 - 2c(\alpha-1) \\ &= (2\lambda+1)(1+c^2)t + 2c(\alpha+2\lambda)t^2 + 2c(1-\alpha).\end{aligned}$$

□

For $c \in (0,1)$ one has $F \geq (1-c)^2 > 0$, so $F^{\alpha-1}$ is bounded on $[-1,1]$. The boundary factor $(1-t^2)^{\lambda+1/2}$ vanishes at $t=\pm1$ because $\lambda > -1/2$. The derivative on the left of Lemma 4 is integrable on $(-1,1)$, since near each endpoint it is comparable to $(1-t^2)^{\lambda-1/2}$ and $\lambda - 1/2 > -1$. Consequently

$$\int Q\,F^{\alpha-2}\,d\mu_\lambda = 0.$$

Now let $I(c) = \int F_c^\alpha \, d\mu_\lambda$, so that $y(z) = I(\sqrt{z})$. Then

$$y_z = \frac{I_c}{2c}, \qquad y_{zz} = \frac{I_{cc}}{4c^2} - \frac{I_c}{4c^3}.$$

Substituting $z = c^2$ into the differential expression (4) and clearing the denominator $4c^3$ produces, under the integral sign, the integrand identity

$$\begin{aligned}&z(1-z)\partial_{zz}(F^\alpha) + \big[\lambda+1-(1-2\alpha-\lambda)z\big]\partial_z(F^\alpha) - \alpha(\alpha+\lambda)F^\alpha \\ &\quad = -\frac{\alpha(c^2-1)}{2c}\,Q\,F^{\alpha-2}.\end{aligned} \tag{5}$$

(The two sides are rational functions of c and t, once $F^{\alpha-2}$ is divided out, and the resulting polynomial identity is the expansion of Lemma 4 against the chain rule; it is recorded in the appendix.) Integrating (5) and using $\int QF^{\alpha-2}\,d\mu_\lambda = 0$ shows that y satisfies (4). This proves (2).

For $\alpha > 0$ and $z \in (0,1)$ define

$$\tau_\alpha(z) = \frac{1}{\alpha}\,\partial_z \log y(\alpha, z) = \frac{\partial_z y(\alpha, z)}{\alpha\, y(\alpha, z)},$$

where $y(\alpha, z)$ denotes the right-hand side of (2). Then τ_α extends continuously to $z = 0$ by

$$\tau_\alpha(0) = \frac{\alpha + \lambda}{\lambda + 1},$$

and the substitution $\partial_z \log y = \alpha\tau_\alpha$ turns (4) into the Riccati equation

$$z(1-z)\tau' = (\alpha + \lambda) - \big[\lambda + 1 - (1 - 2\alpha - \lambda)z\big]\tau - z(1-z)\,\alpha\,\tau^2. \tag{6}$$

In particular,

$$\frac{1}{\alpha}\log y(\alpha, z) = \int_0^z \tau_\alpha(t)\,dt. \tag{7}$$

Lemma 5. *If $\alpha > 0$ and $\alpha + \lambda > 0$, then $\tau_\alpha(z) > 0$ for every $z \in [0, 1)$.*

Proof. The value at the origin is positive. If $z \in (0,1)$ were a first zero, (6) at $\tau = 0$ would give $z(1-z)\tau' = \alpha + \lambda > 0$, so $\tau' > 0$, which is incompatible with reaching zero from above. □

4 An upper bound for the logarithmic derivative

Lemma 6. *For every $\lambda > -1/2$, every $\alpha > 0$ and every $z \in (0,1)$,*

$$\tau_\alpha(z) < \frac{1}{\sqrt{z}\,(1 + \sqrt{z})}.$$

In particular $\tau_\alpha(z) < 1/(2z)$.

Proof. Write $\sigma(z) = 1/(\sqrt{z}\,(1+\sqrt{z}))$ and $s = \sqrt{z}$, so that $\sigma(z) = 1/(s(s+1))$. As $z \downarrow 0$ one has $\sigma(z) \to \infty$, while $\tau_\alpha(z)$ tends to the finite value $(\alpha + \lambda)/(\lambda + 1)$. Thus $\tau_\alpha < \sigma$ on a right neighbourhood of the origin.

Set $t = 1/(s(s+1))$ and substitute $\tau = t$, $z = s^2$ into the right-hand side of (6). The quadratic term contributes

$$z(1-z)\,\alpha\,\tau^2 \cdot s(s+1) = \alpha\, s^2(1-s^2) \cdot \frac{1}{s(s+1)} = \alpha\, s(1-s),$$

where $1 - s^2 = (1-s)(1+s)$ and $(1+s)/(s+1) = 1$. Multiplying the whole right-hand side by $s(s+1)$ therefore gives

$$\begin{aligned}(\alpha + \lambda)s(s+1) - \big[\lambda + 1 - (1 - 2\alpha - \lambda)s^2\big] - \alpha s(1-s)\\ = s^2 + \lambda s - (\lambda + 1) = (s-1)(s + \lambda + 1).\end{aligned}$$

(The coefficient of s^2 is $(\alpha+\lambda) + (1 - 2\alpha - \lambda) + \alpha = 1$, and the coefficient of s is $(\alpha+\lambda) - \alpha = \lambda$.) Therefore, at every point where $\tau_\alpha = \sigma$,

$$\tau_\alpha' = \frac{(s-1)(s+\lambda+1)}{s(s+1)\cdot s^2(1-s^2)} = -\frac{s + \lambda + 1}{s^3(s+1)^2},$$

where $1-s^2=(1-s)(1+s)$ and $s-1=-(1-s)$ were used. On the other hand,

$$\frac{d}{ds}\frac{1}{s(s+1)}=-\frac{1+2s}{s^2(s+1)^2},\qquad \frac{ds}{dz}=\frac{1}{2s},$$

so

$$\sigma'(z)=-\frac{1+2s}{2s^3(s+1)^2}.$$

Subtracting these evaluations yields

$$\tau_\alpha'-\sigma'=\frac{-2(s+\lambda+1)+(1+2s)}{2s^3(s+1)^2}=-\frac{2\lambda+1}{2s^3(s+1)^2}<0,$$

because $\lambda>-1/2$. At a first zero of $\sigma-\tau_\alpha$ in $(0,1)$ one would have $\tau_\alpha'\geq\sigma'$. This contradiction keeps $\tau_\alpha<\sigma$ on the whole interval.

Finally, $z+\sqrt{z}\geq 2z$ for $z\in(0,1]$, so $\sigma(z)\leq 1/(2z)$, and the inequality is strict for $z\in(0,1)$. □

Lemma 7. *For every $\alpha>0$ with $\alpha+\lambda>0$, the map $c\mapsto N_\alpha(c)$ is continuous and strictly increasing on $[0,\infty)$.*

Proof. For $c\in(0,1)$, (2) and the chain rule give

$$\frac{N_\alpha'(c)}{N_\alpha(c)}=2c\,\tau_\alpha(c^2).$$

Lemma 5 makes the right-hand side positive. For $c>1$, (3) gives

$$\frac{d}{dc}\log N_\alpha(c)=\frac{2}{c}-\frac{1}{c^2}\cdot\frac{N_\alpha'(1/c)}{N_\alpha(1/c)}=\frac{2}{c}-\frac{2}{c^3}\,\tau_\alpha(c^{-2}).$$

Lemma 6 yields $\tau_\alpha(c^{-2})<c^2/2$, so the derivative is positive. Continuity on $[0,\infty)$ holds because $0\leq F_c\leq(1+c)^2$ and $\alpha>0$. □

Lemma 7 removes the restriction to the critical multiplier. The law of z is symmetric, so b may be replaced by $|b|$. For $0\leq r\leq r_*$ and $c\geq 0$,

$$N_{q/2}(rc)\leq N_{q/2}(r_*c).$$

Thus (1) holds for every r with $|r|\leq\sqrt{(p+m)/(q+m)}$ as soon as it holds at the positive endpoint.

5 Scales at most one

Set $\gamma=p/2$ and $\delta=q/2$. The hypotheses $q\geq p>0$ and $p+m>0$ become $0<\gamma\leq\delta$ and $\gamma+\lambda>0$. Set

$$\kappa=\frac{\gamma+\lambda}{\delta+\lambda}\in(0,1].$$

Inequality (1) at the critical multiplier is equivalent to

$$N_\gamma(c)\geq N_\delta(c\sqrt{\kappa})\qquad\text{for every } c\geq 0. \tag{8}$$

Lemma 8. *For every $z\in(0,1)$,*

$$\tau_\gamma(z)>\kappa\,\tau_\delta(\kappa z)$$

whenever $\delta>\gamma$, and therefore $N_\gamma(\sqrt{z})\geq N_\delta(\sqrt{\kappa z})$. In particular, (8) holds for every $c\in[0,1]$.

Proof. The case $\gamma = \delta$ is the identity $\kappa = 1$, so assume $\delta > \gamma$. Write $\psi(z) = \kappa\,\tau_\delta(\kappa z)$. Then $\tau_\gamma(0) = \psi(0) = (\gamma+\lambda)/(\lambda+1)$. Differentiate (6). The derivative of the left side is $(1-2z)\tau' + z(1-z)\tau''$, which equals $\tau'(0)$ at $z = 0$. At $z = 0$ and $\tau = t_0 = (\alpha+\lambda)/(\lambda+1)$ both sides of (6) vanish. Differentiating the right side, the product rule contributes $(1-2\alpha-\lambda)\tau$ from the coefficient of τ and $-\alpha\tau^2$ from $z(1-z)\alpha\tau^2$, together with the terms that still contain τ'. At the origin this equals

$$(1-2\alpha-\lambda)t_0 - (\lambda+1)\tau'(0) - \alpha t_0^2.$$

Solving for the derivative gives

$$\tau'_\alpha(0) = \frac{t_0}{\lambda+2}\big(1-2\alpha-\lambda-\alpha t_0\big). \tag{9}$$

The chain rule yields $\psi'(0) = \kappa^2\tau'_\delta(0)$. Since $\kappa t_\delta = t_\gamma$, with $t_\alpha = (\alpha+\lambda)/(\lambda+1)$,

$$\tau'_\gamma(0) - \psi'(0) = \frac{t_\gamma}{\lambda+2}\Big((1-2\gamma-\lambda-\gamma t_\gamma) - \kappa(1-2\delta-\lambda-\delta t_\delta)\Big).$$

The expression in parentheses equals

$$\frac{\delta-\gamma}{(\delta+\lambda)(\lambda+1)}\big(\gamma\delta + \lambda(\gamma+\delta) + 2\lambda^2 + 2\lambda + 1\big).$$

(The verification is the expansion in the appendix.) The quadratic form factors as

$$\gamma\delta + \lambda(\gamma+\delta) + 2\lambda^2 + 2\lambda + 1 = (\gamma+\lambda)(\delta+\lambda) + (\lambda+1)^2 > 0.$$

Hence $\tau'_\gamma(0) - \psi'(0) > 0$. Combined with $\tau_\gamma(0) = \psi(0)$, this forces $\tau_\gamma > \psi$ on a right neighbourhood of the origin.

Suppose there exists a point of $(0,1)$ at which $\tau_\gamma = \psi$. Let z_* be the infimum of such points. Then $z_* > 0$, $\tau_\gamma(z_*) = \psi(z_*)$, and $\tau_\gamma - \psi \geq 0$ on $[0, z_*]$, so the derivative of the difference at z_* is at most zero. It remains to rule that out.

Let $z \in (0,1)$ satisfy $\tau_\gamma(z) = \psi(z) = t$. The equation (6) at exponent γ reads

$$z(1-z)\tau'_\gamma = N_\gamma(t),$$

where

$$N_\gamma(t) = (\gamma+\lambda) - \big[\lambda+1-(1-2\gamma-\lambda)z\big]t - z(1-z)\gamma t^2.$$

For the comparison function, set $u = \kappa z$ and $\tau_\delta(u) = \psi(z)/\kappa$. Then $\psi'(z) = \kappa^2\, d\tau_\delta/du$. Substitute into (6) at exponent δ, multiply through by κ^2, and use $\kappa^2(\delta+\lambda) = \kappa(\gamma+\lambda)$. Dividing by $\kappa > 0$ produces

$$z(1-\kappa z)\psi' = N_\psi(t), \tag{10}$$

where

$$N_\psi(t) = (\gamma+\lambda) - \big[\lambda+1-(1-2\delta-\lambda)\kappa z\big]t - z(1-\kappa z)\delta t^2.$$

Therefore

$$\tau'_\gamma(z) - \psi'(z) = \frac{N_\gamma(t)\,(1-\kappa z) - N_\psi(t)\,(1-z)}{z(1-z)(1-\kappa z)}.$$

The denominator is positive. The numerator equals

$$\frac{z(\delta-\gamma)}{\delta+\lambda}\,Q(t),$$

where

$$Q(t) = A_Q t^2 - 2z(\gamma+\lambda)\,t + (\gamma+\lambda), \qquad A_Q = (1-z)\big(\delta+\lambda - z(\gamma+\lambda)\big).$$

This polynomial identity is checked coefficient-wise in the appendix. The prefactor $z(\delta-\gamma)/(\delta+\lambda)$ is positive, so the sign of $\tau_\gamma' - \psi'$ is the sign of $Q(t)$.

Since $\delta \geq \gamma$ and $z \in (0,1)$,

$$A_Q - (1-z)^2(\gamma+\lambda) = (1-z)(\delta-\gamma) \geq 0,$$

and hence

$$Q(t) \geq (\gamma+\lambda)\big((1-z)^2t^2 - 2zt + 1\big).$$

For $z \in (0, 1/2]$ the discriminant of the quadratic in t is $4(2z-1) \leq 0$, so the quadratic is positive for every finite t except at the single point $z = 1/2$, $t = 2$, where it vanishes. For $z \in (1/2, 1)$ the roots are

$$\frac{1}{z \pm \sqrt{2z-1}},$$

because

$$\big(z - \sqrt{2z-1}\big)\big(z + \sqrt{2z-1}\big) = (1-z)^2.$$

The quadratic is then nonnegative for $0 \leq t \leq 1/(z+\sqrt{2z-1})$. Lemma 6 supplies the stricter bound

$$t = \tau_\gamma(z) < \frac{1}{z+\sqrt{z}} \leq \frac{1}{z+\sqrt{2z-1}}$$

on $[1/2, 1)$, the comparison of the denominators being $\sqrt{z} \geq \sqrt{2z-1}$. At $z = 1/2$ the same bound gives $\tau_\gamma(1/2) < 1/(1/2+\sqrt{1/2}) < 2$. Thus $Q(\tau_\gamma(z)) > 0$, and $\tau_\gamma' > \psi'$ at every contact point in $(0,1)$.

The graphs cannot meet in $(0,1)$. Therefore $\tau_\gamma(z) > \psi(z)$ for every $z \in (0,1)$. Integrating and applying (7) gives

$$\frac{1}{\gamma}\log y(\gamma, z) \geq \frac{1}{\delta}\log y(\delta, \kappa z),$$

which is $N_\gamma(\sqrt{z}) \geq N_\delta(\sqrt{\kappa z})$. The endpoint $c = 1$ follows by continuity of N_α. □

6 Scales larger than one

Lemma 9. *For each fixed $\alpha > 0$ with $\alpha + \lambda > 0$, the map $a \mapsto N_\alpha(a)/a$ is strictly decreasing on $(0,1]$.*

Proof. For $a \in (0,1)$,

$$\frac{d}{da}\log\frac{N_\alpha(a)}{a} = 2a\,\tau_\alpha(a^2) - \frac{1}{a}.$$

Lemma 6 yields $2a\,\tau_\alpha(a^2) < 2a\cdot(1/(2a^2)) = 1/a$, so the derivative is negative. If $0 < a < 1$, the same derivative on $(a,1)$ gives $N_\alpha(a)/a > N_\alpha(1)/1$. Thus the map is strictly decreasing on $(0,1]$. □

In particular, $0 < b < a \leq 1$ implies $N_\alpha(b)/b > N_\alpha(a)/a$.

Proposition 10. *Inequality* (8) *holds for every $c > 1$.*

Proof. Write $\rho(c) = N_\gamma(c)/N_\delta(c\sqrt{\kappa})$. Lemma 8 at scale $1/c \in (0,1)$ gives

$$N_\gamma(1/c) \geq N_\delta\big((1/c)\sqrt{\kappa}\big).$$

Suppose first that $c\sqrt{\kappa} > 1$, and set $a = 1/(c\sqrt{\kappa}) \in (0,1)$. Then $(1/c)\sqrt{\kappa} = \kappa a$. The scaling relation (3) gives

$$\rho(c) = \frac{N_\gamma(1/c)}{\kappa\, N_\delta(a)} = \frac{N_\gamma(1/c)}{N_\delta(\kappa a)} \cdot \frac{N_\delta(\kappa a)}{\kappa\, N_\delta(a)}.$$

The first factor is at least 1 by Lemma 8. If $\kappa < 1$, then $0 < \kappa a < a < 1$, so Lemma 9 makes the second factor strictly larger than 1. If $\kappa = 1$, both factors equal 1. Thus $\rho(c) \geq 1$.

Suppose instead that $c > 1$ and $c\sqrt{\kappa} \leq 1$. Then necessarily $\kappa < 1$. Set $x = c\sqrt{\kappa} \in (\sqrt{\kappa}, 1]$ and $y = \kappa/x$. The inequality $x > \sqrt{\kappa}$ forces $y < x$, and both points lie in $(0,1]$. Also $y = (1/c)\sqrt{\kappa}$ and $c^2 = x^2/\kappa$. Hence

$$\rho(c) = \frac{c^2 N_\gamma(1/c)}{N_\delta(x)} \geq \frac{x^2}{\kappa} \cdot \frac{N_\delta(y)}{N_\delta(x)},$$

where the inequality is Lemma 8. The relation $\kappa = xy$ rewrites the desired bound $x^2 N_\delta(y) \geq \kappa N_\delta(x)$ as

$$\frac{N_\delta(y)}{y} \geq \frac{N_\delta(x)}{x},$$

which is Lemma 9. Therefore $\rho(c) \geq 1$. □

Lemmas 8 and 10 are (8) for every $c \geq 0$. By the reduction of Section 7 below, this is Theorem 1 for $m > -1$.

7 The endpoint and the sphere

For $m > -1$, the identity $\|1 + bz\|_{L^{2\alpha}(\nu_m)} = N_\alpha(|b|)^{1/2}$ converts (8) into (1) at

$$r = \sqrt{\kappa} = \sqrt{\frac{p+m}{q+m}}.$$

Lemma 7 extends the inequality to every smaller multiplier, and Proposition 3 gives the matching necessity.

If $m = -1$ and $p > 1$, then $p + m' > 0$ for every $m' \in (-1, 0)$. The case $m > -1$ of the theorem, already proved, gives (1) for $\nu_{m'}$ at the multiplier $\sqrt{(p+m')/(q+m')}$. As $m' \downarrow -1$,

$$\int (\operatorname{Re} z)^2 \, d\nu_{m'} = \frac{1}{m'+2} \to 1.$$

Every weak-star limit point ν satisfies $\int (\operatorname{Re} z)^2 \, d\nu = 1$. Then $\int \big(1 - (\operatorname{Re} z)^2\big) \, d\nu = 0$. The integrand is continuous and nonnegative, so ν is supported on $\{\pm 1\}$. Each $\nu_{m'}$ is invariant under $z \mapsto -z$, so the same is true of ν, and therefore $\nu = \nu_{-1}$. The circle is compact, so $\nu_{m'}$ converges weak-star to ν_{-1}. Fix $b \in \mathbb{R}$ and set $r(m') = \sqrt{(p+m')/(q+m')}$. Then $r(m')$ tends to $r_* = \sqrt{(p-1)/(q-1)}$, and $z \mapsto |1 + r(m')bz|^q$ converges uniformly to $z \mapsto |1 + r_* bz|^q$. The function $z \mapsto |1 + bz|^p$ is continuous. Weak-star convergence preserves both integrals, so the norms pass to the limit and (1) holds for ν_{-1} at the multiplier r_*. The expansion recorded after Proposition 3 gives the matching necessity.

Proof of Theorem 2. If $x = 0$, then $|ay|$ is the constant $|a|$ for every $y \in \mathbb{S}^{n-1}$, and the asserted bound reduces to $|r| \le 1$. This holds because $q \ge p$. Both sides are homogeneous of degree one in (x, a), so one may assume $|x| = 1$. For $y \in \mathbb{S}^{n-1}$,

$$|x + ay|^2 = 1 + 2a\langle x, y\rangle + a^2.$$

The set $\{y \in \mathbb{S}^{n-1} : \langle x, y\rangle \in dt\}$ has area proportional to $(1 - t^2)^{(n-3)/2}\, dt$. With $\lambda = (n-2)/2$, the exponent $(n-3)/2$ equals $\lambda - 1/2$, so the law of $\langle x, y\rangle$ is μ_λ, and the corresponding circle weight is $m = n - 2$. Symmetry in t gives

$$\|x + ay\|_{L^p(\sigma)} = N_{p/2}(|a|)^{1/2},$$

and the same identity with a replaced by ar at the exponent q. Theorem 1 at weight $m = n - 2$ is therefore the spherical estimate. Sharpness is Proposition 3 at this weight. □

Appendix: the polynomial identities

The integrand form of the hypergeometric operator

Let $z = c^2$, $F = 1 + 2ct + c^2$ and $\partial_c F = 2(c + t)$. Write $L[F^\alpha]$ for the left-hand side of (5). The conversions $\partial_z = (1/(2c))\partial_c$ and

$$\partial_{zz} = \frac{1}{4c^2}\partial_{cc} - \frac{1}{4c^3}\partial_c$$

give, after multiplication by $4c^3F^{2-\alpha}$,

$$\begin{aligned}\partial_c(F^\alpha)\, F^{2-\alpha} &= 2\alpha(c + t)F,\\ \partial_{cc}(F^\alpha)\, F^{2-\alpha} &= 4\alpha(\alpha - 1)(c + t)^2 + 2\alpha F.\end{aligned}$$

The multiplied operator is the quadratic polynomial

$$\begin{aligned}E = (1 - c^2)\big(c^3\,\partial_{cc}^{\mathrm{cl}} - c^2\,\partial_c^{\mathrm{cl}}\big)\\ + 2c^2\big[\lambda + 1 - (1 - 2\alpha - \lambda)c^2\big]\,\partial_c^{\mathrm{cl}} - 4\alpha(\alpha + \lambda)c^3F^2,\end{aligned} \tag{11}$$

where the cleared derivatives are the two displays above. Their coefficients in t are

$$\begin{aligned}(c + t)F &= 2c\,t^2 + (1 + 3c^2)t + c(1 + c^2),\\ \partial_c^{\mathrm{cl}} &= 2\alpha\big(2c\,t^2 + (1 + 3c^2)t + c(1 + c^2)\big),\\ \partial_{cc}^{\mathrm{cl}} &= 4\alpha(\alpha - 1)\,t^2 + 4\alpha c(2\alpha - 1)\,t + 2\alpha\big(1 + (2\alpha - 1)c^2\big).\end{aligned}$$

The linear coefficient of $\partial_{cc}^{\mathrm{cl}}$ is $8\alpha(\alpha-1)c + 4\alpha c = 4\alpha c(2\alpha - 1)$, and its constant term is $4\alpha(\alpha-1)c^2 + 2\alpha(1 + c^2) = 2\alpha\big(1 + (2\alpha - 1)c^2\big)$. The square $F^2 = (1 + c^2 + 2ct)^2$ has coefficients $4c^2$, $4c(1 + c^2)$ and $(1 + c^2)^2$.

Coefficient of t^2. The combination $c^3\partial_{cc}^{\mathrm{cl}} - c^2\partial_c^{\mathrm{cl}}$ has t^2-coefficient $4\alpha c^3(\alpha-1) - 4\alpha c^3 = 4\alpha c^3(\alpha - 2)$. Multiplication by $1 - c^2$ contributes $4\alpha c^3(\alpha - 2)(1 - c^2)$. The middle summand of (11) contributes $8\alpha c^3\big[\lambda + 1 - (1 - 2\alpha - \lambda)c^2\big]$, and the last summand contributes $-16\alpha(\alpha + \lambda)c^5$. Dividing the sum by $4\alpha c^3$ produces

$$(\alpha - 2)(1 - c^2) + 2\big[\lambda + 1 - (1 - 2\alpha - \lambda)c^2\big] - 4(\alpha + \lambda)c^2.$$

The terms independent of c^2 are $\alpha - 2 + 2\lambda + 2 = \alpha + 2\lambda$. The coefficient of c^2 is

$$-(\alpha - 2) - 2(1 - 2\alpha - \lambda) - 4(\alpha + \lambda) = -\alpha + 2 - 2 + 4\alpha + 2\lambda - 4\alpha - 4\lambda = -(\alpha + 2\lambda).$$

The quotient equals $(\alpha + 2\lambda)(1 - c^2)$, so the coefficient of t^2 in E is $-4\alpha c^3(\alpha + 2\lambda)(c^2 - 1)$.

Coefficient of t. The three contributions are

$$\begin{aligned}&(1 - c^2)\big(4\alpha(2\alpha - 1)c^4 - 2\alpha c^2(1 + 3c^2)\big),\\ &4\alpha c^2(1 + 3c^2)\big[\lambda + 1 - (1 - 2\alpha - \lambda)c^2\big],\\ &-16\alpha(\alpha + \lambda)c^4(1 + c^2).\end{aligned}$$

Dividing by $2\alpha c^2$ produces

$$(1 - c^2)\big((4\alpha - 5)c^2 - 1\big) + 2(1 + 3c^2)\big[\lambda + 1 - (1 - 2\alpha - \lambda)c^2\big] - 8(\alpha + \lambda)c^2(1 + c^2).$$

The first summand equals $-(4\alpha - 5)c^4 + 4(\alpha - 1)c^2 - 1$. The second equals $2(\lambda + 1) + 6(\lambda + 1)c^2 - 2(1 - 2\alpha - \lambda)c^2 - 6(1 - 2\alpha - \lambda)c^4$. The third equals $-8(\alpha + \lambda)c^2 - 8(\alpha + \lambda)c^4$. The coefficient of c^4 in the sum is

$$\begin{aligned}&-(4\alpha - 5) - 6(1 - 2\alpha - \lambda) - 8(\alpha + \lambda)\\ &\quad = -4\alpha + 5 - 6 + 12\alpha + 6\lambda - 8\alpha - 8\lambda = -(2\lambda + 1).\end{aligned}$$

The coefficient of c^2 is

$$\begin{aligned}&4(\alpha - 1) + 6(\lambda + 1) - 2(1 - 2\alpha - \lambda) - 8(\alpha + \lambda)\\ &\quad = 4\alpha - 4 + 6\lambda + 6 - 2 + 4\alpha + 2\lambda - 8\alpha - 8\lambda = 0.\end{aligned}$$

The remaining term is $-1 + 2(\lambda + 1) = 2\lambda + 1$. The quotient is therefore $(2\lambda + 1)(1 - c^4) = (2\lambda + 1)(1 - c^2)(1 + c^2)$, and the coefficient of t in E is $-2\alpha c^2(c^2 - 1)(c^2 + 1)(2\lambda + 1)$.

Constant term. The three contributions are $4\alpha(\alpha - 1)c^5(1 - c^2)$, $4\alpha c^3(1 + c^2)\big[\lambda + 1 - (1 - 2\alpha - \lambda)c^2\big]$ and $-4\alpha(\alpha + \lambda)c^3(1 + c^2)^2$. Dividing by $4\alpha c^3$ produces

$$(\alpha - 1)c^2(1 - c^2) + (1 + c^2)\big[\lambda + 1 - (1 - 2\alpha - \lambda)c^2\big] - (\alpha + \lambda)(1 + c^2)^2.$$

The coefficient of c^4 is $-(\alpha - 1) - (1 - 2\alpha - \lambda) - (\alpha + \lambda) = 0$. The coefficient of c^2 is

$$(\alpha - 1) + (\lambda + 1) - (1 - 2\alpha - \lambda) - 2(\alpha + \lambda) = \alpha - 1 + \lambda + 1 - 1 + 2\alpha + \lambda - 2\alpha - 2\lambda = \alpha - 1,$$

and the remaining term is $(\lambda + 1) - (\alpha + \lambda) = 1 - \alpha$. The quotient equals $(\alpha - 1)(c^2 - 1)$, so the constant term of E is $4\alpha c^3(\alpha - 1)(c^2 - 1)$.

These are the coefficients of $-2\alpha c^2(c^2 - 1)Q$. Both sides are quadratic polynomials in t, so $E = -2\alpha c^2(c^2 - 1)Q$. The divisions by α and by powers of c compare coefficients of polynomials and do not restrict the identity: both sides vanish to at least those orders. Dividing by $4c^3F^{2-\alpha}$ gives (5).

The initial slope

Start from (9), which is the differentiation of (6) at $z = 0$. With $t_\alpha = (\alpha + \lambda)/(\lambda + 1)$,

$$\begin{aligned}(1 - 2\alpha - \lambda - \alpha t_\alpha)(\lambda + 1) &= (1 - 2\alpha - \lambda)(\lambda + 1) - \alpha(\alpha + \lambda)\\ &= 1 - \lambda^2 - 2\alpha - 2\alpha\lambda - \alpha^2 - \alpha\lambda = M(\alpha),\end{aligned}$$

where $M(\alpha) = (1-\lambda^2) - \alpha^2 - \alpha(3\lambda+2)$. Thus $1 - 2\alpha - \lambda - \alpha t_\alpha = M(\alpha)/(\lambda+1)$. With $\kappa t_\delta = t_\gamma$,

$$(1 - 2\gamma - \lambda - \gamma t_\gamma) - \kappa(1 - 2\delta - \lambda - \delta t_\delta) = \frac{M(\gamma)}{\lambda+1} - \frac{\gamma+\lambda}{\delta+\lambda} \cdot \frac{M(\delta)}{\lambda+1}.$$

The numerator of the common-denominator form is

$$(\delta+\lambda)M(\gamma) - (\gamma+\lambda)M(\delta).$$

The summand $(1-\lambda^2)$ contributes $(\delta-\gamma)(1-\lambda^2)$. The quadratic summand contributes

$$\begin{aligned} -(\delta+\lambda)\gamma^2 + (\gamma+\lambda)\delta^2 &= \gamma\delta(\delta-\gamma) + \lambda(\delta^2-\gamma^2) \\ &= (\delta-\gamma)\big(\gamma\delta + \lambda(\gamma+\delta)\big). \end{aligned}$$

The remaining summand contributes

$$(3\lambda+2)\big(-(\delta+\lambda)\gamma + (\gamma+\lambda)\delta\big) = (3\lambda+2)\big(-\lambda\gamma + \lambda\delta\big) = \lambda(3\lambda+2)(\delta-\gamma).$$

These are the three contributions

$$(1-\lambda^2)(\delta-\gamma), \qquad (\delta-\gamma)\big(\gamma\delta + \lambda(\gamma+\delta)\big), \qquad \lambda(3\lambda+2)(\delta-\gamma).$$

Their sum is

$$(\delta-\gamma)\big(1 + 2\lambda + 2\lambda^2 + \gamma\delta + \lambda(\gamma+\delta)\big) = (\delta-\gamma)\big((\gamma+\lambda)(\delta+\lambda) + (\lambda+1)^2\big).$$

Dividing by $(\delta+\lambda)(\lambda+1)$ and multiplying by $t_\gamma/(\lambda+2)$ gives

$$\tau'_\gamma(0) - \psi'(0) = \frac{(\delta-\gamma)(\gamma+\lambda)\big((\gamma+\lambda)(\delta+\lambda) + (\lambda+1)^2\big)}{(\delta+\lambda)(\lambda+1)^2(\lambda+2)}.$$

Coefficients of the contact polynomial

Write $P(t) = N_\gamma(t)(1-\kappa z) - N_\psi(t)(1-z)$ and

$$S = \frac{z(\delta-\gamma)}{\delta+\lambda}.$$

The constant terms contribute

$$(\gamma+\lambda)\big((1-\kappa z) - (1-z)\big) = (\gamma+\lambda)z(1-\kappa) = S(\gamma+\lambda),$$

since $1-\kappa = (\delta-\gamma)/(\delta+\lambda)$.

The coefficients of t^2 contribute

$$\begin{aligned} &- z(1-z)\gamma(1-\kappa z) + z(1-\kappa z)\delta(1-z) \\ &\quad = z(1-z)(1-\kappa z)(\delta-\gamma). \end{aligned}$$

The relation $1-\kappa z = (\delta+\lambda - z(\gamma+\lambda))/(\delta+\lambda)$ rewrites the right-hand side as SA_Q.

The coefficient of t in N_γ is $-\big[\lambda+1-(1-2\gamma-\lambda)z\big]$, and the coefficient of t in N_ψ is $-\big[\lambda+1-(1-2\delta-\lambda)\kappa z\big]$. Write $A = \lambda+1$, $G = 1-2\gamma-\lambda$ and $D = 1-2\delta-\lambda$. The coefficient of t in P is

$$\begin{aligned} &- (A - Gz)(1-\kappa z) + (A - D\kappa z)(1-z) \\ &\quad = -A + A\kappa z + Gz(1-\kappa z) + A(1-z) - D\kappa z(1-z) \\ &\quad = z\big(A(\kappa-1) + G(1-\kappa z) - \kappa D(1-z)\big) = zB, \end{aligned}$$

where

$$B = (\lambda+1)(\kappa-1) + (1-2\gamma-\lambda)(1-\kappa z) - \kappa(1-2\delta-\lambda)(1-z).$$

Multiply by $\delta+\lambda$ and use $\kappa(\delta+\lambda) = \gamma+\lambda$:

$$\begin{aligned}(\delta+\lambda)B = (\lambda+1)(\gamma-\delta) &+ (\delta+\lambda)(1-2\gamma-\lambda)(1-\kappa z)\\ &- (\gamma+\lambda)(1-2\delta-\lambda)(1-z).\end{aligned}$$

The right-hand side is affine in z. Its constant term is

$$\begin{aligned}K &= (\lambda+1)(\gamma-\delta) + (\delta+\lambda)(1-2\gamma-\lambda) - (\gamma+\lambda)(1-2\delta-\lambda)\\ &= \lambda\gamma - \lambda\delta + \gamma - \delta + \delta - 2\gamma\delta - \lambda\delta + \lambda - 2\gamma\lambda - \lambda^2\\ &\quad - \gamma + 2\gamma\delta + \gamma\lambda - \lambda + 2\delta\lambda + \lambda^2.\end{aligned}$$

The pairs $\lambda\gamma + \gamma\lambda - 2\gamma\lambda$, $-\lambda\delta - \lambda\delta + 2\delta\lambda$, $-2\gamma\delta + 2\gamma\delta$, $-\lambda^2 + \lambda^2$, $\lambda - \lambda$, $\gamma - \gamma$ and $-\delta + \delta$ each sum to zero, so $K = 0$.

The coefficient of z receives $-\kappa z$ from the second summand and $-(1-z)$ from the third:

$$\begin{aligned}&- (\delta+\lambda)(1-2\gamma-\lambda)\kappa + (\gamma+\lambda)(1-2\delta-\lambda)\\ &\quad = (\gamma+\lambda)\big(-(1-2\gamma-\lambda) + (1-2\delta-\lambda)\big)\\ &\quad = 2(\gamma+\lambda)(\gamma-\delta).\end{aligned}$$

Hence $(\delta+\lambda)B = 2z(\gamma+\lambda)(\gamma-\delta)$, and $zB = -2z(\gamma+\lambda)S$. Adding the three contributions yields $P(t) = S\,Q(t)$.

This is an AI-generated solution to a problem that was prompted by Paata Ivanisvili in Cursor with Grok 4.7.

Paata Ivanisvili
Department of Mathematics, University of California, Irvine
510C Rowland Hall, Irvine, CA 92697-3875, USA