

1 Maks $|b| > 10^\circ$

$30 < \ell < 120$

All pairs (auto+cross)

Table 1: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 120$).

Data	Mode	$A_s [10^{-3} \mu\text{K}^2]$	α_s	β_s	$A_d [10^{-3} \mu\text{K}^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$8.863^{+0.746}_{-0.734}$	$-3.059^{+0.177}_{-0.185}$	$-3.242^{+0.120}_{-0.125}$	$3.931^{+0.019}_{-0.019}$	$-1.998^{+0.010}_{-0.010}$	$1.563^{+0.007}_{-0.006}$	$0.075^{+0.010}_{-0.010}$	3.55
WMAP+Planck	BB	$1.912^{+0.604}_{-0.569}$	$-3.107^{+0.609}_{-0.741}$	$-3.339^{+0.312}_{-0.345}$	$3.032^{+0.013}_{-0.013}$	$-1.952^{+0.010}_{-0.010}$	$1.576^{+0.006}_{-0.006}$	$0.121^{+0.021}_{-0.019}$	1.79
QUIJOTE+WMAP+Planck	EE	$8.400^{+0.365}_{-0.362}$	$-3.072^{+0.096}_{-0.098}$	$-3.164^{+0.036}_{-0.036}$	$3.931^{+0.019}_{-0.019}$	$-1.998^{+0.010}_{-0.010}$	$1.563^{+0.007}_{-0.006}$	$0.071^{+0.007}_{-0.007}$	2.95
QUIJOTE+WMAP+Planck	BB	$0.786^{+0.332}_{-0.803}$	$-4.634^{+0.731}_{-0.664}$	$-3.030^{+0.496}_{-0.145}$	$3.032^{+0.013}_{-0.013}$	$-1.951^{+0.010}_{-0.010}$	$1.576^{+0.006}_{-0.006}$	$0.088^{+0.020}_{-0.950}$	1.64

No autos QUIJOTE

Table 2: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 120$).

Data	Mode	$A_s [10^{-3} \mu\text{K}^2]$	α_s	β_s	$A_d [10^{-3} \mu\text{K}^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$8.862^{+0.737}_{-0.723}$	$-3.057^{+0.175}_{-0.183}$	$-3.242^{+0.120}_{-0.123}$	$3.931^{+0.019}_{-0.019}$	$-1.998^{+0.011}_{-0.010}$	$1.563^{+0.007}_{-0.007}$	$0.075^{+0.011}_{-0.010}$	3.55
WMAP+Planck	BB	$1.909^{+0.601}_{-0.571}$	$-3.118^{+0.612}_{-0.734}$	$-3.337^{+0.315}_{-0.344}$	$3.032^{+0.013}_{-0.013}$	$-1.952^{+0.010}_{-0.010}$	$1.576^{+0.006}_{-0.006}$	$0.121^{+0.021}_{-0.019}$	1.79
QUIJOTE+WMAP+Planck	EE	$8.469^{+0.377}_{-0.375}$	$-3.047^{+0.101}_{-0.103}$	$-3.180^{+0.045}_{-0.045}$	$3.931^{+0.019}_{-0.019}$	$-1.998^{+0.010}_{-0.010}$	$1.563^{+0.007}_{-0.006}$	$0.070^{+0.007}_{-0.007}$	2.99
QUIJOTE+WMAP+Planck	BB	$1.092^{+0.301}_{-0.276}$	$-4.368^{+0.525}_{-0.613}$	$-2.946^{+0.121}_{-0.123}$	$3.032^{+0.013}_{-0.012}$	$-1.952^{+0.010}_{-0.010}$	$1.576^{+0.006}_{-0.006}$	$0.098^{+0.016}_{-0.015}$	1.64

Cross only

Table 3: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 120$).

Data	Mode	$A_s [10^{-3} \mu\text{K}^2]$	α_s	β_s	$A_d [10^{-3} \mu\text{K}^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$10.896^{+1.437}_{-1.309}$	$-2.942^{+0.212}_{-0.220}$	$-3.615^{+0.223}_{-0.246}$	$3.981^{+0.038}_{-0.037}$	$-1.996^{+0.014}_{-0.014}$	$1.584^{+0.011}_{-0.011}$	$0.076^{+0.010}_{-0.010}$	2.71
WMAP+Planck	BB	$3.015^{+1.047}_{-0.899}$	$-2.487^{+0.559}_{-0.630}$	$-3.689^{+0.470}_{-0.550}$	$3.142^{+0.028}_{-0.028}$	$-1.968^{+0.014}_{-0.014}$	$1.628^{+0.011}_{-0.011}$	$0.114^{+0.020}_{-0.018}$	1.55
QUIJOTE+WMAP+Planck	EE	$8.506^{+0.414}_{-0.409}$	$-3.034^{+0.107}_{-0.112}$	$-3.183^{+0.047}_{-0.048}$	$3.987^{+0.037}_{-0.037}$	$-1.996^{+0.014}_{-0.014}$	$1.586^{+0.011}_{-0.011}$	$0.072^{+0.007}_{-0.007}$	2.33
QUIJOTE+WMAP+Planck	BB	$1.237^{+0.314}_{-0.305}$	$-4.148^{+0.492}_{-0.581}$	$-2.893^{+0.121}_{-0.124}$	$3.144^{+0.028}_{-0.027}$	$-1.966^{+0.014}_{-0.014}$	$1.629^{+0.011}_{-0.011}$	$0.102^{+0.016}_{-0.015}$	1.46

$30 < \ell < 200$

All pairs (auto+cross)

Table 4: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 200$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$9.486^{+0.647}_{-0.646}$	$-2.848^{+0.148}_{-0.156}$	$-3.201^{+0.115}_{-0.121}$	$3.751^{+0.016}_{-0.016}$	$-2.242^{+0.006}_{-0.006}$	$1.565^{+0.006}_{-0.006}$	$0.081^{+0.010}_{-0.010}$	4.05
WMAP+Planck	BB	$2.634^{+0.542}_{-0.542}$	$-2.379^{+0.449}_{-0.517}$	$-3.488^{+0.295}_{-0.327}$	$2.911^{+0.011}_{-0.011}$	$-2.162^{+0.006}_{-0.006}$	$1.572^{+0.005}_{-0.005}$	$0.112^{+0.019}_{-0.017}$	3.12
QUIJOTE+WMAP+Planck	EE	$8.805^{+0.324}_{-0.330}$	$-2.951^{+0.081}_{-0.082}$	$-3.150^{+0.036}_{-0.035}$	$3.751^{+0.016}_{-0.016}$	$-2.242^{+0.006}_{-0.006}$	$1.565^{+0.006}_{-0.006}$	$0.079^{+0.007}_{-0.007}$	3.31
QUIJOTE+WMAP+Planck	BB	$1.246^{+0.295}_{-0.282}$	$-4.104^{+0.456}_{-0.536}$	$-3.083^{+0.109}_{-0.109}$	$2.910^{+0.011}_{-0.011}$	$-2.162^{+0.006}_{-0.006}$	$1.572^{+0.005}_{-0.005}$	$0.092^{+0.014}_{-0.014}$	2.66

No autos QUIJOTE

Table 5: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 200$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$9.478^{+0.650}_{-0.644}$	$-2.848^{+0.146}_{-0.156}$	$-3.201^{+0.115}_{-0.121}$	$3.751^{+0.016}_{-0.016}$	$-2.242^{+0.006}_{-0.006}$	$1.565^{+0.006}_{-0.006}$	$0.081^{+0.010}_{-0.010}$	4.05
WMAP+Planck	BB	$2.625^{+0.541}_{-0.537}$	$-2.385^{+0.446}_{-0.511}$	$-3.481^{+0.294}_{-0.323}$	$2.911^{+0.011}_{-0.011}$	$-2.162^{+0.006}_{-0.006}$	$1.572^{+0.005}_{-0.005}$	$0.112^{+0.018}_{-0.017}$	3.12
QUIJOTE+WMAP+Planck	EE	$8.926^{+0.337}_{-0.337}$	$-2.905^{+0.084}_{-0.087}$	$-3.168^{+0.044}_{-0.044}$	$3.751^{+0.016}_{-0.016}$	$-2.242^{+0.006}_{-0.006}$	$1.565^{+0.006}_{-0.006}$	$0.079^{+0.007}_{-0.007}$	3.35
QUIJOTE+WMAP+Planck	BB	$1.538^{+0.312}_{-0.311}$	$-3.597^{+0.421}_{-0.491}$	$-2.948^{+0.116}_{-0.118}$	$2.910^{+0.011}_{-0.011}$	$-2.162^{+0.006}_{-0.006}$	$1.572^{+0.005}_{-0.005}$	$0.091^{+0.014}_{-0.014}$	2.67

Cross only

Table 6: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 200$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$11.582^{+1.179}_{-1.113}$	$-2.664^{+0.162}_{-0.172}$	$-3.492^{+0.200}_{-0.222}$	$3.795^{+0.032}_{-0.031}$	$-2.239^{+0.008}_{-0.008}$	$1.587^{+0.010}_{-0.010}$	$0.081^{+0.009}_{-0.009}$	3.15
WMAP+Planck	BB	$3.615^{+0.991}_{-0.853}$	$-2.125^{+0.399}_{-0.456}$	$-3.814^{+0.459}_{-0.539}$	$3.050^{+0.024}_{-0.024}$	$-2.160^{+0.008}_{-0.008}$	$1.638^{+0.010}_{-0.010}$	$0.107^{+0.018}_{-0.016}$	2.28
QUIJOTE+WMAP+Planck	EE	$9.078^{+0.362}_{-0.364}$	$-2.866^{+0.088}_{-0.092}$	$-3.162^{+0.047}_{-0.046}$	$3.801^{+0.032}_{-0.032}$	$-2.239^{+0.008}_{-0.008}$	$1.589^{+0.010}_{-0.010}$	$0.080^{+0.007}_{-0.007}$	2.62
QUIJOTE+WMAP+Planck	BB	$1.684^{+0.313}_{-0.312}$	$-3.450^{+0.390}_{-0.444}$	$-2.859^{+0.118}_{-0.120}$	$3.052^{+0.024}_{-0.023}$	$-2.159^{+0.008}_{-0.008}$	$1.640^{+0.010}_{-0.010}$	$0.099^{+0.015}_{-0.014}$	2.01

2 Maks $|b| > 15^\circ$

$30 < \ell < 120$

All pairs (auto+cross)

Table 7: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 120$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$8.308^{+0.778}_{-0.770}$	$-2.868^{+0.202}_{-0.215}$	$-3.304^{+0.143}_{-0.151}$	$2.477^{+0.017}_{-0.017}$	$-1.808^{+0.016}_{-0.016}$	$1.575^{+0.009}_{-0.009}$	$0.059^{+0.012}_{-0.012}$	1.36
WMAP+Planck	BB	$2.029^{+0.630}_{-0.606}$	$-2.729^{+0.638}_{-0.783}$	$-3.261^{+0.351}_{-0.401}$	$1.815^{+0.011}_{-0.011}$	$-2.363^{+0.014}_{-0.014}$	$1.586^{+0.008}_{-0.008}$	$0.085^{+0.021}_{-0.020}$	1.49
QUIJOTE+WMAP+Planck	EE	$7.612^{+0.368}_{-0.369}$	$-2.903^{+0.110}_{-0.112}$	$-3.147^{+0.042}_{-0.042}$	$2.476^{+0.017}_{-0.017}$	$-1.808^{+0.015}_{-0.015}$	$1.574^{+0.009}_{-0.009}$	$0.039^{+0.009}_{-0.009}$	1.34
QUIJOTE+WMAP+Planck	BB	$0.936^{+0.307}_{-0.268}$	$-4.507^{+0.604}_{-0.703}$	$-3.030^{+0.133}_{-0.135}$	$1.815^{+0.011}_{-0.011}$	$-2.362^{+0.014}_{-0.014}$	$1.585^{+0.008}_{-0.008}$	$0.055^{+0.017}_{-0.016}$	1.51

No autos QUIJOTE

Table 8: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 120$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$8.323^{+0.781}_{-0.770}$	$-2.864^{+0.201}_{-0.213}$	$-3.306^{+0.143}_{-0.150}$	$2.477^{+0.017}_{-0.017}$	$-1.808^{+0.016}_{-0.016}$	$1.575^{+0.010}_{-0.009}$	$0.059^{+0.012}_{-0.012}$	1.36
WMAP+Planck	BB	$2.024^{+0.634}_{-0.602}$	$-2.733^{+0.641}_{-0.781}$	$-3.259^{+0.352}_{-0.402}$	$1.815^{+0.011}_{-0.011}$	$-2.362^{+0.014}_{-0.014}$	$1.586^{+0.008}_{-0.008}$	$0.085^{+0.021}_{-0.021}$	1.49
QUIJOTE+WMAP+Planck	EE	$7.684^{+0.378}_{-0.380}$	$-2.872^{+0.115}_{-0.119}$	$-3.163^{+0.053}_{-0.053}$	$2.476^{+0.017}_{-0.017}$	$-1.808^{+0.016}_{-0.015}$	$1.574^{+0.009}_{-0.009}$	$0.039^{+0.009}_{-0.009}$	1.36
QUIJOTE+WMAP+Planck	BB	$1.148^{+0.348}_{-0.330}$	$-4.033^{+0.584}_{-0.725}$	$-2.883^{+0.142}_{-0.147}$	$1.815^{+0.011}_{-0.011}$	$-2.362^{+0.014}_{-0.014}$	$1.585^{+0.008}_{-0.008}$	$0.057^{+0.017}_{-0.016}$	1.50

Cross only

Table 9: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 120$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$11.425^{+1.703}_{-1.524}$	$-2.601^{+0.231}_{-0.246}$	$-3.891^{+0.292}_{-0.342}$	$2.529^{+0.034}_{-0.033}$	$-1.806^{+0.021}_{-0.021}$	$1.607^{+0.016}_{-0.016}$	$0.063^{+0.011}_{-0.011}$	1.19
WMAP+Planck	BB	$3.406^{+1.255}_{-1.039}$	$-2.079^{+0.620}_{-0.684}$	$-3.991^{+0.626}_{-0.804}$	$1.857^{+0.023}_{-0.022}$	$-2.383^{+0.018}_{-0.019}$	$1.622^{+0.015}_{-0.015}$	$0.081^{+0.020}_{-0.018}$	1.36
QUIJOTE+WMAP+Planck	EE	$7.749^{+0.408}_{-0.406}$	$-2.825^{+0.121}_{-0.125}$	$-3.173^{+0.056}_{-0.056}$	$2.530^{+0.033}_{-0.033}$	$-1.805^{+0.020}_{-0.021}$	$1.608^{+0.016}_{-0.016}$	$0.041^{+0.009}_{-0.009}$	1.25
QUIJOTE+WMAP+Planck	BB	$1.249^{+0.359}_{-0.346}$	$-3.860^{+0.568}_{-0.696}$	$-2.874^{+0.144}_{-0.149}$	$1.857^{+0.022}_{-0.022}$	$-2.382^{+0.018}_{-0.018}$	$1.623^{+0.015}_{-0.015}$	$0.061^{+0.017}_{-0.017}$	1.42

$30 < \ell < 200$

All pairs (auto+cross)

Table 10: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 200$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$8.718^{+0.680}_{-0.678}$	$-2.699^{+0.172}_{-0.181}$	$-3.246^{+0.139}_{-0.144}$	$2.352^{+0.015}_{-0.015}$	$-2.108^{+0.008}_{-0.009}$	$1.572^{+0.008}_{-0.008}$	$0.059^{+0.011}_{-0.011}$	2.26
WMAP+Planck	BB	$2.713^{+0.553}_{-0.553}$	$-1.946^{+0.500}_{-0.592}$	$-3.296^{+0.336}_{-0.385}$	$1.784^{+0.009}_{-0.009}$	$-2.417^{+0.009}_{-0.009}$	$1.583^{+0.008}_{-0.008}$	$0.078^{+0.019}_{-0.018}$	1.73
QUIJOTE+WMAP+Planck	EE	$7.874^{+0.332}_{-0.331}$	$-2.818^{+0.095}_{-0.095}$	$-3.131^{+0.041}_{-0.041}$	$2.352^{+0.015}_{-0.015}$	$-2.108^{+0.009}_{-0.009}$	$1.572^{+0.008}_{-0.008}$	$0.046^{+0.008}_{-0.008}$	1.97
QUIJOTE+WMAP+Planck	BB	$1.247^{+0.370}_{-0.355}$	$-3.891^{+0.574}_{-0.712}$	$-2.997^{+0.133}_{-0.135}$	$1.783^{+0.009}_{-0.009}$	$-2.417^{+0.009}_{-0.009}$	$1.582^{+0.008}_{-0.008}$	$0.054^{+0.016}_{-0.015}$	1.64

No autos QUIJOTE

Table 11: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 200$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$8.707^{+0.679}_{-0.672}$	$-2.697^{+0.169}_{-0.182}$	$-3.241^{+0.137}_{-0.145}$	$2.352^{+0.015}_{-0.015}$	$-2.109^{+0.009}_{-0.009}$	$1.572^{+0.008}_{-0.008}$	$0.059^{+0.011}_{-0.011}$	2.26
WMAP+Planck	BB	$2.510^{+0.670}_{-2.542}$	$-1.868^{+0.593}_{-0.630}$	$-3.396^{+0.399}_{-0.917}$	$1.784^{+0.009}_{-0.009}$	$-2.417^{+0.009}_{-0.009}$	$1.583^{+0.008}_{-0.008}$	$0.072^{+0.022}_{-0.548}$	1.73
QUIJOTE+WMAP+Planck	EE	$7.988^{+0.339}_{-0.340}$	$-2.769^{+0.097}_{-0.099}$	$-3.146^{+0.052}_{-0.052}$	$2.352^{+0.015}_{-0.014}$	$-2.108^{+0.009}_{-0.008}$	$1.572^{+0.008}_{-0.008}$	$0.046^{+0.008}_{-0.008}$	2.00
QUIJOTE+WMAP+Planck	BB	$1.622^{+0.377}_{-0.388}$	$-3.247^{+0.506}_{-0.620}$	$-2.856^{+0.140}_{-0.144}$	$1.783^{+0.009}_{-0.009}$	$-2.417^{+0.009}_{-0.009}$	$1.583^{+0.008}_{-0.008}$	$0.055^{+0.016}_{-0.015}$	1.64

Cross only

Table 12: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 200$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$11.568^{+1.427}_{-1.245}$	$-2.402^{+0.178}_{-0.194}$	$-3.698^{+0.261}_{-0.307}$	$2.391^{+0.029}_{-0.028}$	$-2.108^{+0.011}_{-0.011}$	$1.600^{+0.015}_{-0.014}$	$0.061^{+0.010}_{-0.011}$	1.92
WMAP+Planck	BB	$4.032^{+1.231}_{-1.010}$	$-1.670^{+0.465}_{-0.518}$	$-4.074^{+0.623}_{-0.778}$	$1.850^{+0.020}_{-0.020}$	$-2.420^{+0.012}_{-0.012}$	$1.636^{+0.014}_{-0.014}$	$0.075^{+0.018}_{-0.016}$	1.51
QUIJOTE+WMAP+Planck	EE	$8.141^{+0.370}_{-0.369}$	$-2.700^{+0.100}_{-0.104}$	$-3.146^{+0.055}_{-0.056}$	$2.393^{+0.029}_{-0.028}$	$-2.108^{+0.011}_{-0.011}$	$1.601^{+0.014}_{-0.014}$	$0.048^{+0.008}_{-0.008}$	1.73
QUIJOTE+WMAP+Planck	BB	$1.706^{+0.360}_{-0.380}$	$-3.145^{+0.469}_{-0.580}$	$-2.830^{+0.141}_{-0.147}$	$1.852^{+0.020}_{-0.020}$	$-2.419^{+0.012}_{-0.012}$	$1.638^{+0.014}_{-0.014}$	$0.060^{+0.016}_{-0.015}$	1.47

3 Maks $|b| > 20^\circ$

$30 < \ell < 120$

All pairs (auto+cross)

Table 13: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 120$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$7.311^{+0.785}_{-0.780}$	$-3.185^{+0.228}_{-0.247}$	$-3.298^{+0.149}_{-0.157}$	$1.759^{+0.015}_{-0.015}$	$-2.036^{+0.020}_{-0.020}$	$1.534^{+0.012}_{-0.012}$	$0.046^{+0.013}_{-0.013}$	1.99
WMAP+Planck	BB	$1.866^{+0.663}_{-0.618}$	$-2.668^{+0.708}_{-0.875}$	$-3.041^{+0.364}_{-0.423}$	$1.193^{+0.009}_{-0.009}$	$-2.338^{+0.018}_{-0.018}$	$1.578^{+0.012}_{-0.011}$	$0.085^{+0.025}_{-0.024}$	1.47
QUIJOTE+WMAP+Planck	EE	$6.806^{+0.372}_{-0.372}$	$-3.157^{+0.122}_{-0.127}$	$-3.149^{+0.045}_{-0.045}$	$1.758^{+0.015}_{-0.015}$	$-2.035^{+0.020}_{-0.020}$	$1.532^{+0.012}_{-0.012}$	$0.021^{+0.009}_{-0.009}$	1.87
QUIJOTE+WMAP+Planck	BB	$0.998^{+0.339}_{-0.303}$	$-4.242^{+0.628}_{-0.757}$	$-2.977^{+0.148}_{-0.147}$	$1.192^{+0.009}_{-0.009}$	$-2.338^{+0.018}_{-0.018}$	$1.577^{+0.011}_{-0.011}$	$0.059^{+0.020}_{-0.019}$	1.46

No autos QUIJOTE

Table 14: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 120$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$7.316^{+0.789}_{-0.785}$	$-3.184^{+0.230}_{-0.249}$	$-3.299^{+0.149}_{-0.156}$	$1.759^{+0.015}_{-0.015}$	$-2.035^{+0.020}_{-0.020}$	$1.534^{+0.012}_{-0.012}$	$0.046^{+0.013}_{-0.013}$	1.99
WMAP+Planck	BB	$1.887^{+0.673}_{-0.633}$	$-2.653^{+0.706}_{-0.881}$	$-3.043^{+0.364}_{-0.428}$	$1.193^{+0.009}_{-0.009}$	$-2.339^{+0.018}_{-0.018}$	$1.578^{+0.011}_{-0.011}$	$0.085^{+0.025}_{-0.024}$	1.47
QUIJOTE+WMAP+Planck	EE	$6.870^{+0.386}_{-0.389}$	$-3.128^{+0.128}_{-0.134}$	$-3.156^{+0.055}_{-0.056}$	$1.758^{+0.015}_{-0.015}$	$-2.035^{+0.020}_{-0.020}$	$1.532^{+0.012}_{-0.012}$	$0.021^{+0.009}_{-0.009}$	1.89
QUIJOTE+WMAP+Planck	BB	$1.225^{+0.373}_{-0.360}$	$-3.753^{+0.609}_{-0.756}$	$-2.844^{+0.161}_{-0.164}$	$1.193^{+0.009}_{-0.009}$	$-2.338^{+0.018}_{-0.018}$	$1.578^{+0.011}_{-0.011}$	$0.061^{+0.020}_{-0.019}$	1.46

Cross only

Table 15: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 120$).

Data	Mode	$A_s [10^{-3} \mu K^2]$	α_s	β_s	$A_d [10^{-3} \mu K^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$10.479^{+1.754}_{-1.559}$	$-2.941^{+0.262}_{-0.283}$	$-3.995^{+0.330}_{-0.387}$	$1.760^{+0.029}_{-0.029}$	$-2.054^{+0.027}_{-0.027}$	$1.542^{+0.020}_{-0.020}$	$0.048^{+0.012}_{-0.012}$	1.57
WMAP+Planck	BB	$2.962^{+1.328}_{-1.065}$	$-2.105^{+0.711}_{-0.767}$	$-3.728^{+0.665}_{-0.882}$	$1.216^{+0.021}_{-0.020}$	$-2.355^{+0.026}_{-0.025}$	$1.609^{+0.022}_{-0.021}$	$0.081^{+0.023}_{-0.022}$	1.42
QUIJOTE+WMAP+Planck	EE	$6.924^{+0.419}_{-0.419}$	$-3.082^{+0.134}_{-0.143}$	$-3.169^{+0.058}_{-0.059}$	$1.759^{+0.029}_{-0.028}$	$-2.053^{+0.026}_{-0.027}$	$1.541^{+0.020}_{-0.019}$	$0.021^{+0.009}_{-0.009}$	1.59
QUIJOTE+WMAP+Planck	BB	$1.298^{+0.386}_{-0.366}$	$-3.638^{+0.600}_{-0.705}$	$-2.851^{+0.164}_{-0.167}$	$1.216^{+0.020}_{-0.020}$	$-2.354^{+0.025}_{-0.025}$	$1.610^{+0.022}_{-0.021}$	$0.063^{+0.020}_{-0.020}$	1.43

$30 < \ell < 200$

All pairs (auto+cross)

Table 16: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 200$).

Data	Mode	$A_s [10^{-3} \mu\text{K}^2]$	α_s	β_s	$A_d [10^{-3} \mu\text{K}^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$7.912^{+0.719}_{-0.720}$	$-2.954^{+0.201}_{-0.215}$	$-3.263^{+0.146}_{-0.153}$	$1.697^{+0.013}_{-0.013}$	$-2.203^{+0.011}_{-0.011}$	$1.526^{+0.010}_{-0.010}$	$0.050^{+0.012}_{-0.012}$	1.81
WMAP+Planck	BB	$2.712^{+0.593}_{-0.615}$	$-1.659^{+0.551}_{-0.674}$	$-3.112^{+0.351}_{-0.406}$	$1.190^{+0.008}_{-0.008}$	$-2.334^{+0.011}_{-0.011}$	$1.579^{+0.010}_{-0.010}$	$0.070^{+0.022}_{-0.021}$	1.54
QUIJOTE+WMAP+Planck	EE	$7.218^{+0.352}_{-0.345}$	$-3.010^{+0.107}_{-0.111}$	$-3.134^{+0.044}_{-0.044}$	$1.696^{+0.013}_{-0.013}$	$-2.203^{+0.011}_{-0.011}$	$1.525^{+0.010}_{-0.010}$	$0.030^{+0.009}_{-0.009}$	1.68
QUIJOTE+WMAP+Planck	BB	$1.359^{+0.410}_{-0.399}$	$-3.577^{+0.605}_{-0.750}$	$-2.931^{+0.148}_{-0.149}$	$1.190^{+0.008}_{-0.008}$	$-2.334^{+0.011}_{-0.011}$	$1.578^{+0.010}_{-0.010}$	$0.058^{+0.019}_{-0.018}$	1.47

No autos QUIJOTE

Table 17: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 200$).

Data	Mode	$A_s [10^{-3} \mu\text{K}^2]$	α_s	β_s	$A_d [10^{-3} \mu\text{K}^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$7.913^{+0.717}_{-0.723}$	$-2.955^{+0.199}_{-0.219}$	$-3.264^{+0.146}_{-0.150}$	$1.697^{+0.013}_{-0.013}$	$-2.203^{+0.011}_{-0.011}$	$1.526^{+0.010}_{-0.010}$	$0.050^{+0.012}_{-0.012}$	1.81
WMAP+Planck	BB	$2.696^{+0.600}_{-0.655}$	$-1.653^{+0.552}_{-0.689}$	$-3.117^{+0.358}_{-0.429}$	$1.190^{+0.008}_{-0.008}$	$-2.334^{+0.011}_{-0.012}$	$1.579^{+0.010}_{-0.010}$	$0.070^{+0.023}_{-0.022}$	1.54
QUIJOTE+WMAP+Planck	EE	$7.348^{+0.356}_{-0.359}$	$-2.951^{+0.112}_{-0.117}$	$-3.144^{+0.054}_{-0.055}$	$1.696^{+0.013}_{-0.013}$	$-2.203^{+0.011}_{-0.011}$	$1.525^{+0.010}_{-0.010}$	$0.030^{+0.009}_{-0.009}$	1.69
QUIJOTE+WMAP+Planck	BB	$1.781^{+0.385}_{-0.416}$	$-2.867^{+0.509}_{-0.637}$	$-2.804^{+0.153}_{-0.160}$	$1.190^{+0.008}_{-0.008}$	$-2.334^{+0.011}_{-0.011}$	$1.579^{+0.010}_{-0.010}$	$0.056^{+0.018}_{-0.018}$	1.47

Cross only

Table 18: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes and spectral indices, and on the synchrotron–dust correlation coefficient ρ , from fits to the EE and BB spectra using the 10^0 mask (multipole range $\ell = 30 - 200$).

Data	Mode	$A_s [10^{-3} \mu\text{K}^2]$	α_s	β_s	$A_d [10^{-3} \mu\text{K}^2]$	α_d	β_d	ρ	χ^2_{red}
WMAP+Planck	EE	$11.209^{+1.557}_{-1.374}$	$-2.651^{+0.214}_{-0.229}$	$-3.870^{+0.299}_{-0.349}$	$1.700^{+0.025}_{-0.024}$	$-2.209^{+0.015}_{-0.015}$	$1.534^{+0.018}_{-0.017}$	$0.051^{+0.011}_{-0.011}$	1.56
WMAP+Planck	BB	$3.935^{+1.386}_{-1.157}$	$-1.456^{+0.547}_{-0.641}$	$-4.007^{+0.736}_{-0.907}$	$1.234^{+0.019}_{-0.018}$	$-2.337^{+0.016}_{-0.016}$	$1.630^{+0.020}_{-0.019}$	$0.071^{+0.020}_{-0.018}$	1.44
QUIJOTE+WMAP+Planck	EE	$7.496^{+0.399}_{-0.384}$	$-2.881^{+0.119}_{-0.123}$	$-3.149^{+0.058}_{-0.058}$	$1.700^{+0.026}_{-0.025}$	$-2.210^{+0.015}_{-0.015}$	$1.535^{+0.018}_{-0.017}$	$0.030^{+0.009}_{-0.009}$	1.51
QUIJOTE+WMAP+Planck	BB	$1.767^{+0.383}_{-0.421}$	$-2.889^{+0.492}_{-0.625}$	$-2.801^{+0.160}_{-0.164}$	$1.235^{+0.019}_{-0.018}$	$-2.336^{+0.016}_{-0.016}$	$1.633^{+0.020}_{-0.019}$	$0.061^{+0.019}_{-0.018}$	1.39