



Universidad de La Laguna
Instituto de Astrofísica de Canarias

Trabajo fin de Master

SPECTRAL PROPERTIES OF POLARIZED SYNCHROTRON AND THERMAL DUST EMISSION FROM POWER SPECTRUM ANALYSES WITH QUIJOTE, WMAP AND PLANCK DATA

Pablo de Ferra Chermaz

Supervisores:

José Alberto Rubiño Martín (IAC)

Resumen

CONTENTS

1	INTRODUCTION	3
2	OBJECTIVES	4
3	DATA	5
3.1.	Q-U-I JOint TEnerife (QUIJOTE)	5
3.2.	Wilkinson Microwave Anisotropy Probe (WMAP)	6
3.3.	Planck	6
3.3.1	Low Frequency Instrument (LFI)	6
3.3.2	High Frequency Instrument (HFI)	6
4	METHODOLOGY	6
4.1.	Angular power spectrum (APS) analysis	6
4.2.	Masks	7
4.3.	Data corrections	8
4.3.1	Beam and pixel window functions	8
4.3.2	Unit conversion	8
4.3.3	Noise power spectrum	9
4.3.4	Colour corrections	9
4.4.	Simulations	11
4.4.1	Sky signal simulations	11
4.4.2	Noise simulations	11
4.5.	Foreground emission model	12
4.5.1	Synchrotron emission	12
4.5.2	Thermal dust emission	13
4.5.3	Synchrotron-dust cross-correlation	13
4.5.4	CMB subtraction	13
4.6.	Bayesian inference and MCMC fitting	13
4.6.1	Likelihood	14
4.6.2	Priors	14
4.6.3	Sampler	14
4.7.	Fitting approaches	14
4.7.1	Power-law fitting	15
4.7.2	Bin-to-bin analysis	15
5	RESULTS	15
5.1.	Power-law fitting	15
5.2.	Bin-to-bin fitting	16
6	CONCLUSIONS	20
	REFERENCES	20
A	Analysis fitting only cross-spectra	22
1.1.	Power-law fitting	22
1.2.	Bin-to-bin fitting	22
B	Alternative multipole fitting range $30 \leq \ell \leq 199$	26
2.1.	Fitting all data (auto- and cross-spectra)	26
2.2.	Fitting only cross-spectra	26
C	Reduced χ^2 values of the power-law fittings	27

1 INTRODUCTION

The Cosmic Microwave Background (CMB) provides a unique window to probe the early stages of the Universe and its structure formation. Furthermore, CMB polarization enables to probe the amplitude of primordial Gravitational Waves (GWs; Ade et al., 2021) generated during the inflationary epoch (Kamionkowski, 1999; Kosowsky, 1996). These GWs give rise to the so-called B-mode of CMB polarization. In recent years, many experiments have been observing the CMB anisotropies. For instance, the Wilkinson Microwave Anisotropy Probe (WMAP; Hinshaw et al., 2013), scanning the full sky in five frequency bands between 23 and 94 GHz, and the Planck satellite (Planck Collaboration, 2020) in 9 bands, with polarization measurements in 7 of them, spanning from 30 to 353 GHz. These observations have significantly advanced our understanding of the Universe and motivated further scientific exploration, marking the beginning of a new era of high-precision cosmology with future missions centred in the study of the CMB, such as LiteBIRD (Allys et al., 2022) or other cosmology experiments such as the Square Kilometre Array (SKA; Huynh & Lazio, 2013), offering enhanced sensitivity and resolution.

The primordial B-mode signal might have an amplitude high enough to be within reach of current and upcoming experiments. However, its detection does not depend solely on instrumental sensitivity. A successful discovery critically relies on accurate component separation, since the cosmological signal is contaminated by polarized foreground emission from the Galaxy with significantly higher amplitude (Ade et al., 2021; Planck Collaboration, 2015). Component separation is therefore a central challenge in the design of future CMB experiments such as LiteBIRD (Allys et al., 2022), directly linking the search for primordial B modes to the statistical characterization and astrophysical understanding of polarized emission from the magnetized interstellar medium (ISM).

Microwave observations also contain several astrophysical emission mechanisms other than the CMB, collectively known as foregrounds. Because the primordial B-mode signal has a very low amplitude, the removal of these foregrounds in polarization has become a crucial step for its detection. Two foreground components are known to produce strongly linearly polarized radiation: synchrotron emission, generated by relativistic cosmic-ray electrons spiraling around Galactic magnetic field (GMF) lines and dominating at low frequencies; and thermal dust emission from interstellar dust grains aligned with the magnetic field, which dominates at high frequencies. Both components can exhibit polarization fractions of up to $\sim 20\%$ in certain regions of the sky (Planck Collaboration, 2015).

Anomalous microwave emission (AME), on the other hand, is observed to have a very low polarization fraction (G nova-Santos et al., 2017). If AME originates from spinning dust grains, theoretical models predict negligible polarization levels (Draine et al., 2016; Herman, D. et al., 2023). Another relevant foreground is free-free emission, produced by free electrons interacting with ionized gas; however, this mechanism generates radiation that is effectively unpolarized (Gold et al., 2011).

Synchrotron emission accounts for most of the radio emission from Active Galactic Nuclei (AGNs), which are thought to be powered by supermassive black holes in galaxies and quasars (Blandford & Znajek, 1977; Ishibashi, W. & Courvoisier, T. J.-L., 2011). This emission also dominates the radio continuum from star-forming galaxies, such as our own, at frequencies below $\nu \sim 30$ GHz. The relativistic electrons in nearly all synchrotron sources typically exhibit power-law energy distributions. The energy flux distribution for synchrotron radiation is given by $S_\nu \propto \nu^\alpha$, or equivalently, in terms of the brightness temperature as $T_\nu \propto \nu^\beta$, where $\beta = \alpha - 2$ (Rybicki & Lightman, 1979). This frequency dependence of the brightness temperature will be referred to as the spectral index. The spectral index depends on the energy-independent spectral energy slope s of the cosmic-ray electron spectrum, which is typically assumed to have a value of $s = -3$ in the frequency range from 10 to 30 GHz, but varying at lower frequencies (Padovani et al., 2021). Spatial variations of β reflect spatial variations of the cosmic-ray electrons and magnetic field properties in the interstellar medium along the line of sight. Full-sky maps of Galactic synchrotron

emission have already been derived in [Krachmalnicoff et al. \(2018\)](#) or [Rubiño-Martín et al. \(2023\)](#), and their analyses clearly show spatial variations of the spectral index, as discussed below.

At higher frequencies, the dominant polarized foreground is thermal emission from interstellar dust grains aligned with the GMF. This emission dominates the microwave sky above approximately 100 GHz and is partially linearly polarized. The frequency spectrum of thermal dust emission is well described by a modified black body (grey body), $I_d(\nu) \propto \nu^{\beta_d} B_\nu(T_d)$, where $B_\nu(T_d)$ is the Planck black-body spectrum at dust temperature T_d . Using combined Planck and WMAP data, [Planck Collaboration et al. \(2015\)](#) determined the dust spectral index in both intensity and polarization, finding $\beta_d \simeq 1.59$ for $T_d \simeq 19.6$ K, while [?](#) derived a mean polarization spectral index of $\beta_d = 1.53 \pm 0.02$ from PR3 maps at 353 GHz, with the dust EE and BB power spectra following power laws in multipole and exhibiting a significant E/B asymmetry. Since both dust and synchrotron polarization are aligned by the same GMF, a spatial cross-correlation between the two components is expected and has indeed been detected ([Choi & Page, 2015](#); [?](#)).

The sensitivities of WMAP and Planck alone do not allow a sufficiently accurate characterization of the polarized synchrotron signal at intermediate and high Galactic latitudes, leaving important uncertainties in the modelling of this foreground component. This is significantly improved by the QUIJOTE experiment ([Rubiño-Martín et al., 2023](#)), which observes the polarized Northern sky in the 10–40 GHz range, precisely where synchrotron emission is strongest and where Faraday rotation effects are not expected to be significant ([Fuskeland et al., 2021](#)). Combining QUIJOTE with WMAP and Planck data, [de la Hoz et al. \(2023\)](#) performed one of the first reconstructions of the polarized synchrotron spectral index in real space, finding statistically significant spatial variations across the sky, with an average value of $\beta = -3.08$ and a dispersion of 0.13. More recently, [Casas et al. \(2025\)](#) studied the spatial variability of polarized synchrotron emission using QUIJOTE MFI data alone, reinforcing the evidence for spatial variability in polarized synchrotron emission and highlighting the importance of combining QUIJOTE observations with complementary data from WMAP, Planck, and future low-frequency surveys in order to reduce noise contamination and improve the robustness of spectral-index estimates.

In this work, we characterize the spectral properties of polarized synchrotron and thermal dust emission through power spectrum analyses of the combined QUIJOTE MFI DR1 (11, 13, 17, and 19 GHz), WMAP 9-year (K, Ka, Q, V, and W bands), and Planck PR3 (LFI and HFI, 28.4–353 GHz) datasets. Both foreground components are modelled simultaneously by fitting a parametric SED model to the EE and BB angular power spectra (APS) and their cross-spectra across this wide frequency range. A key aspect of this analysis is the explicit treatment of the spatial cross-correlation between polarized synchrotron and thermal dust emission. Their cross-spectra encode complementary information about the geometry and coherence of the Galactic magnetic field along the line of sight ([Planck Collaboration et al., 2015](#); [Choi & Page, 2015](#)). The analysis is carried out over the full Northern sky footprint of QUIJOTE, using three Galactic latitude masks to assess the sensitivity of the results to Galactic plane contamination. The resulting foreground characterization is intended to serve as legacy input for component separation in upcoming CMB polarization experiments such as LiteBIRD ([Allys et al., 2022](#)).

2 OBJECTIVES

The main objective of this work is to characterize the spectral properties of the two dominant polarized Galactic foregrounds: synchrotron and thermal dust emission, using harmonic-space (power spectrum) tools. The analysis is performed on the combined dataset of QUIJOTE MFI DR1 maps at 11 and 17 GHz ([Rubiño-Martín et al., 2023](#)), all five WMAP 9-year frequency bands (K, Ka, Q, V, and W, [Bennett et al., 2013](#)), and the three LFI ([Planck Collaboration, 2016a](#)) and four HFI polarization bands ([Planck Collaboration et al., 2020a](#)) from Planck PR3, spanning a total frequency range from 11 to 353 GHz. This broad frequency coverage enables us to simultaneously fit a parametric spectral energy distribution (SED) model for both foreground components.

A second key objective is the characterization of the spatial cross-correlation between polarized synchrotron and thermal dust emission. Since both foregrounds are polarized by the same GMF, their EE and BB angular power spectra are expected to be partially correlated. Measuring this cross-correlation, alongside the auto-spectra of each component, provides complementary constraints on the geometry and coherence of the GMF and improves the accuracy of foreground separation in the CMB polarization context (Choi & Page, 2015).

Our results will be compared to prior power-spectrum studies that did not include QUIJOTE data. Martire et al. (2022) characterized the polarized synchrotron emission using WMAP and Planck data alone, obtaining a spectral index of $\beta_s = -3.00 \pm 0.10$ and a ratio of B- to E-mode amplitudes of $BB/EE \simeq 0.2$. Krachmalnicoff et al. (2018) obtained a steeper mean value of $\beta_s = -3.22 \pm 0.08$ when including the lower-frequency S-PASS data at 2.3 GHz.

For the thermal dust component, our results will be compared with those of Planck Collaboration et al. (2020b), who used Planck PR3 maps at 353 GHz to characterise dust polarization power spectra over sky regions covering 24 to 71% of the sky. They found power-law multipole indices $\alpha_{EE} = -2.42 \pm 0.02$ and $\alpha_{BB} = -2.54 \pm 0.02$, a significant E/B power asymmetry, and variations of the spectral exponents across sky regions. The dust polarization SED was well described by a modified black body with spectral index $\beta_d = 1.53 \pm 0.02$, with no evidence for frequency decorrelation between 100 and 353 GHz. For the synchrotron-dust cross-correlation, the benchmark is provided by Choi & Page (2015), who showed that a joint foreground model including the synchrotron-dust cross-correlation fits all WMAP and Planck frequency cross-spectra. These results constitute the main observational benchmarks against which our joint synchrotron-plus-dust analysis will be validated.

Beyond the immediate scientific goals, this work aims to provide robust foreground constraints and data products, including spectral index estimates and validated foreground models, that will serve as essential inputs for component separation in forthcoming CMB polarization experiments, including (Allys et al., 2022) and other ground-based and space facilities.

The analysis is carried out over the full sky footprint available to the experiments, using three Galactic masks with different latitude cuts to assess the sensitivity of our results to Galactic plane emission.

3 DATA

All data used in this work consist of three independent maps for each frequency band: one intensity map, corresponding to the Stokes parameter I , and two maps describing linear polarization, given by the Stokes parameters Q and U . In this work, we consider polarization data from three experiments: QUIJOTE-MFI, WMAP, and Planck, which are described below.

3.1. Q-U-I JOint TEnerife (QUIJOTE)

The QUIJOTE experiment (Rubiño-Martín et al., 2023) observes the visible sky from the Teide Observatory (latitude $+28.3^\circ$), covering all regions with elevations greater than 30° in both intensity and polarization. The experiment consists of two telescopes equipped with three polarimetric instruments: the Multifrequency Instrument (MFI), operating in the 10–20 GHz range; the Thirty-GHz Instrument (TGI); and the Forty-GHz Instrument (FGI).

In this work we use data from the 11 and 17 GHz bands of the MFI, which have angular resolutions of 55 and 38 arcmin, respectively. The data are obtained from the Legacy Archive for Microwave Background Data Analysis (LAMBDA)¹. The 11 GHz band is particularly useful for characterizing the polarization of low-frequency foregrounds, primarily synchrotron emission (Adak et al., 2025; Casas et al., 2025) and Anomalous Microwave Emission (AME, Fernández-Torreiro et al., 2023; González-González et al., 2025).

¹<https://lambda.gsfc.nasa.gov/product/quijote/index.html>

3.2. Wilkinson Microwave Anisotropy Probe (WMAP)

WMAP was a space-based experiment that observed the total intensity and polarization of the full sky across five frequency bands, spanning the frequency range from 23 to 94 GHz. In this work, we make use of all five frequency bands from the 9-year data release (Bennett et al., 2013): the K-band (23 GHz, $FWHM = 52.8$ arcmin), Ka-band (33 GHz, $FWHM = 39.6$ arcmin), Q-band (41 GHz, $FWHM = 31.2$ arcmin), V-band (61 GHz, $FWHM = 21.0$ arcmin), and W-band (94 GHz, $FWHM = 13.2$ arcmin). All maps are downloaded from the LAMBDA database².

3.3. Planck

Planck was a space-based experiment operating two complementary instruments: the Low Frequency Instrument (LFI) and the High Frequency Instrument (HFI), together observing the full sky in intensity and polarization across nine frequency channels, from 30 to 857 GHz. In this work, we use data from the third public release (PR3) of the Planck mission, available through the Planck Legacy Archive³ (PLA).

3.3.1 Low Frequency Instrument (LFI)

The LFI instrument covered three frequency bands at 28.4, 44.1, and 70.4 GHz. We use the PR3 frequency maps from all three LFI channels (Planck Collaboration, 2016a). The original angular resolutions correspond to effective beams of 32.65, 27.94, and 13.01 arcmin at 28.4, 44.1, and 70.4 GHz, respectively. All maps are downgraded to a HEALPIX⁴ resolution of $N_{\text{side}} = 512$ to maintain a common pixelization with the QUIJOTE and WMAP datasets.

3.3.2 High Frequency Instrument (HFI)

The HFI instrument provides six frequency bands extending up to 857 GHz. In this analysis we use the PR3 frequency maps from the HFI channels up to 353 GHz (100, 143, 217, and 353 GHz) (Planck Collaboration et al., 2020a). These four channels provide polarization information, including Stokes Q and U maps, and are therefore used in our polarization analysis. The higher-frequency HFI channels at 545 and 857 GHz are not included, as they do not provide polarization measurements. To ensure a consistent pixelization across the QUIJOTE, WMAP, and Planck datasets, all selected maps are downgraded to $N_{\text{side}} = 512$.

4 METHODOLOGY

In this section we describe the methodology followed to characterise Galactic foreground emission using multi-frequency angular power spectrum analysis. The approach relies entirely on harmonic-space statistics, avoiding the complications of real-space fitting, and exploits the broad frequency coverage provided by the combination of QUIJOTE, WMAP, and Planck datasets. We first introduce the mathematical framework of angular power spectrum analysis (Section 4.1.), then describe the construction of the analysis masks (Section 4.2.), the data processing steps (beam in Section 4.3.1, unit corrections in Section 4.3.2, noise in Section 4.3.3, and colour corrections in Section 4.3.4), the simulations used to estimate uncertainties (Section 4.4.), the foreground models (Section 4.5.), and finally, the Bayesian parameter inference (Section 4.6.).

4.1. Angular power spectrum (APS) analysis

The APS is a statistical tool that describes the distribution of signal power as a function of angular scale on the sky. It is a particularly powerful instrument for studying the statistical properties of diffuse emission, since it allows one to separate contributions from different physical processes through their characteristic frequency and multipole dependencies. Rather than working directly

²<https://lambda.gsfc.nasa.gov/product/wmap/current/index.html>

³<https://pla.esac.esa.int/>

⁴<https://healpy.readthedocs.io/en/latest/>

in pixel space, all analyses presented in this work are carried out in harmonic space, where the APS constitutes the fundamental observable.

The APS is expressed in terms of the spherical harmonic coefficients $a_{\ell m}$, which encode the sky signal at each multipole ℓ and azimuthal order m . The multipole ℓ is inversely proportional to the angular scale $\theta \sim 180^\circ/\ell$, so that large angular scales correspond to low- ℓ values, and small angular scales to high- ℓ values. For reference, $\ell = 200$ corresponds to approximately 1° , which is comparable to the angular resolution of the QUIJOTE-MFI instrument. The cross-power spectrum between two frequency maps X and Y is defined as:

$$C_\ell^{X \times Y} = \left\langle \frac{\sum_m a_{\ell m}^X (a_{\ell m}^Y)^*}{2\ell + 1} \right\rangle, \quad (1)$$

where X and Y denote the two frequency bands. When $X = Y$ this reduces to the auto-power spectrum. Cross-spectra probe the signal component common to two independent observations, suppressing the contribution of uncorrelated noise. By combining auto- and cross-spectra from 14 frequency bands (2 QUIJOTE, 5 WMAP and 7 Planck channels), we form a total of 105 spectra (14 auto and 91 cross), greatly improving the constraining power on the foreground model parameters.

All power spectra in this work are computed using the NAMASTER PYTHON library⁵ (Alonso et al., 2019), which implements the pseudo- C_ℓ method with an analytical treatment of the mode-coupling matrix induced by the sky mask. In particular, the E- and B-mode polarization fields are treated with B-mode purification (Smith, 2006), which suppresses E-to-B leakage arising from the application of a partial sky mask. The power spectra are computed in nine multipole bins of width $\Delta\ell = 20$ over the range $20 \leq \ell \leq 199$, with bin edges defined as

$$\ell_{\text{bin}}^{(k)} \in [\ell_1^{(k)}, \ell_2^{(k)}] = [20 + 20(k-1), 39 + 20(k-1)], \quad k = 1, \dots, 9. \quad (2)$$

The effective multipole ℓ_{eff} reported for each bin is computed by NAMASTER as the weighted average of the individual multipoles within the bin:

$$\ell_{\text{eff}}^{(k)} = \frac{\sum_{\ell \in \text{bin } k} w_\ell \ell}{\sum_{\ell \in \text{bin } k} w_\ell}, \quad (3)$$

where w_ℓ are the per-multipole weights assigned to the bandpower. In this work uniform weights are used, so $\ell_{\text{eff}}^{(k)}$ reduces to the arithmetic mean of all integer multipoles in the k -th bin, yielding $\ell_{\text{eff}} = 29, 49, 69, \dots, 189$ for the nine bins.

4.2. Masks

The analysis is restricted to the sky region accessible to the QUIJOTE-MFI telescope (Section 3.1), which operates from the Teide Observatory and therefore covers a declination band of approximately $-30^\circ < \delta < +90^\circ$. To mitigate contamination from the bright Galactic plane, three analysis masks are constructed by combining the QUIJOTE low-declination survey footprint (Rubiño-Martín et al., 2023) with Galactic latitude cuts at $|b| > 10^\circ$, $|b| > 15^\circ$, and $|b| > 20^\circ$. Each binary mask is apodized using the C2 scheme of NAMASTER with an apodization scale of 5° , which tapers the mask smoothly to zero at its edges and suppresses ringing effects in harmonic space. The three masks are shown in Figure 1. The effective sky fractions after apodization are approximately $f_{\text{sky}} \approx 0.30, 0.26$, and 0.23 for the 10, 15, and 20 degree cuts, respectively. The masks are defined on a HEALPIX grid at $N_{\text{side}} = 512$ and are identical for all frequency bands.

⁵<https://namaster.readthedocs.io/en/latest/>




Figure 1: Analysis masks combining the QUIJOTE low-declination survey footprint with Galactic latitude cuts of $|b| > 10^\circ$ (left), $|b| > 15^\circ$ (middle), and $|b| > 20^\circ$ (right), after C2 apodisation with a 5° border.

4.3. Data corrections

The raw power spectra computed from the observed maps contain systematic contributions from instrumental effects that must be accounted for before a physical interpretation can be drawn. In the ideal case, the measured C_ℓ coefficients would be proportional to the square of the theoretical spherical harmonic coefficients, $|a_{\ell m}^{\text{theo}}|^2$. In practice, several multiplicative and additive corrections are applied, as described below.

4.3.1 Beam and pixel window functions

The finite angular resolution of each instrument and the pixelization of the HEALPIX maps (Górski et al., 2005) introduce multiplicative window functions that suppress power at high multipoles. The pixel window function, w_ℓ , arises from the averaging of the sky signal over each pixel and depends solely on N_{side} ; for $N_{\text{side}} = 512$ this correction is small but non-negligible at $\ell > 500$. The instrumental beam is characterised by the window function B_ℓ , which captures the effective angular resolution of the experiment and causes an exponential suppression of power above the beam scale.

For each frequency band, the beam window function B_ℓ is loaded from the official data products of each experiment and interpolated to the effective multipoles of the binning scheme. For QUIJOTE-MFI, the beam is provided as a tabulated array per frequency channel in a FITS file. For WMAP, it is provided as a normalised ASCII table. For Planck LFI and HFI bands, the beam window functions are taken from the respective official FITS products, with separate T, E, and B transfer functions for HFI channels.

Both cases, auto- and cross-spectra, are handled by a single correction formula. Defining the effective transfer function for band i as $\mathcal{W}_\ell^i \equiv w_\ell B_\ell^i$, where w_ℓ is the HEALPIX pixel window function (common to all bands at fixed N_{side}) and B_ℓ^i is the instrumental beam window function of band i , the corrected spectrum between bands i and j is:

$$C_\ell^{i \times j, \text{corr}} = \frac{C_\ell^{i \times j, \text{sky}} - \delta_{ij} N_\ell^i}{\mathcal{W}_\ell^i \mathcal{W}_\ell^j}, \quad (4)$$

where δ_{ij} is the Kronecker delta, N_ℓ^i is the noise power spectrum of band i (see Section 4.3.3), and the denominator deconvolves the combined effect of the pixel window and the beam of each band. The Kronecker delta ensures that the noise bias is subtracted only for auto-spectra ($i = j$); for cross-spectra between different frequency bands the noise contributions of the two detectors are statistically independent and thus average to zero in the cross-correlation, so no noise subtraction is required.

4.3.2 Unit conversion

All sky maps are converted to a common unit of K_{RJ} (Rayleigh–Jeans brightness temperature) before computing the power spectra. The unit conversion factor from thermodynamic CMB temperature K_{CMB} to K_{RJ} is:

$$u(\nu) = \frac{c^2}{2k_B\nu^2} \left. \frac{\partial B_\nu}{\partial T} \right|_{T_{\text{CMB}}}, \quad (5)$$

where B_ν is the Planck function, k_B is the Boltzmann constant, c the speed of light, and $T_{\text{CMB}} = 2.725$ K. For Planck HFI bands (100–353 GHz), the conversion uses the official bandpass-averaged unit conversion coefficients from the Planck 2018 data release (Planck Collaboration et al., 2020a), accounting for the instrumental bandpass shape. For all remaining bands the conversion is evaluated analytically at the effective band-centre frequency. The resulting unit conversion factor applied to a power spectrum of the pair (i, j) is $u(\nu_i) \times u(\nu_j)$, where $i = j$ for auto-spectra and $i \neq j$ for cross-spectra.

4.3.3 Noise power spectrum

The instrumental noise contributes an additive bias to the auto-power spectrum that must be estimated and removed (Equation 4). The noise estimation strategy is specific to each experiment.

For QUIJOTE-MFI, the HMDM is constructed from the two half-survey maps provided in the DR1 data release (Rubiño-Martín et al., 2023), together with their associated per-pixel weight maps w_1 and w_2 (where $w_i = 1/\sigma_i^2$):

$$n = \frac{h_1 - h_2}{\sqrt{(w_1 + w_2)(w_1^{-1} + w_2^{-1})}}. \quad (6)$$

This construction ensures that the noise power spectrum of the difference map matches that of the combined weighted-sum map, while containing no sky signal (Planck Collaboration, 2016a).

For Planck, the same methodology is followed but the nature of the data split differs between the two focal plane arrays. For the LFI (30–70 GHz), ring-half maps are used as h_1 and h_2 , with pixel covariance maps providing the weights (Planck Collaboration, 2016a). For the HFI (100–353 GHz), the official half-mission maps from the Planck 2018 data release (Planck Collaboration et al., 2020a) are used. In both cases, the covariance-weighted HMDM is constructed using Equation 6. Since Planck maps are provided at a higher resolution (N_{side}), the resulting HMDM is downgraded to $N_{\text{side}} = 512$ after construction to match the resolution of the other datasets.

For WMAP, the official band-average maps are noise-weighted co-additions of all Differential Assemblies (DAs) at each frequency. No half-mission split is provided directly for the band-average maps; instead, we construct the HMDM from the temporal split available in the public data release. We use the first four years of observations (h_1 , years 1 to 4) and the remaining five years (h_2 , years 5 to 9). Because the two halves average over different numbers of years, the naive difference $h_1 - h_2$ does not reproduce the noise level of the full 9-year combination. We apply the analytical normalization factor that accounts for the unequal time splits:

$$n_{\text{WMAP}} = (h_1 - h_2) \sqrt{\frac{t_1 t_2}{(t_1 + t_2)^2}}, \quad (7)$$

where $t_1 = 4$ and $t_2 = 5$ are the number of years in each half, giving a normalization factor of $\sqrt{20/81}$. This ensures that the noise power spectrum of the HMDM matches that of the full 9-year map, and it is used directly as N_ℓ^i in Equation 4.

4.3.4 Colour corrections

Colour corrections account for the mismatch between the SED of the foreground emission and the spectral response of each instrumental bandpass. When a source with spectral index β is observed through a band with transmission $\tau(\nu)$, the measured flux density differs from the value at the nominal band-centre frequency by a factor that depends on β . This factor is the colour correction $\text{CC}(\beta)$.

The colour correction coefficients for each band and each foreground component (synchrotron and thermal dust) are pre-computed using the `fastcc` code⁶ (Peel et al., 2022), which evaluates the bandpass-weighted ratio between the assumed foreground SED and a flat (Rayleigh-Jeans) reference spectrum.

The treatment differs between foreground components because they follow different SEDs. For synchrotron emission (power law, $I_\nu \propto \nu^\alpha$), the colour correction depends only on a single spectral index α , so it can be parameterized as a 1D polynomial. The same applies to dust emission in non-HFI bands, where the modified blackbody curvature is negligible and a power-law approximation is sufficient. For these components, the coefficients are obtained using `fastcc`. For thermal dust in the Planck HFI bands (100-353 GHz), however, the SED follows a modified blackbody (MBB),

$$I_\nu \propto \nu^{\beta_d} B_\nu(\nu, T_d), \quad (8)$$

where the Planck function $B_\nu(\nu, T_d)$ introduces a non-trivial dependence on the dust temperature T_d that cannot be absorbed into a power-law index. As a result, the HFI dust colour correction depends jointly on both β_d and T_d , and a simple 1D polynomial in α is insufficient. It is therefore computed using the `interpcc` interpolator (provided with `fastcc`), which interpolates a two-dimensional grid of pre-computed colour corrections as a function of both β_d and T_d (Peel et al., 2022). The grid is evaluated at the fixed dust temperature $T_d = 19.6$ K adopted throughout this work, reducing the 2D surface to a 1D function of β_d alone, which is then fitted with a second-order polynomial.

In all cases the resulting colour correction for component $c \in \{s, d\}$ at frequency band ν is expressed as a second-order polynomial in the relevant spectral index:

$$\text{CC}_c(\nu; \alpha) = a_0^c(\nu) + a_1^c(\nu) \alpha + a_2^c(\nu) \alpha^2, \quad (9)$$

where $\alpha = \beta_s + 2$ for synchrotron ($c = s$) and $\alpha = \beta_d + 2$ for dust ($c = d$) in Rayleigh-Jeans units. The polynomial coefficients $\{a_0^c(\nu), a_1^c(\nu), a_2^c(\nu)\}$ are specific to each component c and frequency band ν , and are derived using `fastcc` following the methodology described above.

Rather than being applied as a fixed correction to the data, the colour corrections are folded directly into the foreground model (see Section 4.5.) at each MCMC iteration, using the spectral index proposed by the sampler at that step. Each foreground component carries its own correction, applied at each frequency of the pair. The colour-corrected model terms are:

$$C_\ell^{\text{sync, cc}}(\nu_i, \nu_j) = \frac{A_s (\ell/\ell_{\text{ref}})^{\alpha_s} S_s(\nu_i) S_s(\nu_j)}{\text{CC}_s(\nu_i; \beta_s) \text{CC}_s(\nu_j; \beta_s)}, \quad (10)$$

$$C_\ell^{\text{dust, cc}}(\nu_i, \nu_j) = \frac{A_d (\ell/\ell_{\text{ref}})^{\alpha_d} S_d(\nu_i) S_d(\nu_j)}{\text{CC}_d(\nu_i; \beta_d) \text{CC}_d(\nu_j; \beta_d)}, \quad (11)$$

and for the synchrotron-dust cross-correlation, each frequency leg receives the correction appropriate for its own component:

$$C_\ell^{\text{corr, cc}}(\nu_i, \nu_j) = \rho \sqrt{A_s A_d} \left(\frac{\ell}{\ell_{\text{ref}}} \right)^{(\alpha_s + \alpha_d)/2} \left[\frac{S_s(\nu_i)}{\text{CC}_s(\nu_i; \beta_s)} \frac{S_d(\nu_j)}{\text{CC}_d(\nu_j; \beta_d)} + \frac{S_s(\nu_j)}{\text{CC}_s(\nu_j; \beta_s)} \frac{S_d(\nu_i)}{\text{CC}_d(\nu_i; \beta_d)} \right]. \quad (12)$$

Since the corrections are re-evaluated at every likelihood call, they track the spectral index throughout the chain rather than being fixed at a single approximate value. This propagates the uncertainty in the spectral index self-consistently into the corrected model, improving upon approaches where the colour correction is applied as a fixed factor computed at a mean spectral index (Planck Collaboration et al., 2020b).

⁶<https://github.com/mpel/fastcc>

4.4. Simulations

The statistical uncertainties on the measured power spectra are estimated from Monte Carlo simulations. In the corrected spectrum (Equation 4), two independent terms contribute to the total error: the uncertainty on the sky power spectrum $C_\ell^{i \times j, \text{sky}}$, characterized by σ_{C_ℓ} , and the uncertainty on the noise power spectrum N_ℓ^i , characterized by σ_{N_ℓ} . These are estimated from two independent sets of $N_{\text{sim}} = 100$ simulations, and the total uncertainty is obtained as their quadrature sum:

$$\sigma_\ell = \sqrt{\sigma_{C_\ell}^2 + \sigma_{N_\ell}^2}, \quad (13)$$

where σ_{C_ℓ} is the standard deviation of the corrected power spectra across sky-plus-noise realizations, and σ_{N_ℓ} is the standard deviation of the noise power spectra across noise-only realizations. Note that in the case of cross-spectra, the noise term N_ℓ^i does not appear in Equation 4 (since $\delta_{ij} = 0$), so $\sigma_{N_\ell} = 0$ and the total uncertainty reduces to $\sigma_\ell = \sigma_{C_\ell}$ alone.

4.4.1 Sky signal simulations

Realistic simulated polarization sky maps for each frequency band and experiment are generated using the Python Sky Model (PySM) package⁷ (Thorne et al., 2017). PySM is a widely used PYTHON library for sky emission modelling in CMB analyses (e.g. Planck Collaboration et al. 2020c; Martire et al. 2022).

The simulation includes both polarized and intensity foreground components. Since this work focuses exclusively on polarization, only the polarized components enter the power spectrum analysis; the intensity components are included solely to produce realistic total-intensity maps but do not contribute to the polarized signal and are not used in the fitting. The polarized foreground components are: synchrotron emission modelled as a power law with a spatially constant spectral index (**s1**); thermal dust emission described as a single-component modified blackbody (**d1**); and a lensed CMB polarization realization (**c1**), computed using the Taylens code (Næss & Louis, 2013) with unlensed C_ℓ produced by CAMB⁸. The intensity foreground components included are spinning dust (AME) modelled as a sum of two spinning dust populations (**a1**; Planck Collaboration 2016b), and free-free emission following the analytic Commander model (**f1**); these are included for completeness but play no role in the polarization analysis.

Simulations are produced independently for all 14 frequency bands. The sky emission at each band-centre frequency is obtained from PySM, converted to thermodynamic CMB temperature units, and subsequently convolved in harmonic space with the instrumental beam window function of the corresponding experiment at a resolution of $N_{\text{side}} = 512$.

4.4.2 Noise simulations

Noise simulations are produced independently for each frequency band and experiment, reflecting the distinct noise properties of each instrument.

QUIJOTE-MFI

The noise in the QUIJOTE-MFI maps is spatially correlated and anisotropic, and the two channels within each MFI horn exhibit significant noise correlations both in intensity ($\sim 60\text{--}80\%$) and in polarization ($\sim 20\%$; Rubiño-Martín et al. 2023). The QUIJOTE noise simulations used in this work are those described in Rubiño-Martín et al. (2023), which account for the measured anisotropy, spatial correlations, and inter-channel covariance structure.

WMAP

WMAP data are not affected by $1/f$ noise, so the noise is well described by a per-pixel Gaussian model. Following Bennett et al. (2013), $N_{\text{sim}} = 100$ noise realization maps are generated as independent Gaussian random fields with zero mean and pixel-dependent standard deviation. The per-pixel noise level for polarization is:

⁷<https://pysm3.readthedocs.io/en/latest/>

⁸<https://www.camb.info/>

$$\sigma_{\text{pix}} = \frac{\sigma_0}{\sqrt{N_{\text{obs, eff}}}}, \quad (14)$$

where σ_0 is the band-level noise sensitivity (tabulated in the WMAP data release) and $N_{\text{obs, eff}}$ is the effective number of observations per pixel, which accounts for the covariance between the Q and U Stokes parameters stored in the WMAP maps. Specifically, for the Q component:

$$N_{\text{obs, eff}}^Q = N_{\text{obs}}^Q - \frac{(N_{\text{obs}}^{QU})^2}{N_{\text{obs}}^U}, \quad (15)$$

and analogously for U, where N_{obs}^Q , N_{obs}^U , and N_{obs}^{QU} are the diagonal and off-diagonal elements of the per-pixel observation count matrix given by WMAP for each map and frequency band. This correction ensures that the effective noise per pixel correctly reflects the independent information content of Q and U after accounting for their covariance. The Q and U noise maps are drawn independently using their respective σ_{pix} .

Planck

For Planck LFI and HFI bands, $N_{\text{sim}} = 100$ per-band noise simulations are downloaded from the Planck Legacy Archive (Planck Collaboration et al., 2020c). These simulations are generated using the MCMC-based noise estimation procedures described in the respective data processing papers for LFI (Planck Collaboration, 2016a) and HFI (Planck Collaboration et al., 2020a), which characterise the noise power spectrum including contributions from the $1/f$ noise component of the detectors. Each simulation map is processed identically to the data in the analysis pipeline.

4.5. Foreground emission model

The measured power spectra over the frequency range 11–353 GHz are dominated by two Galactic emission components in polarization: synchrotron radiation and thermal dust emission (Planck Collaboration, 2016b). A cross-correlation term between the two components is included to account for any spatial correlation along the line of sight. This modelling approach is similar to that followed by Choi & Page (2015) and Hensley et al. (2022). The total model power spectrum between frequencies ν_i and ν_j we will use in this work is given by Equation 16.

$$C_\ell(\nu_i, \nu_j) = C_\ell^{\text{sync}}(\nu_i, \nu_j) + C_\ell^{\text{dust}}(\nu_i, \nu_j) + C_\ell^{\text{corr}}(\nu_i, \nu_j). \quad (16)$$

Here C_ℓ^{sync} is the synchrotron power spectrum, modelled as a power law in both frequency and multipole (Section 4.5.1); C_ℓ^{dust} is the thermal dust power spectrum, modelled via a modified blackbody SED (Section 4.5.2); and C_ℓ^{corr} is the synchrotron-dust cross-correlation term, which accounts for any spatial correlation between the two components (Section 4.5.3).

4.5.1 Synchrotron emission

Synchrotron radiation is the dominant polarized foreground at frequencies below ~ 70 GHz. Its angular power spectrum is well described by a power law in ℓ (Choi & Page, 2015; Martire et al., 2022). Defining the synchrotron frequency scaling as

$$S_s(\nu) = \left(\frac{\nu}{\nu_{\text{ref}}} \right)^{\beta_s}, \quad (17)$$

the synchrotron power spectrum is:

$$C_\ell^{\text{sync}}(\nu_i, \nu_j) = A_s \left(\frac{\ell}{\ell_{\text{ref}}} \right)^{\alpha_s} S_s(\nu_i) S_s(\nu_j), \quad (18)$$

where A_s is the synchrotron amplitude at the reference multipole $\ell_{\text{ref}} = 80$ and reference frequency $\nu_{\text{ref}} = 23.0$ GHz, α_s is the angular power spectral index, and β_s is the frequency spectral index in Rayleigh-Jeans units.

4.5.2 Thermal dust emission

Thermal dust emission dominates polarized foreground emission above ~ 70 GHz, with a SED following a modified blackbody (MBB) law (Planck Collaboration et al., 2020b). In Rayleigh-Jeans brightness temperature units, the MBB scaling relative to the dust reference frequency $\nu_0 = 353$ GHz is:

$$S_d(\nu) = \left(\frac{\nu}{\nu_0} \right)^{\beta_d} \frac{B_\nu(\nu, T_d)}{B_\nu(\nu_0, T_d)}, \quad (19)$$

where B_ν is the Planck function and $T_d = 19.6$ K is the dust temperature, fixed in accordance with Planck measurements (Planck Collaboration, 2016b). The dust model power spectrum is:

$$C_\ell^{\text{dust}}(\nu_i, \nu_j) = A_d \left(\frac{\ell}{\ell_{\text{ref}}} \right)^{\alpha_d} S_d(\nu_i) S_d(\nu_j), \quad (20)$$

where A_d is the dust amplitude at $\ell_{\text{ref}} = 80$ and $\nu_0 = 353$ GHz, α_d is the angular power spectral index, and β_d is the MBB frequency spectral index.

4.5.3 Synchrotron-dust cross-correlation

A non-zero spatial correlation between synchrotron and thermal dust emission can arise when the Galactic magnetic field aligns the polarization angle of both components along the line of sight (Choi & Page, 2015). Following Hensley et al. (2022), the cross-correlation term is modelled as:

$$C_\ell^{\text{corr}}(\nu_i, \nu_j) = \rho \sqrt{A_s A_d} \left[S_s(\nu_i) S_d(\nu_j) + S_s(\nu_j) S_d(\nu_i) \right] \left(\frac{\ell}{\ell_{\text{ref}}} \right)^{(\alpha_s + \alpha_d)/2}, \quad (21)$$

where $\rho \in [-1, 1]$ is the synchrotron-dust correlation coefficient. When colour corrections are applied, each per-band scaling factor is divided by its corresponding colour correction polynomial, as described in Section 4.3.4.

4.5.4 CMB subtraction

Before fitting the foreground model, the CMB contribution is subtracted from the observed power spectra. We use the best-fit theoretical Λ CDM power spectrum from Planck 2018 (Planck Collaboration, 2020), obtained from the combined Plik TTTEEE likelihood with low-multipole (lowE) and gravitational lensing constraints. Both C_ℓ^{EE} and C_ℓ^{BB} are subtracted; for BB only the lensing contribution is included, since primordial tensor B-modes have not yet been detected. The CMB spectrum is interpolated to the effective multipoles of the binning scheme before subtraction.

4.6. Bayesian inference and MCMC fitting

We adopt a Bayesian framework to constrain the foreground model parameters from the measured power spectra. The full parameter vector for the power-law fitting is $\boldsymbol{\theta} = \{A_s, \alpha_s, \beta_s, A_d, \alpha_d, \beta_d, \rho\}$, where the EE and BB modes are fitted separately in this work. The posterior distribution is:

$$p(\boldsymbol{\theta} \mid \mathbf{D}) \propto \mathcal{L}(\mathbf{D} \mid \boldsymbol{\theta}) p(\boldsymbol{\theta}), \quad (22)$$

where $p(\boldsymbol{\theta} \mid \mathbf{D})$ is the posterior probability of the parameters given the data $\mathbf{D}$, $\mathcal{L}(\mathbf{D} \mid \boldsymbol{\theta})$ is the likelihood of observing the data under the model with parameters $\boldsymbol{\theta}$, and $p(\boldsymbol{\theta})$ is the prior encoding our knowledge of the parameters before seeing the data. The posterior is thus proportional to the product of the likelihood and the prior, with the normalization constant (the Bayesian evidence) being irrelevant for parameter estimation.

4.6.1 Likelihood

The likelihood is constructed assuming that residuals between the measured and model power spectra are Gaussian distributed. This approximation is well justified in the regime of partial sky coverage and moderate multipoles ($\ell < 200$), where the central limit theorem ensures that the distribution of bandpower estimates is close to Gaussian, and has been widely adopted in foreground power spectrum analyses at similar angular scales (Choi & Page, 2015; Martire et al., 2022; Rubiño-Martín et al., 2023). The data vector $\mathbf{D}$ consists of the measured spectra $\hat{C}_\ell(\nu_i, \nu_j)$ concatenated over all frequency pairs and multipole bins, with corresponding uncertainties $\sigma_\ell(\nu_i, \nu_j)$ estimated from Monte Carlo realizations (Section 4.4.). The log-likelihood is then:

$$\ln \mathcal{L} = -\frac{1}{2} \sum_{i,j,\ell} \frac{[\hat{C}_\ell(\nu_i, \nu_j) - C_\ell(\nu_i, \nu_j; \boldsymbol{\theta})]^2}{\sigma_\ell^2(\nu_i, \nu_j)}. \quad (23)$$

4.6.2 Priors

Two fitting approaches are considered in this work: a power-law fit, in which all nine multipole bins are fitted simultaneously by a global power-law model, and a bin-to-bin fit, in which each multipole bin is fitted independently. Both approaches are described in detail in Section 4.7.. In both cases, uniform (flat) priors are adopted within physically motivated bounds:

$$A_s \geq 0, \quad A_d \geq 0, \quad \beta_s \leq 0, \quad \beta_d \geq 0, \quad \rho \in [-1, 1],$$

For the bin-to-bin analysis, an additional Gaussian prior on the synchrotron frequency spectral index is included to regularize the fit in multipole bins where the signal-to-noise ratio in the low-frequency channels is too low to adequately constrain the synchrotron SED, which can otherwise lead to spurious results for β_s and, consequently, for β_d (Planck Collaboration et al., 2020b):

$$\beta_s \sim \mathcal{N}(-3.1, 0.30^2), \quad (24)$$

consistent with the values found in published WMAP and Planck foreground analyses (Gold et al., 2011; Planck Collaboration et al., 2016). This approach has also been used in various works as in Planck Collaboration et al. (2020b) or Choi & Page (2015).

4.6.3 Sampler

Posterior sampling is performed with the EMCEE affine-invariant ensemble sampler (Foreman-Mackey et al., 2013). Convergence is assessed for all fits using a combination of standard diagnostics: the integrated autocorrelation time $\hat{\tau}$ is monitored to ensure the post-burn-in chain is sufficiently long; the effective sample size ($\text{ESS} = N_{\text{post}}/\hat{\tau}$) is required to exceed 1000 per parameter; the mean acceptance fraction across walkers is verified to lie within the range recommended for the affine-invariant sampler; and the reduced chi-squared χ_{red}^2 at the posterior maximum-likelihood point is inspected to confirm that the model provides an adequate fit and that the uncertainties are well calibrated.

4.7. Fitting approaches

Two complementary fitting approaches are applied to the corrected power spectra over the multipole range $20 \leq \ell \leq 199$, using data from all 14 frequency bands described in Section 3. The baseline analysis includes both auto- and cross-spectra, yielding 105 independent spectra.

Moreover, a complementary study restricted to cross-spectra only, which avoids any noise bias in the auto-spectra, is presented in Appendix A; and a study fitting in the multipole range $30 \leq \ell \leq 199$ suggested in Rubiño-Martín et al. (2023) is presented in Appendix B.

4.7.1 Power-law fitting

In this approach all nine multipole bins are fitted simultaneously by the global power-law foreground model of Section 4.5. The MCMC samples the full parameter vector $\theta = \{A_s, \alpha_s, \beta_s, A_d, \alpha_d, \beta_d, \rho\}$ over all frequencies and multipoles. EE and BB modes are fitted separately.

4.7.2 Bin-to-bin analysis

In this approach each multipole bin is fitted independently, without imposing any global power-law dependence on ℓ . The per-bin parameter vector is $\theta_{\text{bin}} = \{A_s, \beta_s, A_d, \beta_d, \rho\}$; the multipole spectral indices (α_s, α_d) are unconstrained in a single-bin analysis. This approach enables us to examine how the fitted parameters vary with the effective multipole, ℓ_{eff} , revealing any scale-dependent behavior or local deviations from the global power-law assumption.

5 RESULTS

In this section we present the results of the foreground power spectrum analysis described in Section 4. The analysis is performed over the multipole range $20 \leq \ell \leq 199$ using the combined 14-band of QUIJOTE, WMAP and Planck dataset. Two complementary fitting approaches are applied: a global power-law fit (Section 4.7.1) and an independent bin-to-bin fit (Section 4.7.2). For each approach, results are obtained for three Galactic masks defined by latitude cuts at $|b| > 10^\circ$, 15° , and 20° (Section 4.2.), allowing us to assess the robustness of the derived foreground parameters with respect to Galactic plane contamination. In all cases, the EE and BB polarization modes are fitted separately.

The foreground model and the Bayesian inference framework are described in Sections 4.5. and 4.6., respectively. This analysis follows approaches similar to those of Planck Collaboration et al. (2020b), Choi & Page (2015), Hensley et al. (2022), and Martire et al. (2022). The baseline results presented here include both auto- and cross-spectra (105 spectra in total). A complementary analysis restricted to cross-spectra only, which avoids any noise bias contribution from the auto-spectra, is presented in Appendix A.

The multipole range adopted here starts at $\ell = 20$, rather than the $\ell \geq 30$ range used in several comparable studies (Rubiño-Martín et al., 2023; Martire et al., 2022). The choice is motivated by the behaviour of the BB-mode power-law fit: the BB signal is intrinsically fainter than EE and falls steeply with multipole, so the first bin ($\ell_{\text{eff}} = 29$) carries a large fraction of the constraining power on the angular spectral index α_s . When this bin is excluded, the remaining bins at smaller angular scales provide insufficient leverage to anchor the power-law slope, leading to steeper and poorly constrained values of α_s . Including the first bin resolves this degeneracy and yields well-constrained, physically consistent results for the BB mode. This effect is demonstrated in Appendix B, where results obtained over the range $30 \leq \ell \leq 199$ are presented for comparison.

5.1. Power-law fitting

The global power-law model (Section 4.7.1) is fitted simultaneously over all nine multipole bins and all 105 frequency spectra. The posterior constraints on the synchrotron amplitude A_s , multipole spectral index α_s , frequency spectral index β_s , dust amplitude A_d , multipole spectral index α_d , frequency spectral index β_d , and synchrotron-dust correlation coefficient ρ are reported as the median and 68% credible intervals.

Results are obtained separately for EE and BB polarization modes, and for two dataset combinations: WMAP+Planck (WP+Pl) and QUIJOTE+WMAP+Planck (QJ+WP+Pl). These combinations are used to quantify the impact of including QUIJOTE-MFI data in the analysis.

The results for the three Galactic masks are presented below. For each mask, we provide a table of the best-fit parameters, followed by figures illustrating the best-fit synchrotron and dust spectral models alongside the measured frequency spectra.

The posterior constraints for the $|b| > 10^\circ$ mask are given in Table ??, and the corresponding spectral model fits are shown in Figure 2.

The posterior constraints for the $|b| > 15^\circ$ mask are given in Table ??, and the corresponding spectral model fits are shown in Figure 3.

The posterior constraints for the $|b| > 20^\circ$ mask are given in Table ??, and the corresponding spectral model fits are shown in Figure 4.

Table 1: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes, spectral indices, and synchrotron-dust correlation coefficient ρ , from a global power-law fit to the EE and BB polarization auto- and cross-spectra of the 14-band QUIJOTE+WMAP+Planck dataset over the multipole range $20 \leq \ell \leq 199$, for the three Galactic masks.

Mask	Dataset	Mode	$A_s [10^{-3} \mu\text{K}^2]$	α_s	β_s	$A_d [10^{-3} \mu\text{K}^2]$	α_d	β_d	ρ
$ b > 10^\circ$	WP+P1	EE	$6.687^{+0.421}_{-0.411}$	$-3.710^{+0.063}_{-0.065}$	$-3.017^{+0.041}_{-0.042}$	$3.217^{+0.011}_{-0.011}$	$-2.546^{+0.004}_{-0.004}$	$1.530^{+0.005}_{-0.005}$	$0.094^{+0.007}_{-0.007}$
		BB	$1.895^{+0.342}_{-0.317}$	$-3.258^{+0.172}_{-0.187}$	$-3.195^{+0.140}_{-0.145}$	$2.359^{+0.007}_{-0.007}$	$-2.203^{+0.004}_{-0.004}$	$1.530^{+0.005}_{-0.005}$	$0.132^{+0.013}_{-0.013}$
	QJ+WP+P1	EE	$7.398^{+0.260}_{-0.256}$	$-3.619^{+0.037}_{-0.038}$	$-3.085^{+0.016}_{-0.016}$	$3.216^{+0.011}_{-0.011}$	$-2.547^{+0.004}_{-0.004}$	$1.530^{+0.005}_{-0.005}$	$0.096^{+0.005}_{-0.005}$
		BB	$1.657^{+0.186}_{-0.180}$	$-3.334^{+0.112}_{-0.121}$	$-2.999^{+0.052}_{-0.052}$	$2.359^{+0.007}_{-0.007}$	$-2.203^{+0.004}_{-0.004}$	$1.530^{+0.005}_{-0.005}$	$0.111^{+0.010}_{-0.010}$
$ b > 15^\circ$	WP+P1	EE	$5.395^{+0.425}_{-0.415}$	$-3.874^{+0.078}_{-0.082}$	$-3.042^{+0.047}_{-0.048}$	$2.093^{+0.010}_{-0.010}$	$-2.534^{+0.006}_{-0.006}$	$1.552^{+0.007}_{-0.007}$	$0.080^{+0.008}_{-0.008}$
		BB	$1.723^{+0.376}_{-0.342}$	$-3.284^{+0.207}_{-0.229}$	$-3.314^{+0.175}_{-0.185}$	$1.378^{+0.006}_{-0.006}$	$-2.317^{+0.006}_{-0.006}$	$1.530^{+0.007}_{-0.007}$	$0.128^{+0.016}_{-0.015}$
	QJ+WP+P1	EE	$5.990^{+0.260}_{-0.255}$	$-3.767^{+0.045}_{-0.046}$	$-3.079^{+0.017}_{-0.017}$	$2.092^{+0.010}_{-0.009}$	$-2.535^{+0.006}_{-0.006}$	$1.551^{+0.007}_{-0.007}$	$0.068^{+0.006}_{-0.006}$
		BB	$1.367^{+0.198}_{-0.183}$	$-3.410^{+0.141}_{-0.150}$	$-3.017^{+0.061}_{-0.061}$	$1.379^{+0.006}_{-0.006}$	$-2.317^{+0.006}_{-0.006}$	$1.531^{+0.007}_{-0.007}$	$0.116^{+0.013}_{-0.012}$
$ b > 20^\circ$	WP+P1	EE	$5.948^{+0.451}_{-0.439}$	$-3.741^{+0.076}_{-0.080}$	$-3.140^{+0.054}_{-0.055}$	$1.439^{+0.009}_{-0.009}$	$-2.479^{+0.008}_{-0.008}$	$1.487^{+0.008}_{-0.008}$	$0.030^{+0.009}_{-0.009}$
		BB	$1.436^{+0.374}_{-0.336}$	$-3.476^{+0.243}_{-0.272}$	$-3.456^{+0.186}_{-0.198}$	$0.927^{+0.005}_{-0.005}$	$-2.237^{+0.007}_{-0.007}$	$1.532^{+0.009}_{-0.009}$	$0.162^{+0.019}_{-0.018}$
	QJ+WP+P1	EE	$5.880^{+0.258}_{-0.253}$	$-3.732^{+0.046}_{-0.047}$	$-3.113^{+0.019}_{-0.019}$	$1.439^{+0.009}_{-0.009}$	$-2.478^{+0.008}_{-0.008}$	$1.486^{+0.008}_{-0.008}$	$0.023^{+0.007}_{-0.007}$
		BB	$1.347^{+0.208}_{-0.198}$	$-3.371^{+0.151}_{-0.164}$	$-2.984^{+0.065}_{-0.065}$	$0.927^{+0.005}_{-0.005}$	$-2.238^{+0.007}_{-0.007}$	$1.533^{+0.009}_{-0.009}$	$0.121^{+0.015}_{-0.015}$






Figure 2: Best-fit global power-law models for synchrotron and thermal dust emission overlaid on the measured EE and BB frequency power spectra, for the $|b| > 10^\circ$ Galactic mask.

5.2. Bin-to-bin fitting

In the bin-to-bin approach (Section 4.7.2), each of the nine multipole bins is fitted independently, without imposing any global power-law dependence on ℓ . The per-bin parameter vector is $\theta_{\text{bin}} = \{A_s, \beta_s, A_d, \beta_d, \rho\}$, and results are obtained separately for EE and BB modes. This approach allows the identification of scale-dependent trends and potential deviations from the global power-law behaviour found in Section 5. In multipole bins where the signal-to-noise ratio in the low-frequency channels is insufficient to independently constrain the synchrotron SED, the Gaussian prior on β_s described in Section 4.6. is applied to regularize the fit.

Posterior constraints (median and 68% credible intervals) on A_s , β_s , A_d , β_d , and ρ are reported as a function of the effective multipole ℓ_{eff} for each bin. Results are presented for the three Galactic masks below. The corresponding cross-spectra-only results are provided in Appendix A.

Figure 3: Same as Figure 2 but for the $|b| > 15^\circ$ Galactic mask.Figure 4: Same as Figure 2 but for the $|b| > 20^\circ$ Galactic mask.

$|b| > 10^\circ$ mask

BIN-TO-BIN RESULTS – MASK $|b| > 10^\circ$

The bin-to-bin posterior constraints for the $|b| > 10^\circ$ mask are given in Table 2.

$|b| > 15^\circ$ mask

BIN-TO-BIN RESULTS – MASK $|b| > 15^\circ$

The bin-to-bin posterior constraints for the $|b| > 15^\circ$ mask are given in Table 3.

$|b| > 20^\circ$ mask

BIN-TO-BIN RESULTS – MASK $|b| > 20^\circ$

The bin-to-bin posterior constraints for the $|b| > 20^\circ$ mask are given in Table 4.

Table 2: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes, frequency spectral indices, and synchrotron-dust correlation coefficient ρ , fitted independently per multipole bin from the EE (top) and BB (bottom) polarization auto- and cross-spectra of the 14-band QUIJOTE+WMAP+Planck data set, using the $|b| > 10^\circ$ Galactic mask.

ℓ range	ℓ_{eff}	$A_s^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_s^{EE}	$A_d^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_d^{EE}	ρ^{EE}	χ_{red}^2
20–39	29.0	$294.505^{+3.295}_{-3.397}$	$-3.075^{+0.018}_{-0.018}$	$49.086^{+0.270}_{-0.261}$	$1.481^{+0.007}_{-0.007}$	$0.092^{+0.007}_{-0.007}$	1.564
40–59	49.0	$37.590^{+1.238}_{-1.322}$	$-3.135^{+0.050}_{-0.047}$	$7.887^{+0.077}_{-0.084}$	$1.519^{+0.014}_{-0.013}$	$0.117^{+0.013}_{-0.014}$	1.251
60–79	69.0	$13.055^{+0.753}_{-0.759}$	$-3.234^{+0.068}_{-0.069}$	$4.798^{+0.040}_{-0.041}$	$1.531^{+0.012}_{-0.012}$	$0.044^{+0.014}_{-0.014}$	1.169
80–99	89.0	$6.238^{+0.606}_{-0.639}$	$-3.133^{+0.116}_{-0.110}$	$2.673^{+0.022}_{-0.023}$	$1.504^{+0.013}_{-0.013}$	$0.055^{+0.021}_{-0.019}$	1.433
100–119	109.0	$3.519^{+0.507}_{-0.523}$	$-2.966^{+0.165}_{-0.164}$	$1.555^{+0.016}_{-0.017}$	$1.567^{+0.019}_{-0.018}$	$0.037^{+0.028}_{-0.027}$	1.084
120–139	129.0	$2.535^{+0.477}_{-0.500}$	$-2.594^{+0.230}_{-0.219}$	$1.017^{+0.013}_{-0.013}$	$1.420^{+0.021}_{-0.020}$	$0.174^{+0.044}_{-0.041}$	1.495
140–159	149.0	$1.980^{+0.583}_{-0.593}$	$-2.909^{+0.238}_{-0.238}$	$0.679^{+0.010}_{-0.010}$	$1.477^{+0.026}_{-0.025}$	$0.225^{+0.057}_{-0.048}$	0.955
160–179	169.0	$1.299^{+0.524}_{-0.502}$	$-3.094^{+0.278}_{-0.271}$	$0.479^{+0.008}_{-0.008}$	$1.543^{+0.029}_{-0.028}$	$0.073^{+0.062}_{-0.059}$	1.002
180–199	189.0	$1.070^{+0.658}_{-0.617}$	$-2.903^{+0.296}_{-0.299}$	$0.404^{+0.007}_{-0.007}$	$1.635^{+0.032}_{-0.032}$	$0.119^{+0.099}_{-0.068}$	1.212
ℓ range	ℓ_{eff}	$A_s^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_s^{BB}	$A_d^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_d^{BB}	ρ^{BB}	χ_{red}^2
20–39	29.0	$47.841^{+1.913}_{-1.914}$	$-3.010^{+0.059}_{-0.059}$	$22.650^{+0.173}_{-0.177}$	$1.509^{+0.010}_{-0.010}$	$0.117^{+0.014}_{-0.014}$	1.389
40–59	49.0	$9.572^{+0.753}_{-0.772}$	$-3.125^{+0.112}_{-0.110}$	$6.111^{+0.052}_{-0.052}$	$1.493^{+0.012}_{-0.011}$	$0.104^{+0.020}_{-0.020}$	1.394
60–79	69.0	$1.185^{+0.492}_{-0.516}$	$-3.154^{+0.272}_{-0.254}$	$3.378^{+0.023}_{-0.024}$	$1.550^{+0.012}_{-0.011}$	$0.088^{+0.055}_{-0.042}$	1.562
80–99	89.0	$0.718^{+0.438}_{-0.408}$	$-2.988^{+0.232}_{-0.239}$	$2.010^{+0.016}_{-0.016}$	$1.552^{+0.013}_{-0.013}$	$0.153^{+0.096}_{-0.055}$	1.360
100–119	109.0	$0.829^{+0.454}_{-0.446}$	$-2.697^{+0.246}_{-0.259}$	$1.325^{+0.011}_{-0.012}$	$1.572^{+0.015}_{-0.015}$	$0.104^{+0.071}_{-0.049}$	1.156
120–139	129.0	$1.101^{+0.417}_{-0.440}$	$-2.965^{+0.247}_{-0.237}$	$0.826^{+0.008}_{-0.009}$	$1.540^{+0.018}_{-0.017}$	$0.143^{+0.062}_{-0.044}$	1.695
140–159	149.0	$0.152^{+0.351}_{-0.129}$	$-2.976^{+0.333}_{-0.309}$	$0.527^{+0.007}_{-0.007}$	$1.514^{+0.023}_{-0.022}$	$-0.029^{+0.141}_{-0.154}$	1.153
160–179	169.0	$1.057^{+0.486}_{-0.483}$	$-3.070^{+0.319}_{-0.310}$	$0.421^{+0.006}_{-0.006}$	$1.533^{+0.025}_{-0.024}$	$-0.054^{+0.051}_{-0.056}$	1.475
180–199	189.0	$1.095^{+0.653}_{-0.586}$	$-3.169^{+0.326}_{-0.320}$	$0.365^{+0.005}_{-0.005}$	$1.488^{+0.024}_{-0.023}$	$0.068^{+0.079}_{-0.065}$	1.646

Table 3: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes, frequency spectral indices, and synchrotron-dust correlation coefficient ρ , fitted independently per multipole bin from the EE (top) and BB (bottom) polarization auto- and cross-spectra of the 14-band QUIJOTE+WMAP+Planck data set, using the $|b| > 15^\circ$ Galactic mask.

ℓ range	ℓ_{eff}	$A_s^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_s^{EE}	$A_d^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_d^{EE}	ρ^{EE}	χ_{red}^2
20–39	29.0	$276.355^{+3.451}_{-3.415}$	$-3.078^{+0.019}_{-0.019}$	$33.564^{+0.235}_{-0.242}$	$1.467^{+0.009}_{-0.010}$	$0.078^{+0.008}_{-0.008}$	2.025
40–59	49.0	$32.205^{+1.230}_{-1.292}$	$-3.107^{+0.056}_{-0.056}$	$4.943^{+0.069}_{-0.071}$	$1.584^{+0.020}_{-0.019}$	$0.046^{+0.016}_{-0.016}$	1.039
60–79	69.0	$10.944^{+0.757}_{-0.774}$	$-3.234^{+0.084}_{-0.083}$	$2.612^{+0.033}_{-0.034}$	$1.514^{+0.019}_{-0.019}$	$-0.007^{+0.019}_{-0.018}$	1.150
80–99	89.0	$5.791^{+0.602}_{-0.638}$	$-3.148^{+0.120}_{-0.119}$	$1.691^{+0.020}_{-0.021}$	$1.519^{+0.019}_{-0.018}$	$0.080^{+0.023}_{-0.022}$	1.418
100–119	109.0	$3.234^{+0.489}_{-0.500}$	$-2.907^{+0.170}_{-0.170}$	$1.118^{+0.014}_{-0.014}$	$1.552^{+0.021}_{-0.022}$	$0.097^{+0.030}_{-0.029}$	1.294
120–139	129.0	$2.109^{+0.515}_{-0.508}$	$-2.815^{+0.247}_{-0.243}$	$0.623^{+0.011}_{-0.011}$	$1.380^{+0.029}_{-0.027}$	$0.123^{+0.049}_{-0.044}$	1.390
140–159	149.0	$1.952^{+0.606}_{-0.609}$	$-2.822^{+0.259}_{-0.273}$	$0.461^{+0.009}_{-0.009}$	$1.484^{+0.033}_{-0.033}$	$0.148^{+0.058}_{-0.050}$	1.058
160–179	169.0	$1.552^{+0.564}_{-0.546}$	$-2.991^{+0.280}_{-0.273}$	$0.363^{+0.007}_{-0.007}$	$1.568^{+0.035}_{-0.034}$	$0.030^{+0.063}_{-0.059}$	0.932
180–199	189.0	$0.815^{+0.689}_{-0.580}$	$-2.909^{+0.305}_{-0.305}$	$0.288^{+0.007}_{-0.007}$	$1.656^{+0.045}_{-0.046}$	$0.111^{+0.137}_{-0.085}$	1.184
ℓ range	ℓ_{eff}	$A_s^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_s^{BB}	$A_d^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_d^{BB}	ρ^{BB}	χ_{red}^2
20–39	29.0	$41.336^{+2.018}_{-2.036}$	$-3.086^{+0.070}_{-0.068}$	$13.185^{+0.145}_{-0.146}$	$1.484^{+0.014}_{-0.014}$	$0.164^{+0.017}_{-0.017}$	1.514
40–59	49.0	$8.650^{+0.794}_{-0.812}$	$-3.092^{+0.127}_{-0.127}$	$4.407^{+0.044}_{-0.046}$	$1.494^{+0.014}_{-0.014}$	$0.041^{+0.022}_{-0.021}$	1.672
60–79	69.0	$0.866^{+0.520}_{-0.506}$	$-3.118^{+0.261}_{-0.262}$	$2.148^{+0.019}_{-0.019}$	$1.571^{+0.015}_{-0.015}$	$0.121^{+0.091}_{-0.057}$	1.405
80–99	89.0	$0.832^{+0.487}_{-0.471}$	$-2.812^{+0.241}_{-0.258}$	$1.119^{+0.014}_{-0.014}$	$1.560^{+0.020}_{-0.020}$	$0.125^{+0.093}_{-0.062}$	1.172
100–119	109.0	$0.953^{+0.501}_{-0.506}$	$-2.754^{+0.269}_{-0.290}$	$0.688^{+0.009}_{-0.009}$	$1.577^{+0.024}_{-0.024}$	$0.041^{+0.060}_{-0.053}$	1.025
120–139	129.0	$1.012^{+0.441}_{-0.437}$	$-3.018^{+0.271}_{-0.270}$	$0.393^{+0.007}_{-0.007}$	$1.557^{+0.034}_{-0.033}$	$0.091^{+0.066}_{-0.055}$	1.524
140–159	149.0	$0.186^{+0.382}_{-0.156}$	$-2.968^{+0.305}_{-0.309}$	$0.308^{+0.005}_{-0.005}$	$1.521^{+0.033}_{-0.032}$	$0.023^{+0.163}_{-0.135}$	1.193
160–179	169.0	$1.104^{+0.471}_{-0.454}$	$-3.229^{+0.277}_{-0.278}$	$0.237^{+0.006}_{-0.005}$	$1.510^{+0.042}_{-0.041}$	$-0.097^{+0.061}_{-0.068}$	1.258
180–199	189.0	$1.859^{+0.642}_{-0.660}$	$-2.877^{+0.249}_{-0.271}$	$0.190^{+0.005}_{-0.005}$	$1.512^{+0.044}_{-0.042}$	$-0.004^{+0.063}_{-0.061}$	1.337

Table 4: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes, frequency spectral indices, and synchrotron-dust correlation coefficient ρ , fitted independently per multipole bin from the EE (top) and BB (bottom) polarization auto- and cross-spectra of the 14-band QUIJOTE+WMAP+Planck data set, using the $|b| > 20^\circ$ Galactic mask.

ℓ range	ℓ_{eff}	$A_s^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_s^{EE}	$A_d^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_d^{EE}	ρ^{EE}	χ_{red}^2
20–39	29.0	$263.649^{+3.842}_{-3.803}$	$-3.103^{+0.021}_{-0.021}$	$20.685^{+0.229}_{-0.224}$	$1.472^{+0.013}_{-0.014}$	$0.009^{+0.010}_{-0.009}$	1.636
40–59	49.0	$32.168^{+1.267}_{-1.293}$	$-3.119^{+0.056}_{-0.055}$	$4.137^{+0.062}_{-0.061}$	$1.508^{+0.020}_{-0.020}$	$0.067^{+0.015}_{-0.015}$	1.526
60–79	69.0	$10.382^{+0.738}_{-0.721}$	$-3.210^{+0.085}_{-0.085}$	$1.624^{+0.029}_{-0.029}$	$1.446^{+0.025}_{-0.025}$	$-0.091^{+0.022}_{-0.022}$	1.096
80–99	89.0	$5.343^{+0.607}_{-0.608}$	$-3.144^{+0.128}_{-0.130}$	$1.178^{+0.018}_{-0.018}$	$1.485^{+0.024}_{-0.024}$	$0.062^{+0.025}_{-0.024}$	1.263
100–119	109.0	$2.461^{+0.513}_{-0.513}$	$-2.950^{+0.200}_{-0.205}$	$0.820^{+0.014}_{-0.014}$	$1.564^{+0.030}_{-0.029}$	$0.041^{+0.038}_{-0.036}$	1.189
120–139	129.0	$1.955^{+0.504}_{-0.498}$	$-2.783^{+0.237}_{-0.242}$	$0.436^{+0.010}_{-0.010}$	$1.320^{+0.036}_{-0.035}$	$0.180^{+0.059}_{-0.052}$	1.317
140–159	149.0	$2.391^{+0.584}_{-0.601}$	$-2.672^{+0.219}_{-0.236}$	$0.325^{+0.009}_{-0.009}$	$1.450^{+0.046}_{-0.045}$	$0.097^{+0.052}_{-0.047}$	1.069
160–179	169.0	$1.769^{+0.586}_{-0.569}$	$-3.016^{+0.279}_{-0.274}$	$0.265^{+0.007}_{-0.007}$	$1.543^{+0.047}_{-0.045}$	$0.027^{+0.064}_{-0.062}$	0.998
180–199	189.0	$0.829^{+0.681}_{-0.572}$	$-2.911^{+0.299}_{-0.300}$	$0.189^{+0.006}_{-0.006}$	$1.625^{+0.069}_{-0.066}$	$0.122^{+0.145}_{-0.099}$	1.077
ℓ range	ℓ_{eff}	$A_s^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_s^{BB}	$A_d^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_d^{BB}	ρ^{BB}	χ_{red}^2
20–39	29.0	$39.584^{+1.976}_{-1.985}$	$-3.039^{+0.072}_{-0.072}$	$8.333^{+0.113}_{-0.113}$	$1.511^{+0.020}_{-0.019}$	$0.171^{+0.020}_{-0.020}$	1.804
40–59	49.0	$8.077^{+0.815}_{-0.851}$	$-3.086^{+0.142}_{-0.134}$	$2.945^{+0.036}_{-0.037}$	$1.480^{+0.019}_{-0.018}$	$0.035^{+0.028}_{-0.027}$	1.368
60–79	69.0	$0.888^{+0.547}_{-0.533}$	$-2.963^{+0.289}_{-0.288}$	$1.345^{+0.017}_{-0.019}$	$1.607^{+0.024}_{-0.023}$	$0.185^{+0.121}_{-0.072}$	1.464
80–99	89.0	$0.747^{+0.520}_{-0.469}$	$-2.872^{+0.254}_{-0.267}$	$0.763^{+0.012}_{-0.012}$	$1.579^{+0.028}_{-0.029}$	$0.079^{+0.094}_{-0.070}$	1.517
100–119	109.0	$1.143^{+0.550}_{-0.577}$	$-2.750^{+0.280}_{-0.290}$	$0.461^{+0.008}_{-0.008}$	$1.527^{+0.032}_{-0.030}$	$0.046^{+0.064}_{-0.059}$	1.016
120–139	129.0	$0.993^{+0.471}_{-0.454}$	$-3.033^{+0.268}_{-0.263}$	$0.278^{+0.006}_{-0.006}$	$1.587^{+0.044}_{-0.044}$	$0.106^{+0.081}_{-0.064}$	1.419
140–159	149.0	$0.235^{+0.461}_{-0.198}$	$-2.954^{+0.308}_{-0.308}$	$0.235^{+0.005}_{-0.005}$	$1.556^{+0.042}_{-0.041}$	$-0.004^{+0.155}_{-0.144}$	1.239
160–179	169.0	$1.332^{+0.527}_{-0.510}$	$-3.149^{+0.269}_{-0.269}$	$0.173^{+0.005}_{-0.005}$	$1.513^{+0.054}_{-0.054}$	$-0.082^{+0.062}_{-0.067}$	1.092
180–199	189.0	$2.067^{+0.669}_{-0.686}$	$-2.741^{+0.218}_{-0.245}$	$0.137^{+0.005}_{-0.005}$	$1.519^{+0.061}_{-0.060}$	$-0.077^{+0.064}_{-0.067}$	1.197

6 CONCLUSIONS

REFERENCES

- Adak D., et al., 2025, [arXiv eprints](#), p. [arXiv:2510.17761](#)
- Ade P., et al., 2021, [Physical Review Letters](#), 127
- Allys E., et al., 2022, [Progress of Theoretical and Experimental Physics](#), 2023
- Alonso D., et al., 2019, [Monthly Notices of the Royal Astronomical Society](#), 484, 4127–4151
- Bennett C. L., et al., 2013, [The Astrophysical Journal Supplement Series](#), 208, 20
- Blandford R. D., Znajek R. L., 1977, [MNRAS](#), 179, 433
- Casas J. M., et al., 2025, [arXiv eprints](#), p. [arXiv:2510.12860](#)
- Choi S. K., Page L. A., 2015, [Journal of Cosmology and Astroparticle Physics](#), 2015, 020–020
- Draine B. T., et al., 2016, [The Astrophysical Journal](#), 831, 59
- Fernández-Torreiro M., et al., 2023, [MNRAS](#), 526, 1343
- Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, [Publications of the Astronomical Society of the Pacific](#), 125, 306–312
- Fuskeland U., et al., 2021, [Astronomy & Astrophysics](#), 646, A69
- Gold B., et al., 2011, [ApJS](#), 192, 15
- González-González R., et al., 2025, [A&A](#), 695, A245
- Górski K. M., Hivon E., Banday A. J., Wandelt B. D., Hansen F. K., Reinecke M., Bartelmann M., 2005, [ApJ](#), 622, 759
- Génova-Santos R., et al., 2017, [Monthly Notices of the Royal Astronomical Society](#), 464, 4107–4132
- Hensley B. S., et al., 2022, [The Astrophysical Journal](#), 929, 166
- Herman, D. et al., 2023, [A&A](#), 675, A15
- Hinshaw G., et al., 2013, [The Astrophysical Journal Supplement Series](#), 208, 19
- Huynh M., Lazio J., 2013, An Overview of the Square Kilometre Array ([arXiv:1311.4288](#))
- Ishibashi, W. Courvoisier, T. J.-L. 2011, [A&A](#), 525, A118
- Kamionkowski M., 1999, [Nuclear Physics B - Proceedings Supplements](#), 70, 529–536
- Kosowsky A., 1996, [Annals of Physics](#), 246, 49–85
- Krachmalnicoff N., et al., 2018, [Astronomy & Astrophysics](#), 618, A166
- Martire F., Barreiro R., Martínez-González E., 2022, [Journal of Cosmology and Astroparticle Physics](#), 2022, 003
- Næss S. K., Louis T., 2013, [Journal of Cosmology and Astroparticle Physics](#), 2013, 001–001
- Padovani M., Bracco A., Jelić V., Galli D., Bellomi E., 2021, [Astronomy & Astrophysics](#), 651, A116
- Peel M. W., Genova-Santos R., Dickinson C., Leahy J. P., López-Caraballo C., Fernández-Torreiro M., Rubiño-Martín J. A., Spencer L. D., 2022, [Research Notes of the AAS](#), 6, 252
- Planck Collaboration 2015, [Astronomy & Astrophysics](#), 576, A104

- Planck Collaboration 2016a, [Astronomy & Astrophysics](#), 594, A2
- Planck Collaboration 2016b, [Astronomy & Astrophysics](#), 594, A10
- Planck Collaboration 2020, [A&A](#), 641, A6
- Planck Collaboration et al., 2015, [A&A](#), 576, A107
- Planck Collaboration et al., 2016, [A&A](#), 586, A133
- Planck Collaboration et al., 2020a, [A&A](#), 641, A3
- Planck Collaboration et al., 2020b, [A&A](#), 641, A11
- Planck Collaboration et al., 2020c, [A&A](#), 643, A42
- Rubiño-Martín J. A., et al., 2023, [MNRAS](#), 519, 3383
- Rybicki G. B., Lightman A. P., 1979, Radiative processes in astrophysics
- Smith K. M., 2006, [New Astronomy Reviews](#), 50, 1025
- Thorne B., Dunkley J., Alonso D., Naess S., 2017, [Monthly Notices of the Royal Astronomical Society](#), 469, 2821–2833
- de la Hoz E., et al., 2023, [Monthly Notices of the Royal Astronomical Society](#), 519, 3504–3525

A Analysis fitting only cross-spectra

1.1. Power-law fitting

Table 5: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes, spectral indices, and synchrotron-dust correlation coefficient ρ , from a global power-law fit to the EE and BB polarization cross-spectra of the 14-band QUIJOTE+WMAP+Planck dataset over the multipole range $20 \leq \ell \leq 199$, for the three Galactic masks.

Mask	Dataset	Mode	$A_s [10^{-3} \mu\text{K}^2]$	α_s	β_s	$A_d [10^{-3} \mu\text{K}^2]$	α_d	β_d	ρ
$ b > 10^\circ$	WP+P1	EE	$7.196^{+0.572}_{-0.547}$	$-3.716^{+0.076}_{-0.079}$	$-3.177^{+0.065}_{-0.067}$	$3.178^{+0.019}_{-0.019}$	$-2.542^{+0.006}_{-0.006}$	$1.520^{+0.008}_{-0.008}$	$0.095^{+0.007}_{-0.007}$
		BB	$2.273^{+0.494}_{-0.437}$	$-3.117^{+0.191}_{-0.202}$	$-3.296^{+0.208}_{-0.232}$	$2.394^{+0.014}_{-0.014}$	$-2.204^{+0.005}_{-0.005}$	$1.555^{+0.008}_{-0.008}$	$0.132^{+0.013}_{-0.013}$
	QJ+WP+P1	EE	$7.418^{+0.494}_{-0.287}$	$-3.623^{+0.042}_{-0.042}$	$-3.088^{+0.019}_{-0.020}$	$3.181^{+0.019}_{-0.019}$	$-2.542^{+0.006}_{-0.006}$	$1.521^{+0.008}_{-0.007}$	$0.095^{+0.005}_{-0.005}$
		BB	$1.703^{+0.206}_{-0.200}$	$-3.291^{+0.122}_{-0.131}$	$-2.960^{+0.061}_{-0.061}$	$2.395^{+0.014}_{-0.014}$	$-2.204^{+0.005}_{-0.005}$	$1.556^{+0.008}_{-0.008}$	$0.114^{+0.011}_{-0.011}$
$ b > 15^\circ$	WP+P1	EE	$5.727^{+0.580}_{-0.549}$	$-3.875^{+0.095}_{-0.100}$	$-3.182^{+0.073}_{-0.076}$	$2.066^{+0.017}_{-0.017}$	$-2.514^{+0.008}_{-0.008}$	$1.535^{+0.011}_{-0.011}$	$0.081^{+0.008}_{-0.008}$
		BB	$2.163^{+0.568}_{-0.498}$	$-3.180^{+0.228}_{-0.254}$	$-3.585^{+0.286}_{-0.327}$	$1.379^{+0.012}_{-0.012}$	$-2.323^{+0.008}_{-0.008}$	$1.539^{+0.012}_{-0.011}$	$0.123^{+0.015}_{-0.015}$
	QJ+WP+P1	EE	$5.998^{+0.498}_{-0.287}$	$-3.764^{+0.052}_{-0.051}$	$-3.086^{+0.021}_{-0.022}$	$2.066^{+0.017}_{-0.017}$	$-2.514^{+0.008}_{-0.008}$	$1.534^{+0.011}_{-0.011}$	$0.068^{+0.006}_{-0.007}$
		BB	$1.363^{+0.218}_{-0.208}$	$-3.405^{+0.157}_{-0.171}$	$-3.003^{+0.073}_{-0.074}$	$1.380^{+0.012}_{-0.012}$	$-2.323^{+0.008}_{-0.008}$	$1.541^{+0.011}_{-0.011}$	$0.118^{+0.013}_{-0.013}$
$ b > 20^\circ$	WP+P1	EE	$6.864^{+0.654}_{-0.628}$	$-3.717^{+0.092}_{-0.098}$	$-3.379^{+0.091}_{-0.096}$	$1.410^{+0.015}_{-0.015}$	$-2.475^{+0.010}_{-0.010}$	$1.460^{+0.013}_{-0.013}$	$0.032^{+0.009}_{-0.009}$
		BB	$1.805^{+0.584}_{-0.492}$	$-3.421^{+0.275}_{-0.310}$	$-3.885^{+0.331}_{-0.376}$	$0.925^{+0.010}_{-0.010}$	$-2.242^{+0.010}_{-0.010}$	$1.534^{+0.016}_{-0.016}$	$0.154^{+0.019}_{-0.018}$
	QJ+WP+P1	EE	$5.908^{+0.492}_{-0.286}$	$-3.726^{+0.052}_{-0.053}$	$-3.129^{+0.024}_{-0.024}$	$1.411^{+0.015}_{-0.015}$	$-2.475^{+0.010}_{-0.010}$	$1.461^{+0.013}_{-0.013}$	$0.022^{+0.007}_{-0.007}$
		BB	$1.358^{+0.235}_{-0.220}$	$-3.326^{+0.170}_{-0.183}$	$-2.978^{+0.075}_{-0.077}$	$0.926^{+0.011}_{-0.010}$	$-2.242^{+0.010}_{-0.010}$	$1.539^{+0.016}_{-0.016}$	$0.123^{+0.015}_{-0.015}$




Figure 5: Best-fit global power-law models for synchrotron and thermal dust emission using cross-only spectra for the fit, overlaid on the measured EE and BB frequency power spectra, for the $|b| > 10^\circ$ Galactic mask.

1.2. Bin-to-bin fitting



Figure 6: Same as Figure 5 but for the $|b| > 20^\circ$ Galactic mask.Figure 7: Same as Figure 5 but for the $|b| > 20^\circ$ Galactic mask.

Table 6: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes, frequency spectral indices, and synchrotron-dust correlation coefficient ρ , fitted independently per multipole bin from the EE (top) and BB (bottom) polarization cross-spectra of the 14-band QUIJOTE+WMAP+Planck data set, using the $|b| > 10^\circ$ Galactic mask.

ℓ range	ℓ_{eff}	$A_s^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_s^{EE}	$A_d^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_d^{EE}	ρ^{EE}	χ_{red}^2
20–39	29.0	$296.777^{+3.753}_{-3.784}$	$-3.078^{+0.022}_{-0.022}$	$48.896^{+0.490}_{-0.485}$	$1.477^{+0.011}_{-0.011}$	$0.091^{+0.007}_{-0.007}$	1.328
40–59	49.0	$37.435^{+1.334}_{-1.339}$	$-3.148^{+0.057}_{-0.057}$	$7.813^{+0.126}_{-0.125}$	$1.507^{+0.019}_{-0.019}$	$0.117^{+0.013}_{-0.012}$	1.186
60–79	69.0	$12.839^{+0.855}_{-0.888}$	$-3.304^{+0.095}_{-0.096}$	$4.820^{+0.080}_{-0.080}$	$1.537^{+0.020}_{-0.019}$	$0.046^{+0.014}_{-0.014}$	1.072
80–99	89.0	$6.384^{+0.635}_{-0.652}$	$-3.126^{+0.133}_{-0.135}$	$2.703^{+0.041}_{-0.040}$	$1.525^{+0.021}_{-0.020}$	$0.055^{+0.020}_{-0.019}$	1.476
100–119	109.0	$3.714^{+0.531}_{-0.554}$	$-3.013^{+0.179}_{-0.185}$	$1.529^{+0.033}_{-0.033}$	$1.550^{+0.031}_{-0.030}$	$0.033^{+0.026}_{-0.026}$	0.968
120–139	129.0	$2.833^{+0.514}_{-0.534}$	$-2.582^{+0.216}_{-0.215}$	$0.977^{+0.024}_{-0.023}$	$1.368^{+0.034}_{-0.032}$	$0.156^{+0.040}_{-0.038}$	1.457
140–159	149.0	$2.106^{+0.607}_{-0.574}$	$-2.974^{+0.237}_{-0.248}$	$0.677^{+0.017}_{-0.016}$	$1.472^{+0.039}_{-0.038}$	$0.218^{+0.052}_{-0.044}$	1.037
160–179	169.0	$1.386^{+0.527}_{-0.522}$	$-3.168^{+0.298}_{-0.283}$	$0.472^{+0.017}_{-0.016}$	$1.522^{+0.050}_{-0.047}$	$0.072^{+0.065}_{-0.056}$	1.036
180–199	189.0	$1.295^{+0.689}_{-0.671}$	$-2.902^{+0.294}_{-0.306}$	$0.413^{+0.017}_{-0.016}$	$1.685^{+0.062}_{-0.062}$	$0.113^{+0.086}_{-0.063}$	1.236
ℓ range	ℓ_{eff}	$A_s^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_s^{BB}	$A_d^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_d^{BB}	ρ^{BB}	χ_{red}^2
20–39	29.0	$46.872^{+2.152}_{-2.089}$	$-3.014^{+0.069}_{-0.070}$	$22.853^{+0.315}_{-0.301}$	$1.526^{+0.017}_{-0.016}$	$0.120^{+0.014}_{-0.014}$	1.340
40–59	49.0	$9.530^{+0.800}_{-0.804}$	$-2.969^{+0.122}_{-0.123}$	$6.130^{+0.094}_{-0.092}$	$1.503^{+0.019}_{-0.019}$	$0.111^{+0.022}_{-0.021}$	1.318
60–79	69.0	$1.411^{+0.534}_{-0.535}$	$-3.154^{+0.254}_{-0.258}$	$3.440^{+0.052}_{-0.052}$	$1.573^{+0.020}_{-0.020}$	$0.083^{+0.047}_{-0.038}$	1.616
80–99	89.0	$0.935^{+0.453}_{-0.451}$	$-3.013^{+0.232}_{-0.238}$	$2.048^{+0.031}_{-0.032}$	$1.584^{+0.023}_{-0.022}$	$0.139^{+0.072}_{-0.049}$	1.429
100–119	109.0	$0.792^{+0.476}_{-0.468}$	$-2.733^{+0.251}_{-0.270}$	$1.355^{+0.024}_{-0.023}$	$1.605^{+0.025}_{-0.025}$	$0.113^{+0.084}_{-0.051}$	1.217
120–139	129.0	$1.176^{+0.427}_{-0.424}$	$-3.019^{+0.246}_{-0.253}$	$0.832^{+0.016}_{-0.015}$	$1.559^{+0.029}_{-0.028}$	$0.137^{+0.055}_{-0.043}$	1.607
140–159	149.0	$0.320^{+0.495}_{-0.268}$	$-2.952^{+0.317}_{-0.314}$	$0.541^{+0.014}_{-0.014}$	$1.555^{+0.039}_{-0.038}$	$-0.016^{+0.098}_{-0.104}$	1.156
160–179	169.0	$0.536^{+0.505}_{-0.413}$	$-3.063^{+0.308}_{-0.308}$	$0.439^{+0.013}_{-0.013}$	$1.606^{+0.045}_{-0.044}$	$-0.064^{+0.074}_{-0.103}$	1.410
180–199	189.0	$0.794^{+0.586}_{-0.506}$	$-3.283^{+0.307}_{-0.296}$	$0.404^{+0.011}_{-0.010}$	$1.651^{+0.043}_{-0.043}$	$0.103^{+0.097}_{-0.064}$	1.375

Table 7: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes, frequency spectral indices, and synchrotron-dust correlation coefficient ρ , fitted independently per multipole bin from the EE (top) and BB (bottom) polarization cross-spectra of the 14-band QUIJOTE+WMAP+Planck data set, using the $|b| > 15^\circ$ Galactic mask.

ℓ range	ℓ_{eff}	$A_s^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_s^{EE}	$A_d^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_d^{EE}	ρ^{EE}	χ_{red}^2
20–39	29.0	$276.030^{+3.804}_{-3.920}$	$-3.088^{+0.023}_{-0.023}$	$32.899^{+0.460}_{-0.452}$	$1.443^{+0.016}_{-0.016}$	$0.077^{+0.008}_{-0.008}$	1.573
40–59	49.0	$31.665^{+1.343}_{-1.301}$	$-3.115^{+0.068}_{-0.069}$	$4.864^{+0.121}_{-0.118}$	$1.565^{+0.030}_{-0.030}$	$0.046^{+0.015}_{-0.015}$	0.975
60–79	69.0	$10.732^{+0.864}_{-0.867}$	$-3.327^{+0.111}_{-0.115}$	$2.644^{+0.060}_{-0.059}$	$1.529^{+0.030}_{-0.028}$	$-0.004^{+0.018}_{-0.018}$	1.077
80–99	89.0	$6.012^{+0.646}_{-0.648}$	$-3.153^{+0.137}_{-0.143}$	$1.708^{+0.036}_{-0.035}$	$1.541^{+0.029}_{-0.028}$	$0.079^{+0.021}_{-0.021}$	1.461
100–119	109.0	$3.355^{+0.532}_{-0.520}$	$-2.925^{+0.179}_{-0.186}$	$1.096^{+0.029}_{-0.028}$	$1.530^{+0.037}_{-0.037}$	$0.093^{+0.029}_{-0.028}$	1.239
120–139	129.0	$2.466^{+0.557}_{-0.563}$	$-2.738^{+0.246}_{-0.250}$	$0.600^{+0.019}_{-0.019}$	$1.333^{+0.043}_{-0.041}$	$0.109^{+0.045}_{-0.042}$	1.401
140–159	149.0	$2.033^{+0.648}_{-0.631}$	$-2.921^{+0.275}_{-0.280}$	$0.450^{+0.014}_{-0.014}$	$1.443^{+0.051}_{-0.048}$	$0.142^{+0.054}_{-0.049}$	1.114
160–179	169.0	$1.723^{+0.578}_{-0.562}$	$-3.042^{+0.271}_{-0.282}$	$0.369^{+0.016}_{-0.015}$	$1.598^{+0.064}_{-0.061}$	$0.035^{+0.058}_{-0.055}$	0.983
180–199	189.0	$0.969^{+0.733}_{-0.658}$	$-2.901^{+0.307}_{-0.307}$	$0.295^{+0.016}_{-0.015}$	$1.716^{+0.090}_{-0.086}$	$0.103^{+0.119}_{-0.078}$	1.239
ℓ range	ℓ_{eff}	$A_s^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_s^{BB}	$A_d^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_d^{BB}	ρ^{BB}	χ_{red}^2
20–39	29.0	$40.394^{+2.272}_{-2.243}$	$-3.130^{+0.083}_{-0.085}$	$13.261^{+0.253}_{-0.252}$	$1.497^{+0.022}_{-0.022}$	$0.168^{+0.017}_{-0.017}$	1.284
40–59	49.0	$8.505^{+0.860}_{-0.893}$	$-2.987^{+0.152}_{-0.146}$	$4.382^{+0.082}_{-0.086}$	$1.492^{+0.024}_{-0.023}$	$0.045^{+0.024}_{-0.023}$	1.596
60–79	69.0	$1.059^{+0.551}_{-0.556}$	$-3.099^{+0.261}_{-0.263}$	$2.176^{+0.043}_{-0.043}$	$1.586^{+0.027}_{-0.027}$	$0.112^{+0.078}_{-0.053}$	1.476
80–99	89.0	$0.933^{+0.494}_{-0.506}$	$-2.873^{+0.246}_{-0.262}$	$1.114^{+0.027}_{-0.027}$	$1.557^{+0.035}_{-0.035}$	$0.113^{+0.086}_{-0.058}$	1.275
100–119	109.0	$1.052^{+0.498}_{-0.501}$	$-2.853^{+0.278}_{-0.288}$	$0.694^{+0.018}_{-0.018}$	$1.592^{+0.040}_{-0.039}$	$0.038^{+0.056}_{-0.049}$	1.047
120–139	129.0	$1.146^{+0.443}_{-0.439}$	$-3.103^{+0.279}_{-0.281}$	$0.391^{+0.015}_{-0.014}$	$1.567^{+0.058}_{-0.056}$	$0.082^{+0.062}_{-0.050}$	1.331
140–159	149.0	$0.377^{+0.532}_{-0.310}$	$-2.953^{+0.307}_{-0.305}$	$0.335^{+0.014}_{-0.013}$	$1.660^{+0.067}_{-0.064}$	$0.032^{+0.119}_{-0.092}$	1.116
160–179	169.0	$0.610^{+0.509}_{-0.437}$	$-3.186^{+0.292}_{-0.278}$	$0.238^{+0.012}_{-0.011}$	$1.527^{+0.076}_{-0.072}$	$-0.126^{+0.085}_{-0.137}$	1.296
180–199	189.0	$1.485^{+0.677}_{-0.689}$	$-2.987^{+0.263}_{-0.280}$	$0.225^{+0.012}_{-0.011}$	$1.786^{+0.087}_{-0.082}$	$0.029^{+0.066}_{-0.061}$	1.123

Table 8: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes, frequency spectral indices, and synchrotron-dust correlation coefficient ρ , fitted independently per multipole bin from the EE (top) and BB (bottom) polarization cross-spectra of the 14-band QUIJOTE+WMAP+Planck data set, using the $|b| > 20^\circ$ Galactic mask.

ℓ range	ℓ_{eff}	$A_s^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_s^{EE}	$A_d^{\text{EE}} [10^{-3} \mu\text{K}^2]$	β_d^{EE}	ρ^{EE}	χ_{red}^2
20–39	29.0	$264.245^{+4.366}_{-4.499}$	$-3.116^{+0.027}_{-0.027}$	$20.371^{+0.406}_{-0.402}$	$1.456^{+0.021}_{-0.022}$	$0.009^{+0.010}_{-0.010}$	1.518
40–59	49.0	$31.624^{+1.382}_{-1.419}$	$-3.125^{+0.069}_{-0.071}$	$4.020^{+0.105}_{-0.106}$	$1.472^{+0.031}_{-0.031}$	$0.066^{+0.015}_{-0.015}$	1.503
60–79	69.0	$10.313^{+0.794}_{-0.813}$	$-3.266^{+0.112}_{-0.120}$	$1.631^{+0.050}_{-0.049}$	$1.446^{+0.039}_{-0.039}$	$-0.088^{+0.021}_{-0.022}$	0.992
80–99	89.0	$5.609^{+0.661}_{-0.676}$	$-3.129^{+0.153}_{-0.156}$	$1.172^{+0.034}_{-0.033}$	$1.493^{+0.040}_{-0.038}$	$0.062^{+0.023}_{-0.024}$	1.331
100–119	109.0	$2.531^{+0.546}_{-0.545}$	$-2.962^{+0.215}_{-0.224}$	$0.812^{+0.029}_{-0.029}$	$1.556^{+0.053}_{-0.050}$	$0.039^{+0.037}_{-0.035}$	1.110
120–139	129.0	$2.210^{+0.555}_{-0.543}$	$-2.773^{+0.250}_{-0.247}$	$0.412^{+0.017}_{-0.016}$	$1.240^{+0.057}_{-0.053}$	$0.163^{+0.056}_{-0.049}$	1.305
140–159	149.0	$2.518^{+0.630}_{-0.617}$	$-2.777^{+0.236}_{-0.252}$	$0.321^{+0.014}_{-0.014}$	$1.432^{+0.067}_{-0.065}$	$0.094^{+0.050}_{-0.047}$	1.120
160–179	169.0	$1.884^{+0.598}_{-0.566}$	$-3.110^{+0.293}_{-0.289}$	$0.274^{+0.015}_{-0.014}$	$1.611^{+0.084}_{-0.080}$	$0.038^{+0.059}_{-0.058}$	1.063
180–199	189.0	$0.973^{+0.714}_{-0.648}$	$-2.899^{+0.295}_{-0.303}$	$0.191^{+0.015}_{-0.013}$	$1.666^{+0.136}_{-0.125}$	$0.117^{+0.130}_{-0.094}$	1.099
ℓ range	ℓ_{eff}	$A_s^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_s^{BB}	$A_d^{\text{BB}} [10^{-3} \mu\text{K}^2]$	β_d^{BB}	ρ^{BB}	χ_{red}^2
20–39	29.0	$37.722^{+2.289}_{-2.316}$	$-3.070^{+0.085}_{-0.088}$	$8.376^{+0.211}_{-0.209}$	$1.523^{+0.032}_{-0.032}$	$0.177^{+0.021}_{-0.020}$	1.741
40–59	49.0	$7.944^{+0.884}_{-0.899}$	$-3.016^{+0.162}_{-0.160}$	$2.934^{+0.071}_{-0.068}$	$1.480^{+0.031}_{-0.029}$	$0.036^{+0.027}_{-0.027}$	1.258
60–79	69.0	$0.949^{+0.602}_{-0.586}$	$-2.977^{+0.278}_{-0.275}$	$1.318^{+0.038}_{-0.037}$	$1.568^{+0.042}_{-0.040}$	$0.174^{+0.129}_{-0.072}$	1.527
80–99	89.0	$0.906^{+0.528}_{-0.534}$	$-2.941^{+0.262}_{-0.271}$	$0.755^{+0.026}_{-0.025}$	$1.568^{+0.051}_{-0.048}$	$0.066^{+0.083}_{-0.065}$	1.670
100–119	109.0	$1.235^{+0.538}_{-0.528}$	$-2.874^{+0.283}_{-0.295}$	$0.466^{+0.016}_{-0.016}$	$1.543^{+0.052}_{-0.051}$	$0.049^{+0.060}_{-0.054}$	1.043
120–139	129.0	$1.116^{+0.468}_{-0.484}$	$-3.117^{+0.285}_{-0.279}$	$0.278^{+0.014}_{-0.014}$	$1.614^{+0.084}_{-0.082}$	$0.097^{+0.076}_{-0.060}$	1.160
140–159	149.0	$0.457^{+0.599}_{-0.375}$	$-2.938^{+0.305}_{-0.304}$	$0.272^{+0.014}_{-0.014}$	$1.794^{+0.091}_{-0.087}$	$0.017^{+0.115}_{-0.098}$	1.128
160–179	169.0	$0.840^{+0.578}_{-0.542}$	$-3.096^{+0.274}_{-0.276}$	$0.173^{+0.011}_{-0.010}$	$1.523^{+0.102}_{-0.095}$	$-0.103^{+0.080}_{-0.111}$	1.111
180–199	189.0	$1.683^{+0.697}_{-0.731}$	$-2.855^{+0.245}_{-0.271}$	$0.154^{+0.011}_{-0.010}$	$1.706^{+0.123}_{-0.113}$	$-0.054^{+0.069}_{-0.073}$	1.158

B Alternative multipole fitting range $30 \leq \ell \leq 199$

This appendix presents the results of the global power-law fitting analysis carried out over the multipole range $30 \leq \ell \leq 199$, as a robustness check against the baseline range $20 \leq \ell \leq 199$ adopted in Section 5. Starting at $\ell = 30$ is the conventional choice in the literature (Rubio-Martín et al., 2023; Martire et al., 2022), as it avoids the largest angular scales where residual systematic effects and foreground modelling uncertainties may be more significant.

However, as discussed in Section 5, excluding the first multipole bin ($20 \leq \ell \leq 39$, $\ell_{\text{eff}} = 29$) has a marked impact on the BB-mode power-law fit. The BB polarization power spectrum is intrinsically fainter than EE and falls steeply with multipole; as a result, the first bin carries a substantial fraction of the total constraining power on the BB angular spectral index α_s^{BB} . Without it, the remaining bins span a narrower dynamic range in signal amplitude, and the power-law slope becomes poorly constrained. This manifests as an artificial steepening of the inferred α_s^{BB} towards unphysically large negative values, accompanied by significantly enlarged posterior uncertainties. The effect is most pronounced in the $|b| > 10^\circ$ mask when QUIJOTE data are included, where the additional low-frequency sensitivity of QUIJOTE makes the model sensitive to the shape of the synchrotron angular spectrum, amplifying the impact of missing low- ℓ information. The inclusion of the first multipole bin in the baseline analysis resolves this degeneracy, yielding well-constrained and physically consistent estimates for both the EE and BB modes.

The results below are presented for both the auto+cross-spectra and cross-spectra-only configurations, and for the three Galactic masks, to allow a direct comparison with the baseline results of Section 5 and Appendix A.

2.1. Fitting all data (auto- and cross-spectra)

RESULTS AUTO+CROSS-SPECTRA – $30 \leq \ell \leq 199$

Table 9: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes, spectral indices, and synchrotron-dust correlation coefficient ρ , from a global power-law fit to the EE and BB polarization auto- and cross-spectra of the 14-band QUIJOTE+WMAP+Planck dataset over the multipole range $30 \leq \ell \leq 199$, for the three Galactic masks.

Mask	Dataset	Mode	$A_s [10^{-3} \mu\text{K}^2]$	α_s	β_s	$A_d [10^{-3} \mu\text{K}^2]$	α_d	β_d	ρ
$ b > 10^\circ$	WP+Pl	EE	$9.451^{+0.649}_{-0.644}$	$-2.845^{+0.146}_{-0.156}$	$-3.189^{+0.116}_{-0.120}$	$2.999^{+0.012}_{-0.012}$	$-2.244^{+0.006}_{-0.006}$	$1.523^{+0.006}_{-0.006}$	$0.090^{+0.011}_{-0.011}$
		BB	$2.589^{+0.534}_{-0.530}$	$-2.380^{+0.443}_{-0.506}$	$-3.402^{+0.295}_{-0.325}$	$2.336^{+0.008}_{-0.008}$	$-2.163^{+0.006}_{-0.006}$	$1.536^{+0.005}_{-0.005}$	$0.125^{+0.020}_{-0.019}$
	QJ+WP+Pl	EE	$8.806^{+0.325}_{-0.328}$	$-2.949^{+0.079}_{-0.083}$	$-3.148^{+0.036}_{-0.036}$	$2.999^{+0.012}_{-0.012}$	$-2.244^{+0.006}_{-0.006}$	$1.523^{+0.006}_{-0.006}$	$0.088^{+0.008}_{-0.008}$
		BB	$1.266^{+0.296}_{-0.284}$	$-4.074^{+0.449}_{-0.535}$	$-3.065^{+0.109}_{-0.110}$	$2.336^{+0.008}_{-0.008}$	$-2.163^{+0.006}_{-0.006}$	$1.535^{+0.005}_{-0.005}$	$0.103^{+0.016}_{-0.015}$
$ b > 15^\circ$	WP+Pl	EE	$8.697^{+0.678}_{-0.677}$	$-2.695^{+0.169}_{-0.181}$	$-3.234^{+0.138}_{-0.143}$	$1.889^{+0.011}_{-0.011}$	$-2.112^{+0.009}_{-0.009}$	$1.533^{+0.008}_{-0.008}$	$0.066^{+0.012}_{-0.012}$
		BB	$2.667^{+0.557}_{-0.569}$	$-1.946^{+0.497}_{-0.595}$	$-3.232^{+0.338}_{-0.383}$	$1.417^{+0.007}_{-0.007}$	$-2.419^{+0.009}_{-0.009}$	$1.543^{+0.008}_{-0.008}$	$0.088^{+0.021}_{-0.020}$
	QJ+WP+Pl	EE	$7.889^{+0.332}_{-0.335}$	$-2.816^{+0.094}_{-0.095}$	$-3.130^{+0.042}_{-0.042}$	$1.889^{+0.011}_{-0.011}$	$-2.111^{+0.009}_{-0.009}$	$1.533^{+0.008}_{-0.008}$	$0.051^{+0.009}_{-0.009}$
		BB	$1.266^{+0.371}_{-0.355}$	$-3.861^{+0.570}_{-0.697}$	$-2.986^{+0.136}_{-0.135}$	$1.416^{+0.007}_{-0.007}$	$-2.418^{+0.009}_{-0.009}$	$1.542^{+0.008}_{-0.008}$	$0.062^{+0.018}_{-0.017}$
$ b > 20^\circ$	WP+Pl	EE	$7.902^{+0.720}_{-0.716}$	$-2.957^{+0.203}_{-0.216}$	$-3.260^{+0.145}_{-0.151}$	$1.362^{+0.010}_{-0.010}$	$-2.205^{+0.011}_{-0.011}$	$1.487^{+0.011}_{-0.011}$	$0.056^{+0.013}_{-0.013}$
		BB	$2.675^{+0.588}_{-0.613}$	$-1.663^{+0.543}_{-0.661}$	$-3.055^{+0.350}_{-0.403}$	$0.949^{+0.006}_{-0.006}$	$-2.335^{+0.011}_{-0.011}$	$1.540^{+0.011}_{-0.011}$	$0.079^{+0.025}_{-0.023}$
	QJ+WP+Pl	EE	$7.218^{+0.349}_{-0.347}$	$-3.010^{+0.108}_{-0.111}$	$-3.134^{+0.044}_{-0.044}$	$1.362^{+0.010}_{-0.010}$	$-2.205^{+0.011}_{-0.011}$	$1.486^{+0.011}_{-0.011}$	$0.033^{+0.010}_{-0.010}$
		BB	$1.369^{+0.408}_{-0.401}$	$-3.561^{+0.601}_{-0.744}$	$-2.918^{+0.152}_{-0.152}$	$0.949^{+0.006}_{-0.006}$	$-2.335^{+0.011}_{-0.011}$	$1.539^{+0.010}_{-0.010}$	$0.066^{+0.021}_{-0.020}$

2.2. Fitting only cross-spectra

RESULTS CROSS-SPECTRA ONLY – $30 \leq \ell \leq 199$

Table 10: Posterior constraints (median and 68% credible intervals) on the synchrotron and dust amplitudes, spectral indices, and synchrotron-dust correlation coefficient ρ , from a global power-law fit to the EE and BB polarization cross-spectra of the 14-band QUIJOTE+WMAP+Planck dataset over the multipole range $30 \leq \ell \leq 199$, for the three Galactic masks.

Mask	Dataset	Mode	$A_s [10^{-3} \mu\text{K}^2]$	α_s	β_s	$A_d [10^{-3} \mu\text{K}^2]$	α_d	β_d	ρ
$ b > 10^\circ$	WP+Pl	EE	$11.794^{+1.193}_{-1.126}$	$-2.661^{+0.164}_{-0.172}$	$-3.550^{+0.203}_{-0.225}$	$2.962^{+0.022}_{-0.022}$	$-2.240^{+0.008}_{-0.008}$	$1.512^{+0.010}_{-0.009}$	$0.088^{+0.010}_{-0.010}$
		BB	$3.854^{+1.018}_{-0.885}$	$-2.075^{+0.402}_{-0.447}$	$-3.981^{+0.463}_{-0.553}$	$2.372^{+0.016}_{-0.016}$	$-2.160^{+0.008}_{-0.008}$	$1.561^{+0.009}_{-0.009}$	$0.113^{+0.019}_{-0.017}$
	QJ+WP+Pl	EE	$9.027^{+0.360}_{-0.362}$	$-2.868^{+0.089}_{-0.090}$	$-3.170^{+0.046}_{-0.047}$	$2.967^{+0.022}_{-0.022}$	$-2.240^{+0.008}_{-0.008}$	$1.515^{+0.010}_{-0.009}$	$0.087^{+0.008}_{-0.008}$
		BB	$1.642^{+0.319}_{-0.321}$	$-3.482^{+0.406}_{-0.470}$	$-2.891^{+0.120}_{-0.119}$	$2.373^{+0.016}_{-0.016}$	$-2.159^{+0.008}_{-0.008}$	$1.564^{+0.009}_{-0.009}$	$0.106^{+0.016}_{-0.015}$
$ b > 15^\circ$	WP+Pl	EE	$11.700^{+1.432}_{-1.278}$	$-2.396^{+0.177}_{-0.190}$	$-3.737^{+0.265}_{-0.306}$	$1.872^{+0.019}_{-0.019}$	$-2.110^{+0.011}_{-0.011}$	$1.525^{+0.014}_{-0.013}$	$0.066^{+0.011}_{-0.011}$
		BB	$4.234^{+1.211}_{-1.041}$	$-1.638^{+0.464}_{-0.515}$	$-4.233^{+0.637}_{-0.770}$	$1.422^{+0.014}_{-0.013}$	$-2.420^{+0.012}_{-0.012}$	$1.551^{+0.013}_{-0.013}$	$0.081^{+0.019}_{-0.017}$
	QJ+WP+Pl	EE	$8.104^{+0.364}_{-0.363}$	$-2.706^{+0.100}_{-0.104}$	$-3.151^{+0.055}_{-0.055}$	$1.873^{+0.019}_{-0.019}$	$-2.110^{+0.011}_{-0.011}$	$1.526^{+0.013}_{-0.013}$	$0.051^{+0.009}_{-0.009}$
		BB	$1.685^{+0.367}_{-0.387}$	$-3.171^{+0.480}_{-0.594}$	$-2.857^{+0.138}_{-0.145}$	$1.423^{+0.014}_{-0.014}$	$-2.420^{+0.012}_{-0.012}$	$1.553^{+0.013}_{-0.013}$	$0.063^{+0.017}_{-0.017}$
$ b > 20^\circ$	WP+Pl	EE	$11.296^{+1.558}_{-1.375}$	$-2.647^{+0.211}_{-0.228}$	$-3.901^{+0.304}_{-0.352}$	$1.334^{+0.017}_{-0.017}$	$-2.211^{+0.015}_{-0.014}$	$1.460^{+0.017}_{-0.017}$	$0.055^{+0.012}_{-0.012}$
		BB	$4.148^{+1.380}_{-1.156}$	$-1.387^{+0.535}_{-0.621}$	$-4.149^{+0.743}_{-0.914}$	$0.950^{+0.012}_{-0.012}$	$-2.337^{+0.015}_{-0.015}$	$1.543^{+0.018}_{-0.018}$	$0.075^{+0.021}_{-0.019}$
	QJ+WP+Pl	EE	$7.481^{+0.376}_{-0.378}$	$-2.880^{+0.115}_{-0.120}$	$-3.154^{+0.057}_{-0.057}$	$1.334^{+0.017}_{-0.017}$	$-2.210^{+0.014}_{-0.014}$	$1.461^{+0.017}_{-0.017}$	$0.032^{+0.010}_{-0.010}$
		BB	$1.760^{+0.384}_{-0.407}$	$-2.906^{+0.506}_{-0.613}$	$-2.827^{+0.154}_{-0.161}$	$0.952^{+0.012}_{-0.012}$	$-2.337^{+0.016}_{-0.016}$	$1.547^{+0.018}_{-0.018}$	$0.065^{+0.020}_{-0.020}$

C Reduced χ^2 values of the power-law fittings

Table 11: Reduced chi-squared values χ^2_{red} for all global power-law fits, combining the two multipole ranges ($20 \leq \ell \leq 199$ and $30 \leq \ell \leq 199$), both spectral datasets (auto- and cross-spectra, and cross-spectra only), both dataset combinations (WP+Pl and QJ+WP+Pl), and the three Galactic masks.

Mask	Dataset	$20 \leq \ell \leq 199$				$30 \leq \ell \leq 199$			
		Auto+Cross		Cross		Auto+Cross		Cross	
		EE	BB	EE	BB	EE	BB	EE	BB
$ b > 10^\circ$	WP+Pl	10.15	2.91	7.48	2.28	3.99	2.93	3.14	2.27
	QJ+WP+Pl	7.90	2.50	5.83	2.01	3.27	2.51	2.61	2.00
$ b > 15^\circ$	WP+Pl	8.72	1.95	6.46	1.63	2.22	1.63	1.92	1.48
	QJ+WP+Pl	6.85	1.84	5.10	1.59	1.95	1.57	1.73	1.45
$ b > 20^\circ$	WP+Pl	3.36	1.70	2.64	1.56	1.83	1.47	1.57	1.39
	QJ+WP+Pl	2.85	1.60	2.32	1.49	1.69	1.42	1.52	1.36