



Castilla-La Mancha

VACCINATION

|                                                                | Prenatal | 2 months | 4 months     | 6 months   | 11 months | 12 months    | 15 months | 4 years | 5 years | 6 years | 12 years    | 14 years | 19 -64 years | ≥ 60 years    | 65 years | 80 years |
|----------------------------------------------------------------|----------|----------|--------------|------------|-----------|--------------|-----------|---------|---------|---------|-------------|----------|--------------|---------------|----------|----------|
| Hepatitis B <sup>1</sup>                                       |          | HB       | HB           |            | HB        |              |           |         |         |         |             |          |              |               |          |          |
| Diphtheria, Tetanus<br>Pertussis (whooping cough) <sup>2</sup> | dTpa     | DTPa     | DTPa         |            | DTPa      |              |           |         |         | DTPa    |             | Td       | Td           |               | Td       |          |
| Poliomyelitis                                                  |          | VPI      | VPI          |            | VPI       |              |           |         |         | VPI     |             |          |              |               |          |          |
| Haemophilus<br>influenzae type b                               |          | Hib      | Hib          |            | Hib       |              |           |         |         |         |             |          |              |               |          |          |
| Pneumococcal<br>disease                                        |          | VNC      | VNC          |            | VNC       |              |           |         |         |         |             |          |              |               | VNC      |          |
| Meningococcal<br>disease <sup>3</sup>                          |          | MenB     | MenB<br>MenC |            |           | MenB<br>MenC |           |         |         |         | Men<br>ACWY |          |              |               |          |          |
| Measles, Mumps<br>Rubella (MMR) <sup>4</sup>                   |          |          |              |            |           | SRP          |           | SRP     |         |         |             |          | SRP          |               |          |          |
| Varicella (chickenpox) <sup>5</sup>                            |          |          |              |            |           |              | VVZ       | VVZ     |         |         | VVZ         |          | VVZ          |               |          |          |
| Human<br>papillomavirus <sup>6</sup>                           |          |          |              |            |           |              |           |         |         |         | VPH         |          |              |               |          |          |
| Influenza <sup>7</sup>                                         | Flu      |          |              | Flu annual |           |              |           |         |         |         |             |          |              | Flu<br>annual |          |          |
| Herpes Zoster <sup>8</sup>                                     |          |          |              |            |           |              |           |         |         |         |             |          |              |               | HZ       | HZ       |

Systematic administration

Administration to vulnerable people or to those who have not been previously vaccinated

## Legend

### 1. Hepatitis B

#### Childhood vaccination:

Children of HBsAg-positive mothers and unvaccinated mothers who have not been screened should be vaccinated at 0, 2, 4 and 11 months of age. The first dose should be administered within the first 24 hours of life, together with anti-HB immunoglobulin.

#### Vaccination for adolescents and young adults:

For persons up to 18 years of age not previously vaccinated, three doses should be administered with a schedule of 0, 1 and 6 months.

### 2. Diphtheria, Tetanus, Pertussis

#### Prenatal (dTpa):

One dose of dTpa in each pregnancy from the 27th week of gestation, preferably in the 27th or 28th week.

#### Vaccination for adults (Td):

Check previous vaccination status before starting or completing a Td primo-vaccination regimen in adults. Contact health services, including occupational risk prevention services, to check vaccination status. If necessary, administer Td vaccination until 5 doses are complete. Administer a Td dose at around 65 years of age to persons who received 5 doses during childhood and adolescence.

#### Combined DTPa/IPV vaccine at 6 years:

Administer the combined DTPa/IPV vaccine to children vaccinated at 2, 4 and 11 months when they reach 6 years of age. Those who completed the vaccination schedule at 2, 4, 6 and 18 months (4 doses in total) should receive dTpa without IPV at 6 years of age.

### 3. Meningococcal disease (those born after 2023)

It is recommended to administer the vaccine against serogroup B meningococcal disease 1 or 2 weeks apart from other inactivated vaccines indicated at the same age to minimise the risk of reactogenicity with the concomitant administration. It is important to administer it on time at 2 and 4 months of age.

### 4. Measles, Mumps, Rubella (MMR)

The quadrivalent vaccine can be used for the 4-year-old dose to allow concomitant administration with varicella (chickenpox).

#### Vaccination for adults:

Contact health services, including occupational risk prevention services, to check vaccination status. Vaccination is recommended for persons with no documented vaccination history born in Spain after 1970. If necessary, 2 doses of TV may be administered at least 4 weeks apart. For those who have already had a dose, only administer one dose of TV. Vaccination is contraindicated for pregnant women and people with compromised immune systems.

## 5. Varicella (chickenpox)

The quadrivalent vaccine (TV) can be used for the 4-year-old dose to allow concomitant administration with measles, mumps, and rubella.

### Vaccination for adolescents:

For those who at 12 years of age have no history of having had the disease and have not been vaccinated, administer 2 doses of varicella (chickenpox) vaccine at least 4 weeks apart (preferably 8 weeks). If a single dose has been received previously, the schedule should be completed.

### Vaccination for adults:

For those with no evidence of chickenpox immunity in the adult population\*, perform a serological determination (IgG). If serology results are negative, administer 2 doses of varicella (chickenpox) vaccine at least 4 weeks apart (preferably 8 weeks). Having received 2 doses of vaccine is synonymous with immunity.

\* Evidence criteria for varicella (chickenpox) immunity in the adult population: vaccination documentation with two doses, history of varicella (chickenpox), history of herpes zoster or serological confirmation (IgG positive).

Vaccination is contraindicated for pregnant women and people with compromised immune systems.

## 6. Human papillomavirus

### Vaccination at 12 years of age (both sexes, born after 2011):

Two doses should be administered at least 6 months apart.

### Vaccination after 12 and up to 18 years:

Vaccinate only unvaccinated or partially-vaccinated women. If vaccination is started at 15 years of age or older, administer 3 doses with a schedule of 0, 1-2, 6 months (depending on the vaccine used). The schedule should be completed according to the age at the first dose.

## 7. Influenza

### Prenatal:

Pregnant women in any trimester during gestation should be vaccinated during the flu season.

### Childhood vaccination:

Vaccination with one dose during the annual campaign is recommended for those aged between 6 and 59 months.

### Vaccination for the elderly:

Single-dose vaccination during the annual campaign is recommended for those aged 60 years or over.

## 8. Herpes Zoster

Two doses should be administered 2 months apart.

## Abbreviations

**HB:** Hepatitis B vaccine

**DTPa:** Vaccine against diphtheria, tetanus and acellular pertussis with high antigen load.

**dTPa:** Vaccine against diphtheria, tetanus and acellular pertussis with low antigen load.

**Td:** Tetanus and diphtheria vaccination

**VPI:** Polio vaccine

**Hib:** Vaccine against Haemophilus influenzae type B disease

**VNC:** Conjugate vaccine against pneumococcal disease

**Men C:** Conjugate vaccine against serogroup C meningococcal disease

**Men B:** Conjugate vaccine against serogroup B meningococcal disease

**Men ACWY:** Conjugate vaccine against serogroups A, C, W and Y meningococcal disease.

**SRP:** Vaccination against measles, mumps and rubella (MMR).

**VVZ:** Chickenpox (varicella) vaccine

**VPH:** Human papillomavirus vaccine

**Flu:** Influenza vaccine

**HZ:** Herpes Zoster (shingles) vaccine