

**Math 33A**  
**Linear Algebra and Applications**  
**Discussion for February 14-18, 2022**

**Problem 1.**

The following determinant was introduced by Alexandre-Theophile Vandermonde. Consider distinct real numbers  $a_0, \dots, a_n$ , we define the  $(n+1) \times (n+1)$  matrix

$$A = \begin{bmatrix} 1 & 1 & \cdots & 1 \\ a_0 & a_1 & \cdots & a_n \\ a_0^2 & a_1^2 & \cdots & a_n^2 \\ \vdots & \vdots & & \vdots \\ a_0^n & a_1^n & \cdots & a_n^n \end{bmatrix}.$$

Vandermonde showed that  $\det(A) = \prod_{i>j} (a_i - a_j)$ , the product of all differences  $a_i - a_j$ , where  $i$  exceeds  $j$ .

- (a) Verify this formula in the case of  $n = 1$ .
- (b) Suppose the Vandermonde formula holds for  $n - 1$ . You are asked to demonstrate it for  $n$ . Consider the function

$$f(t) = \det \begin{bmatrix} 1 & 1 & \cdots & 1 & 1 \\ a_0 & a_1 & \cdots & a_{n-1} & t \\ a_0^2 & a_1^2 & \cdots & a_{n-1}^2 & t^2 \\ \vdots & \vdots & & \vdots & \vdots \\ a_0^n & a_1^n & \cdots & a_{n-1}^n & t^n \end{bmatrix}.$$

Explain why  $f(t)$  is a polynomial of  $n$ -th degree. Find the coefficient  $k$  of  $t^n$  using Vandermonde's formula for  $a_0, \dots, a_{n-1}$ . Explain why  $f(a_0) = f(a_1) = \cdots = f(a_{n-1}) = 0$ . Conclude that  $f(t) = k(t - a_0)(t - a_1) \cdots (t - a_{n-1})$  for the scalar  $k$  you found above. Substitute  $t = a_n$  to demonstrate Vandermonde's formula.

**Problem 2.**

Find

$$\det \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 & 5 \\ 1 & 4 & 9 & 16 & 25 \\ 1 & 8 & 27 & 64 & 125 \\ 1 & 16 & 81 & 256 & 625 \end{bmatrix}$$

using Vandermonde's formula and using the usual definition of determinant.

**Problem 3.**

For  $n$  distinct scalars  $a_1, \dots, a_n$ , find

$$\det \begin{bmatrix} a_1 & a_2 & \cdots & a_n \\ a_1^2 & a_2^2 & \cdots & a_n^2 \\ \vdots & \vdots & & \vdots \\ a_1^n & a_2^n & \cdots & a_n^n \end{bmatrix}.$$