

LPsP stimulation: lpsp>P2X2 (orange), empty>P2X2 (blue), lpsp - empty (black); mean ± SEM across flies, confirmed ejections only (walking: 951 of 5771 stims shown); ejections numbered within each trial, round 1 / round 2 = first / second trial of a same-side run
left: lpsp_p2x2_reredo (Figure 3 main set)
middle: lpsp_p2x2_walking post-7/8, 300 s trials (29 x 0.5 s pulses every 10 s; lpsp 15 / empty 11 flies; shared y axis; ^ = stim onset)
57 walking trials of 600 s / 59 pulses excluded: no dye response at any stim
reredo: EPG syt8s, 2 s ATP pulses
lpsp 13 flies / empty 12 flies
EPG green ΔF/F (ejection-side 16 glom.)
red (ATP dye) Σ ΔF/F (ejection side)
time from stim onset (s)
round 1: ej 1-5
n = 15/11
round 1: ej 6-10
n = 10/7
round 1: ej 11-15
n = 10/7
round 1: ej 16-20
n = 10/7
round 1: ej 21-25
n = 10/7
round 1: ej 26-29
n = 5/6
round 2: ej 1-5
n = 14/9
round 2: ej 6-10
n = 11/6
round 2: ej 11-15
n = 11/5
round 2: ej 16-20
n = 9/5
round 2: ej 21-25
n = 8/5
round 2: ej 26-29
n = 5/4
stim
round 2: next trial (~5 min later)
2 s
lpsp>P2X2: per-fly mean (mean 0-10 s), 15 flies
empty>P2X2: per-fly mean (mean 0-10 s), 11 flies
mean GCaMP ΔF/F (ejection side)
mean dye Σ ΔF/F (ejection side)
confirmed ejection # within trial (round 1 | round 2)
1-5 6-10 11-15 16-20 21-25 26-29 1-5 6-10 11-15 16-20 21-25 26-29