

LYMPHATIC FILARIASIS

A REVIEW 1862-2002



Rev'd Dr. Wayne Melrose

Lymphatic Filariasis Support Centre
School of Public Health & Tropical Medicine



First published in 2004

This book is copyright. Apart from fair dealing for the purposes of private study, research, criticism or review, as permitted under the Copyright Act, no part may be reproduced by any means without written permission. Enquiries should be made to Warwick Educational Publishing Inc.

© 2004 by Rev'd. Dr. Wayne Melrose

All rights reserved.

Publisher: Warwick Educational Publishing Inc

Enquiries: Warwick Educational Publishing Inc
Secretariat, 29 Kurrajong St., Killarney Qld 4373, Australia
Tel: +61-7-47225760
Email: secretary@wepi.org

National Library of Australia

Author: Melrose, Wayne D.

Title: Lymphatic Filariasis: A Review 1862-2002.

ISBN 0-9579788-4-7

Date of Publication: 1st Mar 2004 Price: free

Format: WEB Size: 297x210 No. of Pages: 80

Disclaimer: While every reasonable care has been taken to ensure that the information in this text is correct, neither the Editor, contributors, nor Warwick Educational Publishing Inc can accept any responsibility for the inclusion or omission of any material or for any inaccuracies in the information supplied.

Editor: Deon Canyon

Cover design: Deon Canyon

Cover photos: Jamie Carron (Island Beach) & Deon Canyon (Blood Collection)

Contents

<i>Foreword</i>	3
<i>Introduction</i>	4
<i>History of Filariasis</i>	7
<i>The Vectors of Filariasis</i>	9
<i>Transmission Dynamics in Filariasis</i>	10
<i>The Global Burden of Lymphatic Filariasis</i>	12
<i>Traditional and Cultural Beliefs</i>	15
<i>Socio-Economic Impact of Filariasis</i>	18
<i>The Spectrum of Filarial Disease</i>	20
<i>Filariasis and Other Illnesses</i>	25
<i>Filariasis in Expatriates</i>	28
<i>Filarial immunity</i>	33
<i>Unravelling the Pathogenesis of Chronic Pathology</i>	38
<i>More than One Pathway to Chronic Pathology?</i>	39
<i>Diagnosis of Lymphatic Filariasis</i>	41
<i>Control of Lymphatic Filariasis</i>	46
<i>Adverse Reactions to Anti-Filarial Drugs</i>	51
<i>Effect of Therapy on Microfilaraemia and Immunological Responses</i>	52
<i>Conclusion</i>	53
<i>References</i>	54

Foreword

This review on lymphatic filariasis by Dr Wayne Melrose has come at an opportune time. The Global Program for the Elimination of Lymphatic Filariasis (GPELF) was launched by the World Health Organization in 1997 as a public/private sector partnership organized as a “Global Alliance”, the object being to eliminate lymphatic filariasis as a public health problem by the year 2020. The program is in its 5th year of operation and recent advances in the diagnosis of infection, prevention of transmission and treatment of infection has generated considerable expectations of the success of the control program among endemic countries. A good review of any disease should give the reader a comprehensive understanding of the “state of the art” in the subject and discuss areas that need further study and research. Dr Wayne Melrose has accomplished this admirably well as he takes us through the history, biology, epidemiology, immunology, disease pathogenesis, socio-economic impact, diagnosis, control and treatment of this important tropical disease. His wide field experience, particularly in Papua New Guinea, is reflected throughout the book and it brings with it the reality of the situation and the challenges faced when one is dealing with such a complex disease. This review should be compulsory reading for all researchers, students and tropical medicine practitioners and others who are keen to learn more about lymphatic filariasis.

Professor Dato

Dr C.P. Ramachandran

Kuala Lumpur

Introduction

Lymphatic filariasis is a mosquito-borne tropical disease, which afflicts around 120 million people worldwide. The World Health Organization (WHO) ranks it as the second most common cause of long-term disability. In 1997, the WHO identified filariasis as one of six potentially eradicable infectious diseases and initiated a global plan for its elimination as a public health problem by the year 2020 (WHO, 1999).

The purpose of this review is three fold: To provide a comprehensive source document which can be used to write educational and health promotional material on lymphatic filariasis; to provide a background for my own original research and to highlight areas which require further investigation in order to stimulate others to carry out further research on this fascinating disease. Extensive use has been made of the reviews by: Sasa, 1976; Ottesen 1980, 1989, 1993, 1994; Piessens and Partono, 1980; Dasgupta, 1984; WHO, 1984, 1987, 1992; 1993, 1994, 1995, 1998, 1999; Partono, 1987; Nanduri and Kazura, 1989; Grove, 1990; Evans *et al.*, 1993; Kazura *et al.*, 1993a; Turner, 1993; McMahon and Simonsen, 1996; Roberts and Janovy, 1996; Ramachandran, 1997; Addiss, 1998;


Freedman, 1998; de Almeida and Freedman, 1999; Michael, 1999.

The filariasis literature is voluminous and this review does not purport to cover all available material. An attempt has been made to cover all the most important aspects of filarial parasites and disease, and to include some of the more obscure and often forgotten material, some of which needs to be revisited in the light of modern filariology. Other excellent reviews on the history of filariasis have been written by Manson-Bahr (1959), Grove (1990) and Routh and Bhowmik (1993).

An Overview of Filarial Parasites

Filarial parasites (filaroids) are long hair-like tissue-dwelling nematodes. All except the Guinea worm *Dracunculus medinensis* (which uses a copepod) employ arthropods as intermediate hosts. All filariids have a similar life cycle that includes an obligatory maturation stage in a blood-sucking insect or copepod, and a reproductive stage in the tissues or blood of a definitive host (Figure 1).

Figure 1. Typical filaroid life cycle



(Source: http://www.dpd.cdc.gov/dpdx/HTML/ImageLibrary/Filariasis_il.asp?body=A-F/Filariasis/body_Filariasis_il_th.htm with permission from CDC)

Adult male and female worms live in the lymphatics, skin, or other tissues (Figure 2). Microfilariae (which are specialised embryos not larvae) are produced by the female worm, circulate in the blood or invade the skin, and are ingested by the vector (Figure 3). Larval development but not multiplication occurs within the muscles of the vector. The infective L3 stage migrates to the proboscis and is transmitted to the new host during feeding (McMahon and Simonsen, 1996). Unlike malaria, the infective stage is not directly injected into the skin of the new host. It is deposited onto the skin whilst the mosquito is feeding and finds its own way through the skin, usually via the puncture made by the mosquito.

Filaroids have been found in all classes of vertebrates except fish and are especially common in birds (Roberts and Janovy 1996). All but one of the species that cause human disease (the Guinea worm) belong to the family Onchocercidae (Roberts and Janovy 1996).

Figure 2. Adult worm in lymphatic vessel. Haematoxylin – eosin stain X400.




Figure 3. Microfilaria in blood. Giemsa stain X200.



Human diseases caused by filarial parasites

Lymphatic Filariasis

Wuchereria bancrofti, *Brugia malayi* and *Brugia timori* are mosquito-borne and cause lymphatic filariasis. In many people's minds lymphatic filariasis is synonymous with "elephantiasis" characterised by

the enlargement of scrotum or limbs (Figure 4). In most endemic areas elephantiasis only occurs in a small proportion of the people suffering from lymphatic filariasis.

Bancroftian filariasis, caused by *W. bancrofti* is responsible for 90% of lymphatic filariasis cases and is found throughout the tropics and in some sub-tropical areas. *Brugia malayi* is confined to Southeast and Eastern Asia. *Brugia timori* is found only in Timor and its adjacent islands (McMahon and Simonsen 1996; Roberts and Janovy 1996). In this review "filariasis" always means lymphatic filariasis unless otherwise specified.

Figure 4. Patient with advanced elephantiasis from Papua New Guinea



Onchocerciasis

Onchocerciasis (also called river blindness) is caused by *Oncocerca volvulus* and the vector is the *Simulid* black fly. The largest numbers of cases occur in Africa and it was spread to Central America, Venezuela and Colombia via the slave trade. There is also a small focus on the Arabian Peninsula. Microfilariae invade the skin and give rise to dermatitis premature aging of the skin and skin atrophy. Development of the adult worm leads to nodule formation. Microfilariae invade the eye and cause an inflammatory reaction that can lead to blindness (McMahon and Simonsen 1996; Roberts and Janovy 1996).

Loiasis

Loiasis is caused by *Loa loa* (sometimes called the African eye worm) and is spread by *Chrysops* flies. It is present in Central and West Africa. Signs and symptoms include fugitive or "Calabar" swellings, itching and joint pains. Sometimes an asymptomatic invasion of the eye surface occurs hence the name "eye worm." (Negesse *et al* 1985; McMahon and Simonsen, 1996; Roberts and Janovy, 1996). It is the most common filaroid infecting travellers from non-endemic

areas (Azimi and Grossman, 1995).

Miscellaneous Filaria Infections

Mansonella perstans (Africa and Central and South America), *Mansonella streptocerca* (Africa) and *Mansonella ozzardi* (Central and South America) are spread by *Culicoid* midges. Most infections are asymptomatic but loiasis-like signs and symptoms can occur (McMahon and Simonsen, 1996).

Zoonotic Filarial Infection

Human infections with various known and unknown animal filarial parasites have regularly been reported throughout the world but especially from the United States (Tobie and Beye, 1962; Beaver and Orihel, 1965; Nelson, 1965; Orihel and Eberhard, 1998). *Dirofilaria immitis* is the cause of heartworm in dogs and occasionally effects humans (for a review see Rodrigues-Silva *et al.*, 1995). Most cases are asymptomatic and the worm becomes calcified in the lung resulting in a "coin lesion" which may be mistaken for carcinoma or tuberculosis. Occasionally asthmatic-like symptoms occur and there may be a marked eosinophilia. Another dog filaroid - *Dirofilaria repens* can cause subcutaneous nodules, peri-orbital lesions, coin lesions in the lungs, and breast lumps in humans and rare cases have been reported throughout the world (Bennett *et al.*, 1989; Pampiglione *et al.*, 1995. High levels of antifilarial IgG, IgE and IgM antibodies are present in patients with dirofilariasis (Simon *et al* 1997) but it should be borne in mind that there is a high background level of dirofilaria-associated antibody in communities in close contact with infected dogs and the prevalence of *D. immitis* antibodies in humans is closely related to the number of infected dogs in the community (Welch and Dobson, 1974). At least six cases of human filarial dermatitis have been attributed to *B. beaveri* (Rosenblatt *et al.*, 1962; Grove, 1990), a filarial species whose natural host is the Raccoon (Ash and Little, 1971) and nine cases of lymph node infection by a unidentified *Brugia* species (Baird *et al* 1986) possibly

from the domestic cat (Beaver and Wong, 1988). Animal filaroids seldom seen to develop and become sexually mature in man and produce a microfilaraemia but it does occur from time to time (Orihel and Eberhard, 1998). Greene *et al.*, (1978) found circulating microfilariae in a patient from Alabama who was suffering from Lupus erythematosus. Although the species could not be identified with certainty, the most likely species appeared to be *Mansonella interstitium*, a filaroid of squirrels. There is also a report of microfilaraemia and eosinophilia in a 70-year-old Greek man who had never been in an area endemic for human filarial parasites and was presumably infected with an animal species (Petrocheilou *et al.*, 1998).

Cases of encephalopathy due to infection with monkey filarial parasite *Meningonema peruzzi* have been reported from Zimbabwe and the Chimpanzee parasite *Microfilaria rodhaini* has been found in human skin biopsies. Two other *Microfilaria* species believed to be of animal origin have been found in human blood, *M. bolivarensis* in American Indians in Venezuela and *M. semi-clarum* in villagers in North-western Zaire (McMahon and Simonsen, 1996).

Dracunculiasis

Dracunculus medinensis is a large filarial worm that causes Dracunculiasis or Guinea worm disease. Its vector is a fresh water copepod and its life cycle differs from that of a typical Filaroids. Instead of invading the tissues the microfilariae are liberated directly from the uterus of the female worm into water.

The female worm migrates to the lower leg or foot and an ulcer is formed from which the microfilariae are discharged. These ulcers often become infected with bacteria resulting in cellulitis or if in the vicinity of a joint, arthralgia. The worm causes ulcers on the lower leg or foot, which often become infected with bacteria.

Chronic inflammation of the joints can lead to stiffness and permanent disability (McMahon and Simonsen, 1996).

History of Filariasis

Filariasis in Antiquity

Not all cases of elephantiasis are caused by filariasis and elephantiasis is not the most common symptom of lymphatic filariasis, it is however the most visible and has been known since early times. Elephantiasis features in the mythology of several societies (Neisius, 1927; Routh and Bhowmik, 1993). Tales of "skiapodes" mythical beings who could "shade themselves from the sun by the greatness of their leg" occur in the mythology of Greece, Rome and India (Price, 1989; Lawence, 1990). Indian mythology also contains the legend of the "curse of St. Thomas" which states that swollen legs occur in the descendants of those who martyred him (Bowers and Carruba, 1970; Lawence, 1970). Elephantiasis is also depicted in art from ancient times. A statue of Pharaoh Mentuhotep 111 (2000 BC) and the "Queen of Punt" stela (1500 BC) graphically depict the deformities associated with elephantiasis of the limbs (Lawence, 1967; Hoeppli, 1969, 1971; Schacher and Sahyuni, 1967) and scrotal swelling is depicted in art from Nigeria (500-200 BC) and South America (500 AD) (Andrews, 1959; Hoeppli, 1969, 1971).

Mythology and art aside, the disease was well known to physicians and medical writers from very early times. Ancient Hindu medical workers knew of the condition and it is referred to in Sanskrit texts dating back to 600 BC. Included in these is a comment that "the disease is prevalent in areas where stagnant water is found" - a very relevant comment given the link between *Culex* mosquitoes, which like to breed in stagnant water and filariasis transmission (Lawence, 1970; Zimmer, 1979). Between 600 and 250 BC men suffering from elephantiasis were prohibited from becoming Buddhist priests (Lawence 1967). The Roman writer Lucretius Carus (100BC) regarded elephantiasis as a characteristic disease of the Egyptians, which was facilitated by the climate around the Nile (Hoeppli, 1969). Persian physicians of the 10th and 13th centuries AD and medieval European physicians provided very accurate descriptions of elephantiasis and hydroceles (Lawence 1967) but it was not until the 19th century that a firm link was established between elephantiasis, hydroceles and chyluria (Routh and Bhowmik, 1993).

The Discoveries

Wuchereria bancrofti

In 1862 Jean-Nicholas Demarquay, a Cuban surgeon, discovered a worm-like "creature" in hydrocele fluid (Demarquay, 1863). In 1866 Brazilian, Otto Wucherer, found the same organism in chylous urine (Wucherer, 1868) but it was not until 1872 that microfilariae were discovered in blood (Lewis, 1872).

In 1876, Brisbane physician and parasitologist, Joseph Bancroft, discovered a female adult worm in an abscess on the arm of a butcher (Cobbold, 1877) and further examples were found in South America (da Silva Araujo, 1877; dos Santos, 1877) and China (Manson, 1881a,b). In 1888, Sibthorpe and Bourne discovered the adult male (Bourne, 1888; Sibthorpe, 1889). The name *Wuchereria bancrofti* was formally adopted in 1921 (Seurat, 1921).

Brugia Species

In 1927 Lichtenstein and Brug discovered a microfilaria in the Dutch East Indies (now Indonesia) which was morphologically different from *W. bancrofti* and called it *Filaria malayi* (Brug, 1927; Lichtenstein, 1927; Brug, 1928; Brug and de Rook, 1930).

This in fact may not have been the first description of *Brugia*. In 1905 Ashburn and Craig described a case from the Philippines, which had microfilariae that differed from *W. bancrofti*, which they called *Filaria philippinesis* (Ashburn and Craig, 1905). Their material was reviewed by Manson-Bahr in 1941, who believed that the microfilariae were identical to Ashburn and Craig's *F. malayi*. Rao and Maplestone first described the adult worm in 1940 (Rao and Maplestone, 1940). The pioneering work of Brug was acknowledged in 1958 when Buckley proposed the new genus *Brugia* and *F. malayi* was re-named *Brugia malayi* (Buckley, 1958; 1960). The zoonotic feature of *Brugia* was discovered in 1939 when microfilariae, later identified as those of *B. malayi* were discovered in a Kra monkey (Poynton and Hodgkin, 1939). Edeson and Wharton, (1958) and Laing *et al.*, (1960) broadened the range of vertebrate host to include other monkey species, domestic cats and dogs.

In the 1960's another filarial species was discovered in Portuguese Timor and given the name *Microfilaria timori* (David and Edeson, 1965). Partono infected Mongolian gerbils with the new species and obtained adult worms. He confirmed that the new species belonged to the genus *Brugia* and called it *B. timori* (Partono *et al.*, 1977).

Mosquito Hosts

On August the 10th, 1887 Patrick Manson fed some mosquitoes on the blood of his microfilaraemic gardener Hin-Lo and was able to demonstrate the development of the larva within the insect (Manson, 1878a, 1878b, 1884). Lewis confirmed his observations in 1872. The mosquitos used by Manson in his initial experiment were later found to *Culex quiquefasciatis* (Manson-Bahr, 1927), which is still regarded as one of the most important vectors of filariasis. By 1976 other species of *Culex* and species of *Anopheles*, *Aedes* and *Mansonia* had been added and the list had grown to several hundred (Sasa, 1976;

Grove, 1990).

Filarial Nocturnal Periodicity

Patrick Manson had two laboratory assistants. One worked during the day and one during the night. Manson noticed that the night worker found more filaria than his day-working colleague and this led him to set up an experiment whereby serial blood samples were taken on the same patients during the day and night. He found that the microfilariae were present in large numbers during the night but were almost absent during the day (Manson, 1880a, 1880b). Manson's observation was greeted with some disbelief, someone even asking, "whether filaria carried watches" (Grove, 1990). Myers, (1881) and MacKenzie, (1881) subsequently silenced the scoffers by confirming Manson's findings. Mackenzie also successfully

reversed the nocturnal periodicity by getting his patient to sleep by day and remain awake at night (MacKenzie, 1881) as did Manson the following year (Manson, 1883) and changes in periodicity have been obtained with as little as a three day reversal in sleeping patterns (Hawking, 1962a). There were a large number of early theories about the mechanism of micro filarial periodicity and they have been given an extensive review by Lane (1948). In 1883 Manson discovered that *D. immitis* microfilariae in a dog hid themselves in the lung when they were absent from the circulation (Manson, 1883; Grove, 1990). Manson confirmed this in a human case in 1897 (Manson, 1899). Thorpe (1896) and Lynch (1905) reported that the microfilariae (subsequently called *W. bancrofti* var *pacific*) showed no periodicity. Turner and Edeson (1957) demonstrated that there are both nocturnal and sub-periodic forms of *Brugia malayi*.

The Vectors of Filariasis

Unlike malaria whose transmission cycle is dependant on *Anopheles* species, filariasis a wide range of mosquitoes act as vectors for filariasis (Manson-Bahr, 1959; WHO, 1992). By far the most widespread and important species is *Culex quinquefasciatus*. The ability of this mosquito to breed in heavily polluted water in urban and semi-urban areas he resulted in a prevalence of "urban" filariasis in areas where the rapid growth of urban areas outstrips the growth of sanitation services (figure 5).



It is a night-biting mosquito and filariasis in the areas where it is the major vector shows nocturnal periodicity. Other *Culex* species are also capable of transmitting filariasis and may be important vectors in certain areas. *Aedes polynesiensis* and *Ae. samoanus* are the most important vectors in the pacific area. The former breeds in crab holes and tree holes making it a very difficult mosquito to control with conventional methods (Figure 6).



They are day-biting mosquitoes thus accounting for the diurnal periodicity of filariasis in this region. *Ae. poecilius*, a night-biting mosquito is a major vector in the Philippines. *Mansonia* species are an important vector of *B. malayi*, and sometimes *W. bancrofti*, in areas where there are extensive areas of aquatic plants

(figure 7).



Anopheles species are important vectors of *W. bancrofti* in parts of Africa and Southern Asia and in Papua New Guinea. *A. barbirostris* breeds in open rice paddies and is the only known vector of *B. timori* (figure 8).



Are any other insects except mosquitoes capable of transmitting lymphatic filariasis? Bed bugs (*Cimex* species) can be naturally and experimentally infected with *Brugia malayi* and *W. bancrofti* (Burton, 1962) and some limited larval development can take place but the mortality of the larvae is extremely high and bed bugs cannot be seriously regarded as a vector of filariasis (Burton, 1963).

Transmission Dynamics in Filariasis

The host-parasite relationship and the transmission dynamics of filariasis have been extensively reviewed by Sasa (1976), Piessens and Partono (1980) and Anderson and May (1991). Two modelling frameworks *lymfasim* (Plaisier *et al.*, 1998) and *Epifil* (Chan *et al.*, 1998) have been developed which also a series of hypotheses can be tested about the life history of the parasite, its transmission, the immuno-regulatory role of the host immune system, the development of disease and the impact of control measures.

In general, the mosquito infection rate, ie the number of mosquitoes that contain microfilariae after a blood meal and the number of microfilariae ingested per mosquito increase with increasing parasitaemia (Piessens and Partono, 1980). Although there is no multiplication in the vector the number of infective larvae is around six times greater than that expected by microfilariae density (Brito *et al.*, 1998). Bockarie (1997) has shown that the annual infective biting rate and the annual transmission potential show a positive correlation with microfilariae rate, microfilariae density and prevalence of leg oedema suggesting that transmission intensity is a major determinant of patent infect and morbidity. Kazura *et al.*, (1997) studied the relationship between annual transmission potential (ATP) and disease status in 1666 individuals from five similar but distinct *W. bancrofti*-infected communities from a highly endemic area of Papua New Guinea. Annual transmission potentials and microfilariae prevalence (MF) in these villages were: 2,344 (MF 94%); 1,338 (MF 82%); 279 (MF 56%); 179 (MF 66%) and 31 (MF 52%) respectively. In all villages the prevalence of leg odema was highly positively related to the ATP (correlation coefficient $r=0.89$ and probability $p=0.04$). The incidence of acute filarial attacks is also related to the transmission intensity. Gyapong *et al.*, (1996) found that the incidence of acute filarial attacks we reduced in the dry season when the transmission potential was at its lowest.

The concept of “facilitation” - that the proportion of microfilariae that develop increases as the number of microfilariae ingested increases (WHO, 1992) is an important consideration when the possibility of eradication of filariasis is being considered. In theory, it should be possible to reach a point where there are insufficient circulating microfilariae in the population to support transmission (WHO, 1992). It is important to note however, that persons with microfilariae densities as low as three per ml can still infect mosquitoes and that residual low-density microfilaraemics after mass treatment programs have the potential to cause rapid resurgence of filariasis (Lowrie *et al.*, 1989; Southgate, 1992). Beckett (1973) found that a heavy uptake of microfilariae from the blood can cause life-threatening damage to the internal organs of the mosquito during larval development but this finding is not supported by the recent work of Brito *et al.*, (1998). The feeding of mosquitoes on

individuals with medium or low micro filarial densities may therefore enhance transmission potential especially if the main vector is a Culicine rather than an Anopheline (Webber, 1981, 1991). The reasons for this difference, termed “limitation” is discussed by Bryan and Southgate (1988a,b) Bryan *et al.*, 1990; Webber (1981, 1991); Southgate and Bryan (1992) who suggest, that in contrast to Culicines, Anopheline mosquitoes have a well developed pharyngeal armature which damages microfilariae when they are ingested. If the number of ingested microfilariae is low there may be insufficient viable microfilariae to infect the mosquito. Loss of microfilariae also occurs in fluid expelled from the anus of anophelines but not culicines (Bryan and Southgate, 1988a,b). The concepts of facilitation and limitation and their role in filariasis transmission has been critically reviewed by Wada *et al.*, (1995) who concluded that “there was no clear evidence to support the existence of facilitation and limitation-based unstable equilibrium in relation to microfilariae prevalence and density below which filariasis would spontaneously disappear even when the vector was *Anopheles*. Instead, the existence of a critical level of man/mosquito contacts for the disappearance of filariasis was suggested”.

Factors that Influence Micro Filarial Prevalence and Density

A spontaneous decrease in microfilaraemia prevalence can occur in the absence of vector control or mass chemotherapy. In a study in Benin the prevalence of microfilaraemia decreased from 9.4% to 0.48% over a 10-year period. The prevalence of people with chronic pathology remained the same. The changes could not be explained by environmental or sociological changes in the region or even by changes in the demographics of the study population (Myung *et al.*, 1998). Microfilariae density is also subject to considerable daily variation (Rouchou, 1954; Nathan *et al.*, 1980) but this is unrelated to lunar phase lactation, or the menstrual cycle (Rachou, 1956; Nathan *et al.*, 1982). Fasting during the month of Ramadan has been shown to reverse nocturnal periodicity (El-Tamami, 1960; Nathan *et al.*, 1982) but solar eclipses have no effect on periodicity (Sahai *et al.*, 1980). The fecund life span of *W. bancrofti* has been calculated by various means and methods and has been variously estimated from 5 to 15 years. A recent study by Vanamail *et al.*, (1996) suggests that the life span is at the lower end of previous estimates - around five years.

Endosymbiont Bacteria

In the 1970's intracellular bacteria were discovered in some species of female filarial parasites (reviewed by Taylor and Hoerauf, 1999; Taylor *et al.*, 2000). It was postulated that they were of the genus *Wolbachia* (now

known to be correct) and had a similar role to the symbiont bacteria found in arthropods where they are known to exert an influence upon the growth and reproduction of their host. Symbiont bacteria may be

implicated in the pathogenesis of filarial pathology and are a potential target for antifilarial activity (see relevant sections below).



The Global Burden of Lymphatic Filariasis

The total global burden of lymphatic filariasis is not known and mapping of its prevalence is on-going (for review see Michael and Bundy, 1997). The disease is endemic in 76 countries and 1.1 billion people are at risk from infection (Ramachandran, 1997). Statistics for many countries are either old, very sketchy, or non-existent. The most recent estimates (Michael and Bundy, 1997; Michael *et al.*, 1996; Ramachandran, 1997) suggest that filariasis infects 120 million people or 2.0% of the world's population. Of these, 83 million people have lymphatic disability, 23 million men have hydroceles and 15 million people have elephantiasis. Millions more suffer from debilitating acute attacks of filarial fevers and lymphadenitis (Ramachandran, 1997). Over 90% of these cases are caused by *W. bancrofti*.

The estimated country-by-country prevalences in the Asia-Pacific region are given in Table 1.

The true prevalences are probably higher than this because most country surveys done in the past were based on the demonstration of microfilariae. Modern surveys using antigen detection (see "Diagnosis" section below) have shown that surveys of microfilaraemia alone can under-estimate the prevalence of filariasis by as much as 30% (Turner, 1993; Chanteau *et al.*, 1995).

Increasing Global Prevalence

There is evidence that the global burden of filariasis is increasing. To quote Piessens and Partono (1980):

"Filariasis shares with schistosomiasis the dubious honour of being one of the major diseases whose prevalence and distribution are increasing in the developing world at a time when other tropical diseases are slowly disappearing. This is all the more remarkable in view of the relative inefficiency of the filarial transmission, the complexity of the parasite's life cycle and the susceptibility of the worms to the widely available and inexpensive drug diethylcarbamazine, that should render filarial nematodes particularly vulnerable to effectively applied control measures."

Table 1. Prevalence of filariasis in various countries (Hawking and Denham, 1976; Michael *et al.*, 1996).

COUNTRY	PREVALENCE (%)
China	0.5
Burma	0.1
French Polynesia	12.6
Indonesia	1.8
India	5.3
Laos	0.1
Malaysia	0.4
Nepal	0.8
New Caledonia	18.7
Micronesia	11.1
Papua New Guinea	72.0
Philippines	4.1
Western Samoa	7.7
Seychelles	10.0
Sri Lanka	1.6
Thailand	2.7
Vanuatu	1.9
Vietnam	1.3

There are several reasons for this. The main vector mosquitoes thrive in urban slum areas and the increased urbanisation around the world's major tropical cities are putting an increasing number of people at risk (Hunter, 1992; Albuquerque *et al.*, 1995; Dhanda *et al.*, 1996). It has long been recognised that one of the adverse effects of economic development projects is the impact on health. Irrigation projects in Ghana have been associated with an increase in filariasis. Hunter (1992). Found that almost every house in villages bordering a man-made irrigation scheme had a case of filariasis. By contrast, only scattered cases were seen in villages two km or more away from the canals. Dzodzomenyo *et al.*, (1999) observed a similar result.

Rural-urban migration and uncontrolled urbanisation often lead to overburdening of sewerage and wastewater systems. The resulting pools of stagnant, polluted water provide an ideal breeding ground for *Culex quinquefasciatus*, a major vector of filariasis (Mak, 1986). In contrast to anophelines culicines can efficiently transmit filariasis in situations where the microfilariae density is low (Webber, 1991). Adhikari *et al.*, (1994) found a significant difference in filariasis prevalence between colliery areas, where there were opencast pits and other vector breeding sites, and non-colliery areas of the same district in India. A similar study in Egypt (Gad *et al.*, 1994) found that inhabitants of houses facing vacant land (which presumably had more opportunity for water to collect and breed mosquitoes) suffered more from filariasis than their

neighbours from the more closely settled areas nearby. This study demonstrates an important aspect of filariasis - that it is very heterogeneous and very focal in its geographical distribution and its prevalence can vary widely even within a small area.

Pursuit of economic activity can also lead to increased transmission in other ways. Rubber trees are often tapped at dawn in an attempt to increase latex yield. This corresponds with a peak in the biting rate of *Mansonia* mosquitoes (Mak, 1986). In many tropical communities and urban areas people congregate for social activities in the early evening - a prime biting time for many filarial vectors (Mak, 1986). It should also be pointed out that malaria is also on the increase, one of the causes being the winding down of vector control program in many parts of the world and increased mosquito resistance to insecticides. Since malaria and filariasis share some common vectors, failure to control mosquitos also contributes to the increase in filariasis. Population movements are playing an increasing part in the spread of filariasis. Filariasis is generally thought to be under control in Thailand but it is feared that there will be resurgence in the disease caused by infected Myanmar migrants (Trierapapab *et al.*, 1997). To quote Ramachandran (1997):

"Statistics become reality when one takes a walk through the villages and shanty towns in endemic countries and visualise people, both young and old, men women and children walking around with enlarged legs, feet, hands and genitals, in addition to other parts of their bodies. Today the disease still continues to spread unabated in many parts of the world due to continued unplanned urban and semi-urban development; man-made environmental and ecological disasters and habitats which continue to enhance mosquito breeding and consequent transmission of disease."

Gender Differences

Animal studies have suggested that estrogenic hormones may play a part in reducing filarial infection in females (Reynouard *et al.*, 1984). Several studies have shown that the microfilaraemia density is lower in women of reproductive age compared with age-matched males. This difference cannot be explained by differences in exposure (say by different clothing) is not associated with pregnancy and may be related to hormonal activity (Dutta and Diesfield, 1994; Kaur, 1997; Alexander and Grenfell, 1999b). Mavoungou *et al.*, (1989) suggest a different hypothesis: that the filarial parasites themselves may contribute to reduced steroid and gonadotropin hormone levels in young women and may lead to a delayed menarche.

The Age-Distribution of Filariasis

The age distribution of microfilaraemia is remarkably consistent throughout the parasite's geographic range. Both prevalence and intensity tend to rise with age and peak in the 15 to 25 year age group and decline in adult hood (Sasa, 1976; Day *et al.*, 1991; Chanteau *et al.*, 1995). In a Haitian study, Lammie *et al.*, (1994) found that filarial antigenaemia increased from 24.5% in children aged 1 to 5 years to 70% in adults older than 50 years. More males are infected than females and the incidence of disease is higher in males (Brabin, 1990; Albuquerque *et al.*, 1995; Michael *et al.*, 1996; Kaur, 1997). The prevalence of antigenaemia and microfilaraemia and the density of microfilaraemia are usually closely related to transmission intensity (Day *et al.*, 1991; Chanteau *et al.*, 1995). Transmission intensity is determined by complex interactions between the species of parasite, the microfilariae rate and density and the vector (Southgate, 1984; Bryan and Southgate, 1988a,b; Bryan *et al.*, 1990; Southgate 1992; Southgate and Bryan, 1992). It has also been suggested that the higher microfilariae prevalence in younger individuals is related to the increased vector-biting rate in younger people (Farid *et al.*, 1997).

Classification of Filarial Endemicity

There have several attempts at the classification of filarial endemicity. Most of these used, as well as parasitological data, subjective clinical findings that are difficult to standardise (Iyengar, 1938; Kumar, 1996). It has been shown that the filarial infection rate as evidenced by microfilaraemia closely correlates with other filariometric indices and can be use to set cut off points for various levels of endemicity (Kumar, 1996).

His proposed classification is given in Table 2.

Table 2. Proposed classification of filarial endemicity (Kumar, 1996)

INFECTION RATE	ENDEMICITY
<10%	Low
10-20%	High
20-40%	Hyperendemic
>40%	Holoendemic

Given that microfilaraemia surveys alone underestimates the prevalence of filariasis by as much as 30% (see above) it would be interesting to do a similar study using filarial antigenaemia as the indicator of infection.

CAN FILARIASIS BE TRANSMITTED OUTSIDE ENDEMIC AREAS?

In theory, yes.

Many mosquito species can transmit the parasite the presence of a microfilaraemia person could result in local transmission or a case of "airport filariasis" could occur if an infected mosquito accidentally transported into the country. Rosenblatt *et al.*, (1962) describe a case, which they believe could have been transmitted

within the city of New York, possibly because of a visit by the patient to the city's international airport. The species of filaria however could not be identified with certainty and it is possible that it was an animal species rather than a human *Brugia* or *Wuchereria bancrofti*.

Traditional and Cultural Beliefs

Social Attitudes Towards Filariasis

An understanding of the community social attitudes to filariasis (or any other disease) is essential if control is to be achieved. To quote Gyapong (1996):

"Disease control programs in developing countries are often unsuccessful or unsustainable because strategies pursued are inappropriate for the community or incompatible with traditional perceptions of aetiology, prevention and control. Feasible interventions to treat and prevent filariasis will require a broad understanding of traditional perceptions of the disease, its cause and consequences and means of prevention. Since perceptions of disease vary from place to place, there is a need to carry out in-depth studies of the social, cultural and economic aspects of the disease before embarking on control measures".

Although Ramachandran (1997) states that: "It cripples them, demoralises them and makes them social outcasts in their own communities and societies due to social stigma and alienation", the social impact of filariasis varies widely from place to place. In general the degree of stigma seems to be associated with the severity and visibility of the disease (Evans *et al.*, 1993; Mujinja *et al.*, 1997). In the Philippines, men with hydroceles continued to lead normal lives although they are subjected to some teasing. On the other hand, women with labial enlargement were presumed to be promiscuous and the shame associated with the disease lead to poor reporting (Evans *et al.*, 1993). In Thailand filariasis is considered to be a "terrible disease" and some sufferers with elephantiasis are shunned by their family and fellow villagers. People refuse to sit with them, eat with them or marry them. The burden is especially heavy for young women who believe that their marriage prospects are nil and they become reclusive and seldom leave their house (Rauyajin *et al.*, 1995). In Tanzania, having filariasis is considered shameful (Muhondwa, 1983; Evans *et al.*, 1993; Mujinja *et al.*, 1997) and in Polynesia during the 1950's men with elephantiasis of the scrotum were considered to have a social and procreative handicap and women with elephantiasis were not considered desirable as wives (Kessel, 1957; Evans *et al.*, 1993). In Ghana people with mild or moderate elephantiasis are well accepted by the community but those with more advanced disease often stay at home because they feel embarrassed about their disease. Men with hydrocele are often teased behind their backs and those with chronic

filariasis cannot be chosen as chiefs. Young women with filariasis usually have problem getting married because prospective husbands consider that the disease will make them unable to work and he will have the cost of paying for her treatment (Hunter, 1992; Gyapong *et al.*, 1996a). In Nigeria, filarial disease seriously hinders a girl's prospect of marriage, or if she is married the stability of the marriage and her prospect for future happiness. The unwillingness of men to marry girls with the disease is compounded by the belief that what happens to the mother will also happen to the child (Amazigo and Obikeze, 1992; WHO, 1995). Women had similar sentiments about marrying men with filariasis but had less of a choice in the matter than men because relatives arranged many of the marriages. Mujinja *et al.*, (1997) in a study in Tanzania, found that stigma, ostracism and discrimination of the filaria-affected, though subtle, have a significant psychological impact. Hunter (1992) records that the marriage situation differs between communities that are filarial-hyperendemic and those where the prevalence of disease is low. Men from the communities where prevalence is low avoid taking girls from the hyperendemic areas as wives. By contrast, there seems to be much less stigma attached to the disease in hyperendemic communities and men will even marry girls who are already showing signs of filariasis. Coreil *et al.*, (1998) however, showed that although young men will take filarial-affected girls as wives, her chances of finding a good husband and marital happiness are diminished. In coastal Ghana sufferers are subjected to teasing and considered to be unsuitable marriage partners. Those who do marry have a higher than normal divorce rate (Ahorlu *et al.*, 1999).

Because the social stigma associated with filariasis is particularly burdensome for women, filariasis has been identified along with schistosomiasis, leishmaniasis, leprosy and onchocerciasis, as diseases which should receive special attention under a "gender and tropical diseases" research program (WHO, 1995). Coreil *et al.*, (1998) showed that Haitian women's lives are heavily burdened both socially and economically by filariasis and collected the following statements:

"I can't harvest the garden produce, because you have to stand - I can't stand."

"I should be out selling, but I don't get up - my foot is too heavy."

"I ask for death because it makes me sick. I've been suffering from this for 33 years."

Bandyopadhyaya (1996) has found that the statement that the chronic manifestations of filariasis can cause "grave social wounds" (WHO, 1992, Remme *et al.*, 1993) is especially true for the women of India. Most affected women had very low self-esteem and some

had a fear of being rejected by their husbands because they were “no longer attractive”. In a society where arranged marriages are the norm, teenage girls with early chronic pathology consider their chances of being selected as a marriage partner to be seriously diminished and many of those with more advanced disease have given up hope of being married altogether. Dreyer *et al.*, (1997) found that there is a silent burden of sexual disability associated with lymphatic filariasis among men but it is seldom acknowledged because shame and fear caused “a conspiracy of silence” surrounding the man and his partner. They found that a wide range of disease-related problems including marriages devoid of sexual activity, complaints of painful intercourse in women whose partners had lymphedema of the penis, recourse to homosexuality by both male and female partners and suicidal thoughts in both sexes.

Children do not escape the social effects of filariasis either due to attacks of ADL interrupting their education or the medical/social effects of chronic pathology. Ramaiah *et al.*, (2000) report the tragic case of a 15-year-old boy with enlarged genitals having to give up his education because of shame, embarrassment and ridicule. Other boys with hydrocele were unable to cycle or walk to school, play, or take part in sport because of scrotal pain.

The Origins of Filariasis

Traditional beliefs about the aetiology of the disease vary widely from place to place. In many developing countries the disease is attributed to sorcery and other supernatural causes. The following examples are cited by Gyapong *et al.*, (1996). In Ghana people believe that elephantiasis of the leg is caused by stepping on “spiritual medicines” thrown on the ground by “juju men” (witch doctors) at funeral dances. Elephantiasis of the arm is believed to be caused by inadvertently picking up an animal tail dropped by a juju man. Others think that it is possible to get the disease from walking on ground on which certain herbs have been thrown or removing a thorn from one’s own foot rather than standing still and letting a “forest dwarf” remove it for you. Some people in the same community believe that the disease is hereditary and is passed to children *in-utero* via an infected father’s semen. Fevers “settling” in the scrotum or breast are blamed for some cases of elephantiasis. Men sometimes blame elephantiasis on their wives use of artificial flavouring agents such as monosodium glutamate or the provision of too much sweet food such as mango and sweet potato. The Duruma of coastal Kenya attribute elephantiasis to witchcraft, sexual intercourse or consumption of burnt food (Wijers, 1977).

Very few people in endemic areas know that mosquitoes transmit filariasis or believe that they can help prevent the disease by minimising contact with mosquitoes (Evans *et al.*, 1993). In a Malaysian study only 9 out of 108 respondents associated filariasis with mosquitoes and most people believed that you could

catch the disease by walking barefoot on dirty ground or consuming contaminated food or drink (Haliza and Mohd, 1986; Evans *et al.*, 1993). Similar views are held by a population in rural Thailand where only school children knew that mosquitoes transmit filariasis and that the disease could be prevented by protecting one’s self from mosquito bites. Most of the older people believed that the disease was hereditary or could be contracted through poor circulation, carrying heavy loads, prolonged standing, bathing in or drinking swampy water, personal contact, or sorcery (Rauyajin *et al.*, 1995). In Tanzania most people knew that mosquitoes spread malaria but very few knew that they also spread filariasis (Muhondwa, 1983). Ahorlu *et al.*, (1999) found that many villagers in a coastal Ghanaian community rejected mosquito transmission, believing instead in other physical, hereditary and spiritual causes. In Tahiti most people discounted the idea that mosquitoes were involved and filariasis was attributed to immersing an injured ankle in the sea or consuming contaminated food or drink (Kessel, 1957; Carme *et al.*, 1979). In the Philippines (Lu *et al.*, 1988; Schultz, 1988) it was mostly the educated people who knew the connection between mosquitoes and filariasis. Many people believed that coming into contact with cold water caused acute disease after heavy work and elephantiasis by taking long walks or squatting while mending fishing nets. Women were thought to catch the disease by washing whilst menstruating. Despite the lack of knowledge about mosquito transmission many people were able to identify that the people most at risk of developing filariasis were outdoor workers such as farmers, abaca strippers and fishermen. In a study in rural south India only 9.3% of apparently uninfected people and 20% of those with chronic filarial pathology knew that filariasis was contracted through mosquito bites. Other causes listed were occupational activity, bad drinking water and poor nutrition (Ramaiah *et al.*, 1996a). Similar findings were reported in Malaysia (Riji, 1986). This study also found that there was no significant difference in knowledge about filariasis between infected and uninfected persons. In Haiti the traditional view is that filariasis is caused by stepping on a magical powder placed on a footpath, injury to a limb, excessive cold or foreign objects such as glass entering the limb (Eberhard *et al.*, 1996; Coreil *et al.*, 1998). In a study among Indian women, Bandyopadhyay (1996) found that 92% of women interviewed did not know that mosquitoes transmit filariasis. Some believed that disease was hereditary while others blamed pregnancy, a previous injury or “bad spirits”. These cultural beliefs are often very deep rooted and may persist long after contact with European culture, religion and medicine and an understanding of them is vital when one is considering a control program. In Tanzania (Muhondwa, 1983) and Tahiti (Kessel, 1957; Carme *et al.*, 1979) for instance, most people steadfastly refused to believe mosquitoes spread filariasis even after an intensive education campaign.

“Where chemotherapy alone is the

main method of control people tend to discount the possibility of mosquito transmission arguing that if mosquitoes were the source of infection, mosquito control measures would have been promoted" (Evans et al., 1993).

Dunn (1976, 1979) and Mak (1986) observed that the impact of human behavioural and socio-economic factors pertinent to the control of filariasis have been largely ignored and there have been few attempts to bridge the gap between modern medical knowledge about filariasis and indigenous perception of the disease. It is also possible that traditional healing methods may influence the course of filarial disease. Ahorlu *et al.*, 1997 in a study from northern Ghana, suggest that scarification (multiple small cuts made on diseased parts of the body to let out "bad blood") may be a risk factor for developing acute adenolymphadenitis and elephantiasis presumably by providing an entry point for secondary bacterial infection, a known risk factor for elephantiasis (see below).

It is of considerable concern to the present writer that even although Papua New Guinea has one of the highest prevalences of filariasis in the world there appear to be no published studies on cultural attitudes to the disease.

Impacts of Cultural Awareness on Filariasis Control Programs

Social class, education, literacy and personality traits can have an influence the outcome of any disease control intervention and a lack of awareness of these issues can affect the success of a filariasis control program (Ramakrishnan 1960; Rao 1982). To quote Sunny *et al.*, (1986):

"An individual's attitude towards filariasis is shaped by the views and awareness of the community of which he/she is a member and may directly or indirectly the acceptance and non-acceptance of the offered control measures."

In the study by Sunny *et al.*, (1986) it was found that there was a marked difference in Diethylcarbamazine (DEC) acceptance between illiterate and literate and non-college educated and college-educated filariasis-infected individuals. Only 30.4% of illiterate suffers accepted the drug whereas 80% who were literate or who had attended school to college level accepted the drug. Interestingly, the incidence of refusal was higher in higher income groups than in low income groups and refusal was more common in females than in males.

The reasons for acceptance and refusal are given in Table 3. These tables provide some very revealing information about while some filarial control programs are successful and some fail. In this study the prime motivators for treatment acceptance was the fact that the medicine was free and it was delivered to their home. The second most important factor was fear of contracting the disease. By far the most important reason for not accepting DEC was the fear of side effects.

Table 3. Reason for acceptance of DEC (Sunny *et al.*, 1986)

Reason	No.	%
Knew it was an effective medicine for	58	17.1
Free medicine supplied at home	154	45.3
Knew that medicine is available to	18	5.3
Afraid that they will get filariasis	80	23.5
Insisted upon by head of household	2	0.5
Persuaded by health worker	24	7.1
No response	4	1.2
Total:	340	

It also highlights the problem of "occult" filariasis - it is difficult for people to accept treatment if they do not believe that they have the disease. This study clearly indicates that active and effectual health promotion is an essential part of any control program and the need to find an effective means of controlling the side effects of DEC without compromising its chemotherapeutic efficacy. Personality traits may also be an important factor in the success of filariasis control programs. In a study in the Philippines by Ventura (1986) 58 acceptors of blood screening and 50 non-acceptors matched for age and sex were given locally-devised personality scales which rated patience, obedience, responsibility, risk-taking, self-control, having an internal or external locus of control, being difficult to talk to and being daring. The study showed that typically, acceptors were those who were responsible, daring, patient, and who had an external locus of control. Being responsible meant that they realised the importance of taking control of their own health, and because they were also daring, they were more likely to have the strength of will to undergo the trauma of a blood test. It was also shown that males tended to comply with a request for a blood test because of a sense of obedience to authority whereas females tended to comply because they were more socially responsible. This study shows that cultural values, sociological attitudes and individual personality traits should be considered when planning a disease control program.

Socio-Economic Impact of Filariasis

"It is necessary to understand the social and economic impact of diseases in order to set priorities within the health system. This requires information on the types of symptom that are experienced; their prevalence, frequency and duration; and the nature of the associated social and economic costs. Much of this information is not available for lymphatic filariasis, and in its absence it is not surprising that the disease is not given higher priority in much of the endemic world" (Evans et al., 1993).

According to Gyapong *et al.*, (1996a, b), a major hurdle for procuring funding for control and research has been the lack of information regarding the social and economic impact of the disease. The estimated socio-economic impact of filariasis varies widely from study to study and documentary evidence of loss of production and income is difficult to find (Andreano and Helminiak, 1988). The current global estimate of the disability associated with lymphatic filariasis is 850 000 disability adjusted life years lost (Gyapong *et al.*, 1996a, c). This is probably a serious underestimate given the lack of recent prevalence information for many of the endemic countries. The global economic burden of the disease is also difficult to estimate but the annual economic cost to India alone is around US\$1,500 million (Addiss, 1998).

There is very strong evidence that late stage chronic disease with its accompanying disability reduces productive capacity (Kessel, 1957; Wijers, 1977; Wegesa *et al.*, 1979; Muhondwa, 1983; Evans *et al.*, 1993), but the economic impact is reduced somewhat by the fact that most people with advanced chronic pathology are beyond their most productive years (Wijers 1977). The impact on some communities can be severe. Gyapong *et al.*, (1996a, b) found that because of chronic filariasis 4.1% of the productive female labour force and 20% of the productive male labour force were disabled by between 10 and 60% and 20% people with advanced chronic disease often had to give up primary - food producing roles and take on a more sedentary occupation such as basket weaving. As a result, the net food production of the community was reduced (Gyapong *et al.*, 1996a, c). Mujinja *et al.*, (1997) have shown that elephantiasis, hydroceles and frequently experienced attacks of adeno-lymphangitis physically incapacitates affected individuals and greatly limits their participation in community economic and social activities. The need to employ labour from outside the family and rises in health care costs have had a negative impact on household income and production. Ramu *et al.*, (1996) found that the productive output of a group of male weavers with chronic filariasis was 27.4% lower than

that of uninfected age-matched controls.

As well as the direct economic impact of chronic pathology-induced disability there is the secondary impact derived from the effects of treatment, particularly the cost of surgery and the loss of income during the hospital stay. In one study of hydrocele surgery there was a very high rate of postoperative sepsis and days in hospital varied from 8 to 43 days (Muhondwa, 1983). After discharge, the men were advised to go on "light duties" for 6 months. In the long-term this period of reduced productivity may be balanced by increased productivity because of enhanced physical activity but evidence for this is entirely anecdotal (Muhondwa, 1983).

There is very little data on the economic impact of acute filariasis but it is obvious that there is potential for a severe economic impact, especially in populations who have a subsistence lifestyle. Kessel (1957) points out that it is very difficult to assess the economic impact of filariasis because most cases occur in rural areas where quantitative measurements of working time lost are seldom made. He states that before filarial control programs were introduced in French Polynesia acute filarial attacks occurred in 27% of the adult population with each individual having 1 to 12 attacks a year and each attack lasting from 1 to 3 days or longer. Fifteen percent of adult males suffered from hydrocele and another 14% had elephantiasis. The work output of those with mild or moderate elephantiasis or hydroceles was reduced and many of those with severe chronic pathology could not work at all and were cared for by their families. Kessel also quotes unpublished reports of a group of 600 Malaysian rubber tappers who reported 150 acute filarial attacks per year with a loss of 450 working days. Gyapong *et al.*, (1996a) found that the average time of incapacitation with Aden-lymphadenitis (ADL) was 3 days per person per year. This may not seem very high but most of the attacks occur during the rainy season - the time of peak agricultural activity. Total incapacitation at this crucial time could mean the family did not produce any food that year. Further evidence for the economic loss inflicted by filariasis comes from India where the estimated cost benefit of filariasis eradication for India in 1971 was said to be in the order of \$US133 million per year (Prescott, 1980). It would of course, be much greater, now. Panicker and Sabesan (1990) and Sabesan *et al.*, (1992) estimate that in the city of Pondicherry alone (population 370,000) 323,205 person working days are lost annually at a cost of \$US635,000 due to acute filarial attacks. A recent study by Ramaiah *et al.*, (1996b, 1997, 1998, 1999) investigated the impact of acute filariasis in south India and found it to be severe both in terms of the amount of money expended on treatment and the loss of individual and community earning capacity. About 66% of infected persons said that their disease impaired their economic activity and

some had completely given up their employment. Most patients however did not leave work but spent only $4.36 \pm$ hours per day on economic activity compared with 5.25 ± 3.52 hours worked by uninfected controls ($P < 0.01$). Performance of women's domestic chores was impeded and most affected people tried to avoid travel. The disability was worse in those with acute disease. In another Indian study, Sabesan *et al.*, (1992) found that an average of 26.5 man-days per patient were lost annually with a cost per person of around \$US20 per person. Bandyopadhyay (1996) in a detailed study of the effect of filariasis on the women of India found that many were unable to work in the house or fields for up to a week at a time, 4 to 5 times per year because of acute filarial attacks. The economic loss per person was estimated at about 24 rupees a day - a small individual sum perhaps, but a large impost on a poor family, and given the large population of India and the high prevalence of filariasis, a large impost upon the country as a whole. Women with chronic pathology, especially those with persisting lymphodema and/or elephantiasis often regarded themselves and were regarded by others as permanently disabled and unable to work. The economic benefits of filariasis

programs are well illustrated by a study in China. The cost of the control program in Miaoxi township, Huzhou City was around \$US4,000 but the cost saving brought about by the reduction of person-working days lost due to acute filarial attacks and lessened demand for medication was around \$US23,000 ie for every \$US1 spent on filariasis control there was a net benefit of about \$US5.75 (Shi *et al.*, 1995). A study in an urban area of Brazil showed that episodes of acute filarial disease frequently caused interruption of daily activities and kept people away from work from 3 to 15 days at a time. Alexander *et al.*, (1999) reports that in a highly endemic area of Papua New Guinea there are 0.31 episodes of adeno-lymphangitis per person per year in the leg alone and can cause complete incapacitation in the sufferer and curtailment of normal activities.

Nor is the impact of acute filariasis is not confined to the residents of endemic areas. A large number of American troops became infected with filariasis while stationed in the South Pacific during World War II. The socio-economic effects of these cases will be discussed in the section "filariasis in expatriates" below.

The Spectrum of Filarial Disease

Traditionally it has been accepted that five groups of people will be found in a filarial endemic area (Dasgupta, 1984; Partono, 1987; Ottesen, 1989; Evans *et al.*, 1993; Ottesen, 1993; Roberts and Janovy, 1997):

1. Exposed, but with no evidence of disease (so called "endemic normals").
2. Microfilaraemic asymptomatic.
3. Acute filarial disease with or without microfilaraemia.
4. Chronic disease, with or without microfilaraemia.
5. Tropical eosinophilia.

Endemic Normals

In a filarial context "endemic normal" refers to people who, despite constant exposure to filariasis and circulating antifilarial antibodies, are amicrofilaraemic and have no clinical evidence of disease (Ottesen, 1989; Weil *et al.*, 1996). Whether these people are truly immune to filariasis is debatable (Kazura *et al.*, 1993a; Weil *et al.*, 1996) have shown that many of these so called "endemic normals" have a cryptic infection because they have circulating filarial antigen which is identical to that found in people with microfilaraemia and ultrasound examination has also revealed that many men who are classified as "endemic normals have adult worms in their scrotum (Dreyer *et al.*, 1996a,c; Noroes *et al.*, 1996; Simonsen *et al.*, 1997b; Faris *et al.*, 1997; Suresh *et al.*, 1997). This indicates that many "endemic normals" rather than being immune, are harbouring adult worms, and rather than being "normal" or "immune" are in a pre-patent stage of the disease. A conclusion supported by Day (1991) who observed, "if you look carefully enough in any infected population, the endemic normal is a rare individual."

Evidence of Immunity

There is however some evidence, much of it from animal studies, that at least partial immunity to filariasis does occur (Denham, 1977; Oothuman *et al.*, 1979; Weil *et al.*, 1982; Kazura *et al.*, 1986; Hayashi *et al.*, 1984; Kazura and Davis, 1982; Kazura *et al.*, 1986; Nilsen *et al.*, 1988; Nanduri and Kazura, 1989). Work by Steel *et al.*, (1996) supports the concept of filarial immunity and suggests that it is probably T cell mediated. Michael and Bundy (1998) have found conclusive evidence that "herd immunity" does exist in filariasis and is modulated by vector exposure and biting rates. The reduction in microfilariae intensity in older people suggests that some degree of protective immunity may occur with long-term exposure (WHO scientific working group, 1981). Such individuals show high lymphocyte proliferative responses to filarial antigens (Nanduri and Kazura, 1989). Experimental animal models have shown that protective immunity is primarily directed against L3

stages (Oothuman *et al.*, 1979; Denham *et al.*, 1983; Yates and Higashi, 1985; Burkot *et al.*, 1996) and can be produced using radiation-attenuated but not dead L3 larvae (Story and Al-Mukhtar, 1982; Maizels and Lawence, 1991) and that the immunity provides cross protection between different filarial species (Oothuman *et al.*, 1979). Data from Papua New Guinea suggests that protective immunity to infective (L3) larvae is acquired over a number of years of exposure to infected mosquitoes and adults but not children, have high titres of antibody to L3 surface antigens (Day and *et al.*, 1991a,b). This stage-specific immunity may protect the individual from "super-infection" while leaving the resident parasites unharmed in the asymptomatic host (Maizels and Lawence, 1991). When worm burdens from infected individuals of different ages are quantified it can be shown that while childhood-acquired new filarial infections, the worm burden in adults remained the same irrespective of microfilariae density. Thus children with a high worm burden may have an even higher worm burden a year later whereas an adult with a low worm burden did not acquire any new infections (Day *et al.*, 1991b).

Asymptomatic Microfilaraemics

Asymptomatic microfilaraemia is often the most common manifestation of filariasis in many endemic populations and has been reported in children as young as 14 months (Lowman, 1944). It is often regarded as a "non-disease" because the individuals concerned have no inkling that their blood contains large numbers of microfilariae and this situation may persist for decades without any progression to clinical disease (Ottesen, 1992). Recent studies however have shown that asymptomatic microfilaraemia is not a benign phase of filariasis and that considerable lymphatic, tissue and organ damage may be occurring (Freedman *et al.*, 1994; Ottesen, 1994; Dissanayake *et al.*, 1995; Freedman *et al.*, 1995). Ultrasound has shown that approximately half of the men with asymptomatic microfilaraemia have nests of motile adult worms in their scrotal lymphatics: the "filarial dance sign" (Amaral *et al.*, 1994). For several centimetres either side of the worms the lymphatics are abnormally dilated but there is no evidence of an inflammatory response (Amaral *et al.*, 1994; Dreyer *et al.*, 1994). Lymphoscintigraphy has also revealed profound changes in asymptomatic microfilaraemics (Ottesen, 1994). Their lymphatics are markedly dilated with collateral channelling and increased lymph flow. By contrast, patients with elephantiasis (see below) show tortuosity, dermal back-flow, obstruction, stasis and poor regional node visualisation (Witte *et al.*, 1993; Dreyer *et al.*, 1994; Azoubel, 1996; Bernas *et al.*, 1996). The absence of an inflammatory response in the lymphatics near the adult worm in the asymptomatic stage of the disease suggests that the

lymphatic changes are induced by parasite or host-derived molecules acting directly on the lymphatics rather than an inflammatory reaction mounted by the host. Conversely, the changes seen in the tissues of those with chronic pathology are wrought by a mechanism which includes a vigorous inflammatory reaction (Ottesen, 1994). Exactly what triggers the change from one to another is one of the on-going debates in filarial research and will be further addressed in the sections on immunology and pathogenesis (below). People with asymptomatic microfilaraemia frequently show evidence of renal disease (see section on renal disease below).

Acute Filarial Disease

As mentioned in the section above on the socio-economic effects of filariasis, the main focus tends to be on the more spectacular chronic manifestations of filariasis such as elephantiasis. This is a focus that has to be altered if effective control of filariasis is to be achieved because:

- (a) In most endemic communities the number of people who have elephantiasis is relatively small compared with those who suffer acute disease and the effects of acute disease on both the individual and the community can be severe.
- (b) Health administrations often express the view that "filariasis is not a problem in our area" when what they really mean is that "elephantiasis is not a problem in our area". This distorts the epidemiological pattern of the disease and makes it difficult to convince the administrations in areas where there is very little chronic pathology that they need to undertake a filariasis control program.
- (c) Acute filariasis can occur without microfilaraemia and may be mis-diagnosed as malaria or other tropical disease. This leads to the wrong treatment being given and a waste of health care resources.

In endemic areas acute attacks can occur in infants as young as 3 months (Dasgupta, 1984) but they usually start appearing in older children and teenagers and continue throughout life (Nanduri and Kazura, 1989). They can occur in amicrofilaraemics or microfilaraemics and are common in people with chronic diseases (Evans *et al.*, 1993; Roberts and Janovy, 1996). In some cases filarial infection markers such as filarial antigen and antifilarial IgG4 antibody are absent. A possible explanation in these cases is an inflammatory response against injected L3 stages before the adult worm has become established (Addiss *et al.*, 1994) but extensive reviews of acute filariasis in American troops during World War II showed that adult worms were present in up to 30% of cases (Wartman, 1947).

The most common manifestation of acute filariasis is Adeno-lymphadenitis (ADL), which is characterised by intense lymphangitis, lymphadenitis with retrograde

extension from the affected node and reddening of the overlying skin. These attacks are usually accompanied by chills and fever (filarial or "elephantoid" fevers). In males there may be orchitis, epididymitis and acute transient hydroceles (Wartman, 1947; Nanduri and Kazura, 1989; Roberts and Janovy, 1996). (For a summary of the diagnostic criteria see the section below on filariasis in expatriates). Each episode lasts approximately a week and resolves spontaneously. Dunyo *et al.*, (1998) have reported that a common finding during the resolution phase is the onset of exfoliation of skin of the affected limb. Das Bidyut *et al.*, (1996) found that circulating Tumour necrosis factor (TNF) is increased during acute filarial episodes suggesting that TNF may play a role in inducing clinical systems and that TNF inhibitors may have a place in its management.

Acute filarial attacks can be extremely debilitating and many people are confined to their houses and unable to take part in productive activity. During a study of 1,300 filariasis patients, Pani and Srividya (1995) found that ADL attacks increased in frequency as the severity of lymphodema increased. In patients with grade 1 lymphodema (mean duration 47.4 days), the mean number of ADL attacks per year was 4.9, while with Grade 2 lymphodema (mean duration 6.2 years), 5.5 per year and Grade 3 lymphodema (mean duration of lymphodema 8.6 years) 10.4 per year. Filarial fevers may occur in the absence of other symptoms and is often treated as "malaria". Health care records which show a high proportion of "antimalarial resistant" fevers is sometimes the first indicator that acute filariasis occurs in a community (personal observation from Papua New Guinea). It has been suggested that the fever is an allergic reaction to excreted worm antigens (Ottesen, 1984; Kar *et al.*, 1993) and studies in ferrets have shown that severe lymphodema is associated with high levels of filarial-specific IgE (Crandall *et al.*, 1993). (See section on humoral immunity below). It has been shown in *in-vivo* studies that disodium cromoglycate selectively inhibits IgE production and enhance the production of IgE-blocking IgG4 antibody by human B cells *in-vitro* (Kimata *et al.*, 1991). It would be interesting to see if the same effects could be demonstrated with lymphocytes stimulated with filarial antigens. Shenoy *et al.*, (1995) found that there was an association between the degree of oedema and the frequency of ADL attacks. They also found that DEC alone or DEC plus antibiotics did not reduce the number of ADL attacks but simple hygienic measures combined with topical antibiotics and antifungal creams were effective in reducing the attacks. Shenoy *et al.*, (1999) have found that care of the feet and local and systemic antibiotic use is an important tool in reducing the frequency and severity of ADL attacks. This suggests that the aetiology of ADL is a complex combination of filarial infection *per sé* and secondary bacterial and or fungal infection (Dreyer *et al.*, 1999c). A important new development has been the finding that the inflammatory responses induced by filaria may be strongly mediated by the lipopolysaccharide-like

substances produced by *Wolbachia* symbiont bacteria carried by the female worms (Taylor *et al.*, 2000). Das *et al.*, (1996) have shown a relationship between Tumour necrosis factor (TNF) and the severity of clinical symptoms of acute disease. Expansion in this area of knowledge would be worthwhile - what happens to phospholipase A₂ (PLA₂) which is modulated by TNF and what is the role of specific TNF and PLA₂ inhibitors in the management of ADL?

Chronic Disease

It has been generally accepted that most people with chronic disease are amicrofilaraemic and that patent infection is negatively related to chronic disease (Jordan, 1955; Nanduri and Kazura, 1989). This basic tenet of filariology has been challenged by metanalysis studies of previously published studies by Michael *et al.*, (1994) who found that, contrary to expectation, that there was an equal prevalence of disease in microfilaraemics and amicrofilaraemics. Indeed, in some studies, especially those involving hydroceles, disease was more prevalent in microfilaraemics than amicrofilaraemics. This demonstrates the need for "traditional filarial dogma" to be re-evaluated in the light of modern diagnostic, immunological and statistical modelling techniques.

Often the first sign of the onset of chronic disease is persistent limb lymphodema or hydroceles after an acute attack (Evans *et al.*, 1993) and the process of development from lymphodema to hydroceles or elephantiasis is a gradual one. The lymphatic dilation, co-channelling and rapid lymph flow seen in asymptomatic microfilaraemic is replaced by tortuosity, dermal back-flow, obstruction, stasis and poor regional node visualisation (Witt *et al.*, 1993; Dreyer *et al.*, 1994). The switch from asymptomatic filariasis to chronic pathology is also accompanied by profound changes in cellular and humoral immunity (see section on immunity below).

There have been several methods devised to try and increase lymph drainage and prevent lymphedema. Hyaluronidase was tried with some success by Jordan (1959) and coumarin, a drug which increases macrophage-associated proteolysis and reduces stasis of protein in the tissues, has been shown to be of benefit in reducing lymphodema and elephantiasis (Jamal *et al.*, 1989a,b; Casley-Smith *et al.*, 1993). Surgical procedures such as the establishment of a shunt between the lymphatic and venous system or the lymphatics and the omentum (Goldsmith *et al.*, 1967; Goel and Misra, 1968; Goldsmith, 1974; Jamal, 1989a; Abalmasov *et al.*, 1994; Binoy, 1998) have been shown to be effective in reducing lymphodema in some patients, and lymphosuction has been proposed as a new treatment modality for chronic lymphedema (Clodius, 1998; Agarwal *et al.*, 1998).

Overall, hydrocele is the most common form of chronic Bancroftian filariasis and in parts of Africa it predominates. Hydrocele is said to have affected up to 70% of males along the Tanzanian coast (Wegesa *et al.*, 1979 and 60% of males in coastal Kenya by the time they reach age 70 years (Wijers, 1977; Estambale *et al.*, 1994a,b). Gyapong (1998) found a very strong positive association between hydrocele and microfilaraemia and only a weak association between elephantiasis and suggests that the pathophysiologies of the two conditions may be different. The strong association between community infection and hydrocele in Ghana, is being used for rapid community assessment of filariasis prevalence (Gyapong *et al.*, 1998a,c).

Elephantiasis is relatively uncommon in Africa and is mostly found in the areas with the most intense transmission (Wijers, 1977; Evans *et al.*, 1993). Hydrocele is also common in India. In Pondicherry, India, 45% of males have a hydrocele by the age of 60 years; Pani *et al.*, 1991) but a whole range of other chronic manifestations ranging from chronic lymphodema to elephantiasis are also seen (Evans *et al.*, 1993). In some populations, despite a high prevalence of microfilaraemia and filarial antigenaemia, filariasis is a mostly a cryptic disease and the number of people with chronic pathology is relatively low. For instance, the prevalence of filarial antigenaemia in many parts of Papua New Guinea is around 80% yet elephantiasis occurs only in scattered foci and is nowhere common (Kazura, 1984). This again draws attention to what I like to call the "elephantiasis only syndrome" whereby we may be tempted to think that because elephantiasis is seldom seen, filariasis is not a community problem.

The first evidence of impending elephantiasis is persisting lymphodema, which is followed by fibrous infiltration, and thickening of the skin. Granulomatous, fat and fibrous connective tissues are laid down in the subcutaneous tissues (Burri *et al.*, 1996). Fissure and ulcer formation (figures 9 and 10) allows access of bacteria and fungi, which have been shown to play an important role in the pathogenesis of elephantiasis.

Figures 9 and 10: Fissure and ulcer formation.




It has been shown that the risk of acquiring chronic filarial disease rises with increasing microfilariae density (Ravindranathan *et al.*, 1980). Chronic manifestation can be graded as follows (Table 4).

Table 4. Grading of lymphedema (Gyapong *et al.*, 1994; McMahon *et al.*, 1996)

Manifestation	Grade. Symptoms
Elephantiasis of limb	0. Normal
	1. Loss of contour or lymphodema
	2. Thickened skin and/or loss of elasticity
	3. Evident elephantiasis
Hydrocele	0. Normal
	1. Swelling of spermatic cord
	2. Swelling up to 10 cm in diameter
	3. > 10 cm in diameter
Scrotal elephantiasis	0. Normal
	1. Lymphodema
	2. Thickened skin and/or loss of elasticity
	3. Evident elephantiasis

Tropical Pulmonary Eosinophilia (TPE).

This is the least common manifestation of filariasis and is at the opposite end of the immunological spectrum from asymptomatic microfilaraemia. In TPE there is a severe hypersensitivity response with marked eosinophilia (figure 11), extreme levels of serum IgE and high titres of antifilarial IgG and IgE.



Circulating microfilariae are not usually found (Van der Sar and Hartz, 1945; Ottesen *et al.*, 1982) but adult worms may be detected by ultrasound (Dreyer *et al.*, 1996d). Paranjape *et al.*, (1985) found that although the mean level of anti-filarial IgG antibody is very high, a small number of patients who had clinical TPE had lower than expected antibody titres.

Clinically TPE is characterised by nocturnal coughing and asthma. Pulmonary infiltration gives rise to a characteristic x-ray picture ("snow flake" lung). Accumulation of eosinophils in the lung with the release of cationic proteins and free radicals can lead to fibrosis and permanent lung damage (Pinkston *et al.*, 1987). The culprit is thought to be an IgE-producing allergen derived from microfilariae (Lobos *et al.*, 1992). For some reason the blocking effects of anti-filarial IgG4 do not seem to work in TPE patients. Tropical pulmonary eosinophilia is most common in South India (Ray *et al.*, 1993) and parts of Southeast Asia but is rarely seen in Papua New Guinea (Nanduri and Kazura, 1989) and Africa (Magnussen *et al.*, 1995).

Recent Ideas on the Classification of Disease


The introduction of assays for circulating filarial antigen (More and Copeman, 1990; Weil, 1987; Weil *et al.*, 1987), the discovery of lymphatic pathology in many "asymptomatic microfilaraemics" (see above), a lack of evidence that there is a negative association between microfilaraemia and disease, and the recognition of the role of bacterial infections suggests that the old classification based on the presence or absence of microfilaraemia and/or chronic pathology is outdated. Freedman (1998) proposes a new classification: asymptomatic infected individuals who are antigen negative; individuals who have overt filariasis and active infection (antigen positive) and individuals who have overt filariasis without active infection (antigen negative). Freedman defines overt filariasis as any of the clinical manifestations associated with the known clinical spectrum of filariasis: adenolymphangitis, hydrocele or elephantiasis (Figure 5).

As cytokine responses appear to segregate according to the presence of Cag rather than the results of the clinical examination, patient classification must

account for both factors. Because all filariasis patients have underlying pathology, this term as used in earlier classifications is now unhelpful in segregating patients. Three groups of individuals are proposed as appropriate for detailed immunological study: asymptomatic microfilaraemics; those with clinical filariasis with active infection (Cag+); and those with clinical infection without active infection (Cag-).

Microfilariae prevalence status refers to microfilaraemia as detectable by conventional parasitological methods, so that some infected individuals may have microfilaraemia undetectable by these methods. Abbreviations: IFN- γ , interferon γ ; IL-4, interleukin 4; MF microfilaraemia (Freedman, 1998, used by permission).

Figure 12. Before the availability of circulating filarial antigen (Cag) assays to sensitively assess current infection status, groupings of humans with filariasis for immunological study have been based on detection of circulating microfilariae (a). Infected individuals with detectable microfilaraemia but without any outwardly discernable manifestations of lymphatic filariasis have been called “asymptomatic microfilaraemic individuals” (top left). Although clinically, filariasis is a spectral disease, those with clinical evidence of lymphatic insufficiency have mostly been grouped together as “chronic pathology” (top right) and have thought to be generally amicrofilaraemic. Separate immunological data on chronic pathology patients who are also microfilaraemic have not been readily available. Infection status as determined by recently available Cag assays permits a different framework for patient classification to be proposed (b).



(With permission of Professor D. Freedman)

Filariasis and Other Illnesses

Filariasis and Asthma

A study from Sri Lanka suggests that filariasis may be a trigger for bronchial asthma (de Sylva *et al.*, 1990). This study of children with cough and wheeze suggests that more than 1/3 had filarial infection. Symptoms improved after DEC therapy probably due both to the elimination of the microfilariae and the anti-inflammatory effects of the DEC. The authors postulate that atrophic individuals have a heightened response to filarial antigen, which may trigger asthma.

Chyluria

The presence of chyle in the urine follows the rupture of dilated lymphatics in the bladder or renal pelvis into the urinary system is a rare but spectacular complication of filariasis (Cahill, 1965; McMahon and Simonsen, 1996).

Renal Disease

There have been several reports of renal abnormalities associated with filarial infections in both humans and animals: hematuria, proteinuria, nephrotic syndrome and glomerulonephritis have all been recorded (Sapico and Baltazar, 1967; Antani and Krishnamurthy, 1971; Casey *et al.*, 1972; Simpson *et al.*, 1972; Crowell *et al.*, 1973; Klei *et al.*, 1974; Mukherjee *et al.*, 1974; Chugh *et al.*, 1978; Dufor and Lepage, 1978; Date *et al.*, 1979; Srivastava *et al.*, 1979; Waugh *et al.*, 1980; Yap *et al.*, 1982; Date *et al.*, 1983; Ormerod *et al.*, 1983; Ngu *et al.*, 1985; Langhammer *et al.*, 1997; Barsoum, 1999). Circulating immune complexes containing filarial antigen have been blamed for the renal damage (Au *et al.*, 1981; Dissanayake *et al.*, 1982; Gajanana *et al.*, 1982; Prasad *et al.*, 1983a,b,c; Das *et al.*, 1987; Prasad and Harinath, 1988; Lunde *et al.*, 1988; Lutsch *et al.*, 1988) and filarial antigen has been located in a renal biopsy (Ngu *et al.*, 1985). In a study of 20 patients with asymptomatic microfilaraemia Dreyer *et al.*, (1992) found that 7 had microscopic hematuria and 4 had proteinuria but their creatinine clearances were normal. Ten individuals who had not shown hematuria and 10 who had not shown proteinuria developed it after DEC treatment but all patients in the study had normal urinalysis 2-3 weeks after therapy. Oliveira (1998) has postulated that one of the causes of membranous nephropathy, an IgG4-mediated disease, are helminths. Amyloidosis also leads to renal disease and filarial infection has been definitely linked to filarial infection in Mongolian gerbils (Crowell *et al.*, 1973) and an association between amyloidosis and human filariasis has been found (see below).

Amyloidosis

Amyloidosis is unusually prevalent in Papua New Guinea (Anders *et al.*, 1976a,b). Cooke and Champness (1970) for instance found amyloid deposits in 7.3% of 1,101 post-mortem material they examined.

Prior to 1978 the tropical diseases known to cause secondary amyloidosis included tuberculosis, leprosy, schistosomiasis, leishmaniasis and chronic suppurative infections (McAdam, 1978). A strong association has been described between filariasis and amyloidosis in a village survey in the Upper Fly region of the Western Province of Papua New Guinea (McAdam, 1978). Six people out of the surveyed population of 224 had evidence of both filariasis and amyloidosis. The level of the amyloidosis-precursor protein SAA (Anders *et al.*, 1976b; Levin *et al.*, 1973) was highest in 2 people who suffered recurrent episodes of filarial adenolymphadenitis and in one of these the level of protein SAA rose 10 fold during a acute attack lasting 3 days. Clearly, this is another health threat imposed by filariasis and another reason why filariasis should be eradicated.

Arthritis and Arthralgia

Filariasis can present with rheumatic features (Adebajo, 1996) and mono-arthritis of a knee or ankle joint is common in filarial-endemic areas. Chaturvedi *et al.*, (1993) investigated 19 cases of arthritis of unknown origin in children and found that 16 of them had filariasis. Nine of these cases were treated with DEC and the arthritis resolved. Low *et al.*, (1996) describe a patient with a hot swollen foot and eosinophilia that responded to DEC. Magnetic resonance imaging (MRI) may be a useful tool for diagnosis (Blacksin *et al.*, 1999).

Polymyositis

Poddar *et al.*, (1994) reported two cases of polymyositis, which presented with generalised painful swelling and weakness of the muscles. Serum creatinine kinase was elevated and inflammatory myopathy was present in biopsies. *W. bancrofti* microfilariae were present in the peripheral blood. The patients responded better to DEC and steroids than to steroids alone and clinical improvement was associated with the clearance of the microfilariae. A patient with acute filarial myositis that responded rapidly to DEC therapy was reported by Sundaray *et al.*, (1992).

Central Nervous System (CNS) Disease

Central nervous system involvement has been reported in Loasis (Brunetiere, 1913; Peruzzi, 1928; Tha Mya, 1928; Van Bogaert *et al.*, 1955). Epileptic seizures decreased after Ivermectin therapy for onchocerciasis (Kip *et al.*, 1992) and a similar finding associated with lymphatic filariasis has been reported by Adamolekun *et al.*, (1993). Garg *et al.*, (1996) reported a Pott's paraplegia-like neurological complication in a filariasis patient. Orihel (1973) reviewed cases of neurological disease which were first thought to be due to infection with *D. perstans* but he was of the opinion the morphology of the microfilariae was consistent with *Meningonema peruzzi* a filaroid species usually found in old world monkeys. Psychotic manifestations were also common in United States servicemen who contracted filariasis during the Second World War (see section "filariasis in expatriates" below) and "nervous symptoms" were noted in Ceylonese students with microfilaraemia by Wijetunge (1967a,b) Fonticiella *et al.*, (1995) reported a case of a new born with intracranial, partially calcified extra-axial collections, which were shown to be due to filariasis transmitted from the mother. Microfilariae of *B. malayi* have been demonstrated in the brains of experimentally infected animals (Paily *et al.*, 1995).

Ocular Involvement

Gupta *et al.*, (1992) report a case of retinal pigment epithelium inflammation and vasculitis in a microfilaraemic patient which rapidly resolved after anti-filarial treatment and Mathai and David (2000) reported a patient who had an adult *W. bancrofti* removed from his eye.

Guillain-Barre Syndrome

Bhatia and Misra (1993) reported a patient who developed recurrent Guillain-Barre syndrome after acute filariasis.

Filariasis and Leucocytes

Apart from eosinophils (see below), filariasis seems to have little effect on the total or differential leucocyte count. There is one paper by Adhikari *et al.*, (1994) which found that the neutrophil percentage was significantly reduced while lymphocyte and eosinophils were increased in filariasis.

Filariasis and Eosinophilia

Eosinophilia is defined as an increase in the number of peripheral blood eosinophils (Zucker-Franklin, 1983). Eosinophilia can be caused by: Parasite infections, allergic disorders, dermatitis, malignant hyper-eosinophilic syndromes and tumours (Zucker-Franklin, 1983). Eosinophilia is often seen in filariasis and very

high eosinophil counts are usually found in tropical eosinophilia syndrome (Van der Sar and Hartz, 1945; McMahon and Simonsen, 1996; Roberts and Janovy, 1996) but many people with filariasis are hosts to other helminth parasites also and it is difficult to measure the direct association between eosinophilia and filariasis. Wong and Guest (1969) found that filarial-associated eosinophilia was present in 11 out of 16 individuals who had strong anti-L3 stage antibodies. By contrast, only 3 out of 14 people with microfilaraemia and 9 out of 16 people with elephantiasis had eosinophilia. Filariasis however, should always be considered when investigating a patient, including returned travellers, for filariasis (Moore and Nutman, 1998).

Cardiac Disease

Tropical endomyocardial fibrosis is common in some tropical areas and is believed to be associated with chronic hyper-eosinophilia caused by helminth infections, including filariasis (Andy *et al.*, 1998). Filarial infection is also a rare cause of pericarditis (Foster, 1956; Chakravarty and Gupta, 1966).

Splenomegaly

Schuurkamp *et al.*, (1987, 1992) have shown that filariasis is an important cause of splenomegaly and that it can be reversed by DEC therapy. Traumatic rupture of a filarial spleen has also been recorded (Behara *et al.*, 1995) and Piyaratn and Pradatsundarasar (1961) reported a case of micro filarial granuloma of the spleen. Splenectomy has been shown to increase microfilariae density in animal studies (Hawking, 1962b; Nooruddin and Ahmed, 1967) but there appears to be no information on the effects of splenectomy in the human.

Filariasis and Anaemia

There has been one report of a significant association between filariasis and anaemia. Saxena *et al.*, (1981) found that 70% of symptomatic cases had an Hb <100g/l. None of these had any other helminth infection. By contrast, none of their control cases had an Hb <100g/l.

Filariasis and Platelets

No association between thrombocytopenia and filariasis has been established but there is evidence that circulating microfilaria modify platelet function. A detailed study by Lui and Weller (1992) showed that *Brugia malayi* microfilariae per 104 platelets completely inhibited platelet aggregation with collagen, arachidonic acid and ionophore A231187, thromboxane generation and thromboxane release when incubated with human platelets. Microfilariae also inhibited the aggregation of platelets in platelet rich plasma stimulated by adenosine diphosphate, compound U46619, or platelet-activating factor. The

effect required close proximity but not direct contact between the microfilariae and was mediated by parasite-derived soluble factors of low molecular weight. This phenomenon explains why platelets do not adhere to microfilariae and why filariasis patients do not seem to suffer from thrombo-occlusive events despite having a large number of circulating foreign objects (microfilariae) that could act as a stimulus for such events. Microfilariae also utilise exogenous and endogenous arachidonic acid to generate and release substances that influence platelet metabolism and function - prostacyclin and prostaglandin E2. The effects on platelet function wrought by microfilariae do raise the question about the possibility of a bleeding tendency in filariasis patients and suggests a study may be worthwhile. Anti-platelet aggregation drugs are widely used to prevent thrombo-occlusion in general medicine and it is possible that if the platelet inhibitor from the filarial parasite could be isolated it could form the basis for a new drug. Paradoxically, infection with the dog heartworm *Dirofilaria immitis* is reported to enhance platelet reactivity (Boudreaux *et al.*, 1989, 1991). Another interaction between microfilariae and platelets has been suggested by the finding by Pancre *et al.*, (1988) that platelets isolated from filariasis patients are cytotoxic to microfilariae *in-vitro* and that

this activity is dependant upon the presence of anti-filarial IgE. The cytotoxic effect was strongly inhibited by antigen-stimulated T lymphocyte supernatants from filarial patients - another protective mechanism induced by parasite-induced immune system modulation.

Miscellaneous Complications

Other complications recorded include acute abdomen in pregnancy (Charles, 1976), cystitis with urethral obstruction (Devasia *et al.*, 1998), microfilariae in ascitic fluid (Handa *et al.*, 1996), fibrosing mediastinitis (Gilbert and Hartman, 1996), cases of tropical vaginal hydroceles (Sivam *et al.*, 1995) and bladder pseudotumours (Gourlay *et al.*, 1999). Wijetunge (1967a,b) studied the clinical manifestation of 212 microfilaraemics and suggests that ocular complications, enlargement of the thyroid gland, nocturnal chest pain, buccal ulceration, parotitis, gastrointestinal pain and arthralgia can be caused by filariasis. There is one report of an association between mild Vitamin A deficiency and onchocerciasis (Zambou *et al.*, 1999) but no comparable studies have been done in lymphatic filariasis.

Filariasis in Expatriates

When people from a non filarial-endemic area move to an endemic area two different patterns of disease are seen depending on whether the migrants are visitors who stay only for a short period or migrants who remain permanently in the endemic area.

Filariasis in US World War II Forces

Many thousands of cases of filariasis occurred in the United States forces stationed in filarial-endemic areas

during World War II, especially in the Pacific Theatre, Papua New Guinea and the Solomon Islands and filariasis was among the leading causes of medical evacuation in the South Pacific (Swartzwelder, 1963). Coggeshall (1945, 1946) and Wartman (1947) state that about 38,300 naval personnel were exposed to filariasis and of these, 10,421 were diagnosed as having the disease. In one unit alone, stationed in a highly endemic area of Samoa, 70% of the exposed troops became infected (Beaver, 1970).

A list of relevant papers is given in Table 5.

Table 5. Filariasis among United States servicemen in the Second World War: number of cases, places and length of exposure and incubation period (Wartman 1947)

Author and Date	Place of Infection	Exposure Length (mths)	Incubation Period (mths)	Number of Cases
Behm and Hayman, 1946	Tonga-Tabu	12	2-18	532
Burhans et al., 1944	South Pacific	nr	3-21.5	46
Coggeshall 1945	Samoa 96%	1-27	9	1200+
	Solomons 4%	av 12.5		
Coggeshall, 1946	South Pacific	Various	1-12	2288
Dickson et al., 1943	Samoa	nr	1-9.5	251
Englehorn and Wellman, 1945	South Pacific	nr	3-14	127
Flynn, 1944	Samoa	2-22	1-5	125
Fogel and Huntington, 1944	South Pacific	nr	nr	large
Glauser, 1945	Samoa	1-23 (av 9)	2-22	172
Goodman et al., 1945	Samoa	2-23	nr	145
Haviland, 1944	South Pacific 97%	1-18	1-19	228
Hodge et al., 1945	Tonga-Tabu	12	nr	62
Huntington et al., 1944	South Pacific	nr	3.5+	251
Johnson, 1944	Samoa	nr	av 9.5	189
Leede and Josey, 1945	Samoa	16.3	2-22	100
Michael, 1944	Samoa	nr	7-9	120
Rifkin and Thompson, 1945	Samoa	13-30	14-20	30
Saphir, 1945	South Pacific	3-6	4-9	35
Smith, 1945	Samoa 97%	3-17	nr	737
Wartman and King, 1944	Samoa	2-11	3+	268
Wartman, 1944	Samoa	1-5	nr	17

Coggeshall, (1946) made a prospective and retrospective study of 2,288 cases. Wartman and King (1944) record 268 American troops with filariasis and state that one fifth of the troops station on a particular island were admitted to hospital with symptoms of filariasis. Engelhorn and Wellman (1945) reported 127 cases of filariasis at a field hospital in Samoa over a 4-month period. On Tonga-tabu Island in the central Pacific 532 men were diagnosed with filariasis in a single year (Behm and Hayman, 1946). Glauser (1945) reported 72 cases in returning marines during a 6-month period. Two thousand five hundred and ninety five cases of filariasis were admitted to the

United States Marine Corps hospital at Klamath Falls, Oregon during a 17-month period (Coggeshall, 1946). Smith (1945) recorded 737 cases which were acquired in Samoa, Wallis Island, New Caledonia and Bougainville and Goodman *et al.*, (1945) studied 145 soldiers with filariasis admitted to Harmon General Hospital who had served in the Cook Islands, Bora Bora Island, Samoa, Woodlark Island (Milne Bay Province, Papua New Guinea) and Ellice Island. Hodge *et al.*, (1945) reported 62 cases at a single field hospital, Flynn (1944) reviewed 125 cases of filariasis admitted to hospital from a single marine battalion that had been stationed in Samoa, Vanuatu and the

Solomon Islands and Dickson *et al.*, (1943) reviewed 251 patients addicted to a Samoan mobile hospital in a single year. Two hundred cases were studied by Thompson *et al.*, (1945), 189 cases by Johnson (1944), 268 by King (1944) and unstated numbers by Fogel (1945) and Michael (1944). According to Swartzwelder (1963) filariasis caused a significant loss of manpower and money and seriously compromised the fighting ability of American army units in the South Pacific. He gives the following as an illustration: The 134th Field Artillery Battalion and the 404th Combat Engineer Company were stationed on Tongatabu Island in the Central Pacific from May 1942 until May 1943 and then on Woodlark Island, Milne Bay Province, Papua New Guinea from July 1942 until January 1944. Because of the large number of personnel infected with filariasis the units were withdrawn to Sydney, Australia where the prevalence of filariasis (based on clinical grounds) among the evacuees was found to be 65% in the 134th Battalion and 55% in the 404th Company. The commanding general of the 6th United States Army recommended that the entire units be withdrawn from active service and return to the United States because their fighting efficiency had been seriously impaired and their rehabilitation would extend over an indefinite period of time. There was also a high economic cost to the American forces - estimated by Napier (1947) at around 100 million dollars. When introducing a speaker at the annual meeting of the American Society of Tropical Medicine and Hygiene the moderator remarked "that in World War II filariasis cost the Navy the equivalent of a battalion of marines for 4 years and enough money to run the Naval Medical Research Institute for 25 years." (American Society for Tropical Medicine and Hygiene, 1957). Filariasis is often thought of as a slowly developing disease, which requires a long period of exposure in order to become infected. A notable feature of the above table is the short exposure time, in some cases only 1 month.

Wartman (1947) provides an extensive epidemiological and clinico-pathological review of

over 6,678 cases recorded by the authors referenced in Table 6 and other clinico-pathological reviews are provided by Thompson *et al.*, (1945), Fogel and Huntington (1945) and Beaver (1970). One of the most notable features is that microfilaraemia was only detected in about 20 cases (Wartman, 1947; Beaver, 1970). Despite the failure to demonstrate microfilariae in most cases there is little doubt about the diagnosis. The clinical signs and symptoms were typical of acute filariasis and, most telling of all, adult worms were recovered in about 30% of the cases (Wartmann, 1947). There are at least two possible explanations for the lack of circulating microfilariae. In expatriate cases the adult worm induced a marked inflammatory reaction with a dense collection of inflammatory cells.

It is possible that the microfilariae get trapped within the lesion and are unable to make their way into the circulation. Another, probably more plausible explanation is that the immune response in previous unexposed expatriates is more vigorous than that seen in "immune tolerate" residents (see below) and the microfilariae are rapidly destroyed. When the rare cases of microfilaraemia did occur the microfilariae showed the same diurnal periodicity pattern that was usually seen in the native population (Eyles *et al.*, 1947).

Almost all the cases occurring in the American troops the cases could be termed "acute filariasis" and would fall into the modern category of filarial adenolymphangitis. It was observed by Huntington *et al.*, (1950) that the signs and symptoms were identical to the syndrome called "mumu" present in Samoan indigenous populations. The attacks commonly lasted from 3 to 5 days with occasional incidences of 2 weeks, or very rarely, one month (Behm and Hayman, 1946). Recurrent attacks were common, occurring at intervals of a few weeks to several months. Hard exercise, fatigue and a hot climate were thought to be precipitating factors (Wartman, 1947). The diagnostic criteria used by Hodge *et al.*, (1945) are provided in Table 6.

Table 6. Diagnostic criteria used for filariasis cases in us forces (Hodge *et al.*, 1945)

A. Involvement of the Genitalia:	B. Involvement of the Superficial Lymphatics of the Extremities
<ol style="list-style-type: none"> 1. Episodes frequently follow periods of manual labour 2. Testicular pain was the most common complaint. 3. The pain may radiate up the spermatic cord or may first appear in the lower abdomen and radiate down the spermatic cord to the testicle. 4. The spermatic cord and epididymis are thickened and indurated but the vas deferens is not involved. 5. Acute hydrocele is often present. 6. Pain in the mesial aspect of the thigh on the affected side may be present. 7. Scrotal edema is often present. 	<ol style="list-style-type: none"> 1. Pain in the extremity radiating distally is the usual complaint. 2. Pain may be first noticed in the proximal lymph nodes. 3. The affected extremity becomes edematous. 4. A definite "red streak" characteristic of acute lymphangitis appears proximally. The affected lymphatic is palpable and usually only a single vessel is involved. 5. There is increased local heat. 6. The lesion extends centrifugally. 7. Resolution begins proximally and extends centrifugally. 8. The lymph nodes draining the affected area are enlarged and tender. 9. Severe constitutional manifestations are usually absent.

The most common presenting signs and symptoms in the American armed forces cases were pain and swelling of the genitalia closely followed by lymphangitis of the arms and legs. Lesions of the genitalia in order of frequency were funiculitis, epididymitis, scrotal oedema and inflammation, lymph scrotum, orchitis, hydrocele and varicocele. Lymphangitis occurred in between 51 and 80% of cases. The most common site was the arm, followed by the leg, groin, buttock and abdomen. An important feature was the tendency for the inflammation to spread in a retrograde direction. Some authors reported skin temperature increases in the effect area but others did not. Lymphodema was occasionally observed but was not common. The prevalence of lymphadenitis varied widely from group to group and ranged from 85% to 7%. Where lymphadenitis was present the most commonly affected nodes were the epitrochlear and the inguinal but other nodes were often involved. Evidence of hypersensitivity was commonplace with many patients suffering urticaria and transient swellings. Some of the latter resembling the "Calabar swellings" or "fugitive swelling" usually associated with *Loa loa* infections. A small number of patients developed a macular erythematous rash. Hypersensitivity-type manifestations were only seen while the patients were in an endemic area, which raises the possibility that they were caused by a reaction to mosquito-injected L3 larvae. One study reported a high incidence of conjunctivitis and some incidences of phobophobia. Filarial fevers were very uncommon and only one case of filarial abscess was reported. Systemic symptoms - malaise and fatigue, were very commonly associated with acute attacks.

The most common laboratory finding was eosinophilia and in a few cases there was a very high eosinophil count associated with a chronic cough - a hallmark of tropical eosinophilia syndrome (King, 1944; Hirst and

McCann, 1945; Pinkston *et al.*, 1987).

Psychological disturbances were common and are discussed by Dickson *et al.*, (1943), Rome and Fogel, 1943, Coggeshall (1945) Englehorn and Wellman, (1945), Glauser (1945), Goodman *et al.*, (1945), Smith (1945), Zeligs (1945), Behm and Hayman (1946); Coggeshall (1946) and Swartzwelder (1963). Typical manifestations were depression, irritability, concentration difficulty, nervousness, anxiety, sleep disorders and fear. It is possible that filariasis *per sé* does produce some mood swings and depression (Rome and Fogel, 1943) but many of these symptoms may have been precipitated by observing Polynesian and Melanesian nationals with disfiguring diseases such as severe hydrocele and elephantiasis and a fear that they would as one writer puts it "go home with their scrotum in a wheelbarrow" or suffer loss of sexual function. As it turns out however, these fears were unfounded. No cases of chronic pathology occurred. Whilst Behm and Hayman (1946) reported 33 patients who believed their sexual potency had been impaired by filariasis and Smith (1945) reported that many of his patients reported a loss of libido, there is no evidence of long-term effect on sexual function and the prevalence of impotence and sterility was no higher in filariasis patients than in the general population (Wartman 1947). Coggeshall (1946) reported that 107 of 504 marriages among marines with a diagnosis of filariasis resulted in pregnancy within 6 months.

How long do the signs and symptoms of filariasis persist after removal from an endemic area? Behm and Hayman (1946) and Wartman (1947) reported that in most cases, even without treatment, attacks became progressively milder when patients were removed from the endemic area. Most men were free of symptoms after they had been out of the endemic area for 20

months but occasional cases persisted for up to 3 years. By contrast however, Trent (1963a,b) who re-evaluated 25 men who had been diagnosed as having filariasis during the Second World War and found that many of them still had evidence of the disease 15 years after leaving the endemic area. Intermittent genital symptoms - pain, swelling and tenderness precipitated or aggravated by standing or physical exercise was present in 32%. Forty percent had abnormal physical findings such as thickening of the spermatic cord, the presence of a tender nodule near the testis, induration of the testis, varicocele, or thickening of the scrotal skin. Recurrent lymphangitis still occurred in 12% with attacks occurring 1 to 3 times per year and lasting for 1 to 3 days. Symptoms consisted of sudden onset of fever, headache, malaise, swelling of affected lymphatics and swelling of regional lymph nodes. In 68% there were episodes of swelling of the extremities and enlargement of the regional lymph nodes draining the area. These episodes tended to occur most frequently in the summer months and the patients complained that the skin of the affected extremities became "puffy, stiff and tight". Winstead (1978) reported a patient who still complained of symptoms 30 years after contracting the disease during military service in Samoa. At the time of his examination (1972) there was no clinical evidence of active filariasis but he had a positive skin test and a weakly positive haemagglutination test. There is no evidence that any of those infected during WW2 developed chronic hydrocele or elephantiasis but in occasional cases microfilaraemia persisted for 15 years (Conn and Greenslit, 1952; Trent, 1963 a, b).

There are several vexing questions to be answered with regard to the American armed forces cases. Why did most of the cases occur in the central Pacific area? Most authors seem to believe that it was because the soldiers were in close contact with infected natives in an area highly endemic for filariasis. Papua New Guinea and the Solomon islands were also highly endemic for filariasis and many American troops were also stationed there. Although some American forces cases did occur in these areas, the numbers seem to be far lower than those that occurred in the Pacific area. Wartman (1947) suggests that the high number of cases in the Central Pacific was due to the fact the vector mosquito is a day time feeder, the microfilariae have diurnal periodicity and since most of the troops activities took place during the day, they were heavily exposed. This does not explain why fewer cases occurred in Papua New Guinea and the Solomon Islands. The vectors there are mainly night feeders but the troops being in an active theatre of war and occupying positions both night and day would be heavily exposed. The other interesting observation is the low number of filariasis cases among Australian troops even although they were stationed in highly endemic areas (Papua new Guinea and the Solomon Islands). Walker (1952) states that there were only 22 cases of filariasis among Australian troops, and as some of these men were Queenslanders (where filariasis was endemic at the time) may have acquired

the infection prior to enlisting in the forces. It may have simply been a lack of familiarity with filariasis and the diagnosis was missed but this was unlikely as the Australian forces medical staff were aware of the risk of filariasis and had been extensively briefed on the American experience in the Pacific (Walker 1952). An interesting observation is that most of the filariasis cases occurred in areas that are not endemic for malaria. Is it possible that malaria somehow modifies the course of filariasis in naive subjects? Although there is no evidence as yet from human studies, this idea merits further study as there is evidence from animal studies that filariasis modifies the course of *P. falciparum* malaria (Yan *et al.*, 1997). Another possibility is that the commonly used anti-malarial prophylactic in use at the time - atabrin, had an anti-filarial action. Although this appears unlikely, it should not be discounted without exploration. Although none of the American servicemen developed elephantiasis, this condition does occur, albeit rarely, in expatriates who have lived in endemic areas for a number of years. One of Patrick Manson's original cases was a French priest from French Polynesia (Manson-Bahr, 1959). Webster reported 15 cases of elephantiasis among white settlers in Samoa. All of these were long-term residents with a mean age of 64.3 years and a mean exposure time of 35.1 years. I, myself discovered an American missionary with severe elephantiasis who had lived on the island of New Britain, Papua New Guinea for over 40 years.

Filariasis in Other Troops

Alhadeff (1955) reported cases of acute filariasis in Mauritian troops stationed in the filarial-endemic areas of Egypt. It difficult to say however whether these were locally acquired infections or not because filariasis also occurs in Mauritius (Sasa 1976).

Filariasis also occurred in expatriate troops serving in Vietnam where both *W. bancrofti* and *B. malayi* occur (Greenberg, 1969; Colwell *et al.*, 1970; Hembree, 1974; Sasa, 1976). In the 1950's a syndrome consisting of hyper-eosinophilia, bronchial asthma and lymphadenopathy was observed in 151 French and North African troops serving in the Tonkin area of North Vietnam. Microfilariae of *B. malayi* were recovered from the lymph nodes of some cases but there are no reports of circulating microfilariae. Treatment with arsenicals or diethylcarbamazine was followed by rapid clinical improvement (Galliard, 1957). Brown and Armstrong (1968) and Sullivan *et al.* (1969) reported some cases of filariasis in American servicemen. By the use of a soluble-antigen fluorescent antibody test Colwell *et al.*, (1970) were able to show that 11% of United States servicemen who were stationed in filarial-endemic regions of South Vietnam had anti-filarial antibodies. By contrast, only 3% of those stationed in apparently non-endemic areas were positive. Moreover, when the servicemen were characterised according to the degree of potential exposure, the prevalence of sero-positivity in field troops was 3 times higher than in base

personnel. One serviceman was also found to have circulating microfilariae. Sullivan *et al.*, (1970) reported a case of clinical filariasis without microfilaraemia that rapidly resolved following DEC therapy. Again, there appears to be reported cases of filariasis among Australia military personnel. Most of Vietnam is malarious and as far as it can be ascertained, American and Australian service persons were on the same malarial prophylaxis regime.

Other Expatriate Cases

McQuay, (1967) studied a group of furloughed missionaries who had served in filarial-endemic areas.

Using haemagglutination and flocculation tests he was able to demonstrate filarial antibodies in 30% of the 1,436 people examined. In some cases microfilariae of *D. perstans*, *L. loa* and *M. ozzardi* were identified but no cases of *W. bancrofti* or *Brugia* species microfilaraemia were found. Bean *et al.*, (1992) reported a case of acute filariasis in an American traveller. The 45-year-old man had spent a short time trekking in Nepal. En route to Nepal he had stayed overnight in Bangkok. Seven months after his return to the United States he developed fatigue, eosinophilia and lymphodema of the arm and chest wall. There was no lymphangitis, lymphadenitis or pain. The symptoms were aggravated by physical exercise. His indirect haemagglutination titre and ELISA assay for filarial antibodies was positive. No circulating microfilariae were detected. Three months after diethylcarbamazine treatment the eosinophilia had fallen and the patient was asymptomatic. During the next 2 years there were several milder episodes followed by complete resolution. Filarial antibodies were present and lymphatic pathology was demonstrated by lymphoscintigraphy. Charters (1981) described a 43-year-old man who had spent 2 years in Papua New Guinea during which he suffered from several bouts of pyrexia of unknown origin. Upon returning to Australia he had several similar episodes accompanied by lymphangitis of the leg. These attacks ceased whilst he was taking

phenoxymethylpenicillin but relapsed within a month of him ceasing the medication. Given that penicillin has not been shown to have any antifilarial action but is usually effective against streptococci, it may be possible that these episodes are related to streptococcal infection which has been shown to be implicated in acute filarial attacks (see below). Libman *et al.*, (1993) screened 1,605 expatriates returning from tropical areas and found that 16 (0.995) had positive filarial serology. Turner and Usurup (1997) report a possible case of filariasis in an expatriate miner working in Papua New Guinea and Moore *et al.*, (1996) reported the case of an American Peace Corps worker who developed acute lymphatic dysfunction (proven by lymphoscintigraphy) within 3 months of arriving in a filarial-endemic area. Koya *et al.*, (1998) found a case of filariasis in an Italian woman who developed acute filariasis after staying for a few weeks in a highly endemic area of India and Manfredi *et al.*, (1998) described a case of *B. malayi* infection in another Italian, a HIV-infected drug user which presented four months after returning to from a two month-long journey to India and Southeast Asia. The interesting thing about these two patients is that they were both microfilaraemic, a rare finding in expatriate cases.

From these case descriptions it is obvious that filariasis is a real threat to expatriates living in, working in, or visiting endemic areas and should always be considered in the differential diagnosis of returning travellers (Doherty *et al.*, 1995). The question of what tests to use in diagnosis is a vexing one and will be discussed in the section on diagnosis below. There is a sharp contrast between expatriates who only visit or work in a filarial-endemic area for short periods of time and expatriates who migrate to filarial endemic areas (or even from one filarial-endemic area to another) permanently. In these people there is an acceleration of the disease progression and they developed chronic pathology such as elephantiasis much faster than the people who are the native inhabitants of the area (Partono *et al.*, 1978).

Filarial immunity

Cellular Immunity

All parasites induce a degree of immuno-suppression as a means of ensuring their survival in the host (Nussenzweig, 1982). Loss of T lymphocyte proliferation and the emergence of a host response dominated by a Th-2- type profile is characteristic of many nematode infections, especially filarioids (Allen and Macdonald, 1998). The immunology of lymphatic filariasis has been very aptly termed an "immunologic maze" (Piessens, 1981) and all that can be attempted here is a summary of the well-established principles. For comprehensive reviews see: Grove, 1978, 1979; Kwa and Mak, 1980; Ottesen, 1980; Ottesen, 1984; Mistry and Subrahmanyam, 1985; Dasgupta *et al.*, 1987; Kumar *et al.*, 1987; Prasad and Harinath, 1987; Ottesen, 1989; Ottesen, 1992; Maizels, 1995; Freedman, 1998.

In general terms, immunological profile differs between people with microfilaraemia, chronic filarial pathology and tropical eosinophilia syndrome (Ottesen, 1992). Microfilaraemics have reduced cellular immunity to filarial antigens and their lymphocytes stimulation index in lymphocyte proliferation tests is usually below 5.

Weller (1978) followed *in-vitro* lymphocyte blastogenesis in rats infected with *B. pahangi*. Responses to infective larva antigens were detectable only in the first few weeks of infection whereas responses to microfilariae antigens only developed as microfilaraemia waned. Cyclic variation in the response to adult worm antigens occurred throughout the period of infection. Work by Deehan *et al.*, (1998) suggests that a phosphocholine-containing filarial secretion inhibits lymphocyte activation by targeting key lymphocyte proliferative signalling pathways. Allen and Macdonald (1998) found that excretory antigen produced from *Brugia malayi* transplanted into mice induced the formation of adherent peritoneal cells that block T cell proliferation. It has been demonstrated by Pancre *et al.*, (1988) that platelets from filariasis patients are cytotoxic to microfilariae and that this effect is inhibited by supernatants from filarial antigen-stimulated lymphocyte cultures - another immune system-modulated protective effect to enhance the survival of the parasite in the host.

It is a common belief that filarial immuno-suppression is limited to filarial antigens and that response to unrelated parasite antigens and such compounds as purified protein derivative (PPD) is unaffected (Ottesen *et al.*, 1977; Piessens *et al.*, 1980a-d; Haque *et al.*, 1983; Regunathan *et al.*, 1997). Some findings challenge this view. Mistry and Subrahmanyam (1985) found that the lymphocyte proliferative response to PPD was marginally reduced in microfilaraemics but normal in patients with elephantiasis and (Srivastava *et al.*, 1981) found that T

cell responses to PPD were normalised after specific antifilarial therapy.

Reports on mitogen response vary. Portaro *et al.*, (1976) Lammie and Katz, 1983; Mistry and Subrahmanyam (1985) and Prasad *et al.*, (1991) report reduced response to mitogen whereas Regunthan *et al.*, 1997 found no difference in mitogen response between groups with microfilaraemia or chronic filarial pathology and endemic normals from the same area. This variation in mitogen response may be due to the timing of the experiments and a variation in responses between subjects. Using a rat model, Weller (1978) demonstrated that the response to phytohaemagglutinin, pokeweed mitogen, and bacterial lipopolysaccharide varied throughout a 1.5-year period of infection with *B. malayi*. Lymphocytes from some animals showed a complete absence of mitogen response between the 16th and 36th week of infection and normal responses at other times.

Other animals however, showed no mitogen-related immuno-suppression throughout the period of infection.

People with elephantiasis usually have normal or increased cellular immunity to filarial antigens with stimulation indexes between 5 and 25 (Ottesen *et al.*, 1977; Weller 1978; Piessens *et al.*, 1980 a, b; Narayanan *et al.*, 1986; Nutman *et al.*, 1987a; Ottesen, 1989; Hitch *et al.*, 1991; Ottesen, 1992; Addis *et al.*, 1995). There is a sub-set of elephantiasis patients, however, which demonstrate a hypo-responsiveness to filarial antigens comparable to that seen in microfilaraemics (Yazdanbakhsh *et al.*, 1993a, b). It was demonstrated that circulating phosphocholine-containing antigens from the adult worm, which can be present in levels as high as 5ug per ml, are capable of suppressing mitogen-induced T cell proliferation (Wadee *et al.*, 1987; Lal *et al.*, 1990). Mahanty *et al.*, (1996a, b, 1997) have found evidence that IL-10 induced by parasite antigen plays an important role in down-regulating antigen-specific lymphocyte proliferation in microfilaraemics and they IL-12 can overcome this down-regulation but does so in a non-antigen-specific manner.

The suppression in cellular immunity is not due to the inability of lymphocytes to respond to antigenic stimulation, but to the presence of suppressor cells (Piessens *et al.*, 1982). It has been shown that the degree of immuno-suppression and rapidity of onset is related to the size of the microfilariae burden and that a reduction in microfilariae decreases the immuno-suppressive effect (Piessens *et al.*, 1981b). This concept however has been challenged by Bosshardt *et al.*, (1995) who found using a Gerbil model, that the microfilaria stage alone is not responsible for the maintenance of cellular immuno-suppression and that the degree of immuno-suppression is related to the total filaria parasite burden. It is interesting to note that Riley *et al.*, (1989) found that the *in-vitro*

immuno-suppression induced by malaria can be partially reversed by indomethacin. This effect is thought to be due to the ability of indomethacin, a cyclo-oxidase inhibitor, to block the production of prostaglandin, which are potent inhibitors of blastogenesis. It would be interesting to carry out a similar experiment using mononuclear cells from filariasis patients and filarial antigens.

Filarial Immunity: Lymphocyte Sub-population Profiles

Dasgupta *et al.*, (1986) found that the total number of circulating lymphocytes in elephantiasis patients was lower than normal due to a depletion of T cells but the percentage of null cells and T rosettes was higher in elephantiasis and microfilaraemics than in amicrofilaraemics. Piessens *et al.*, 1982 found that increased numbers of suppressor T cells were present in 15 of 17 patients with microfilaraemia and in 6 of 11 patients with elephantiasis. The increase correlated with the hypo-reactivity to filarial antigens and the removal of suppressor T cells improved the reactivity to filarial antigens. King *et al.*, (1992) demonstrated that the frequency of parasite-specific CD3+ T cells and the proportion of lymphocytes producing parasite-specific IgE or IgG was significantly lower in microfilaraemics than in patients with chronic pathology.

Cytokines

Cytokine profiles also differ in various manifestations of filarial disease. In microfilaraemic humans and animals the Th1 arm of the immune response is down regulated and the Th2 arm stimulated. Microfilaraemics produce large amounts of IL-4 and IL-5 (type 2 cytokines) and only small amounts of interferon γ and GM-CSF (type 1 cytokines). The situation is reversed in elephantiasis (Nutman *et al.*, 1987b; Maizels and Lawence, 1991; Ottesen, 1992; Pearlman, 1995; Yazdanbakhsh *et al.*, 1993a,b; Dimock *et al.*, 1996; Mahanty *et al.*, 1996; 1997). The cytokine profile has shown to be influenced by clinical status and age of the patient (de Almeida *et al.*, 1996; Sartono *et al.*, 1997). Interferon γ and IL-5 are suppressed in microfilaraemic carriers but IL-4 was not affected. In microfilaraemics IL-4 production was high in young subjects but decreased with age whereas in microfilaraemics IL-4 increased with age. By contrast, interferon-gamma increases with age in amicrofilaraemics but not in microfilaraemics. IL-5 on the other hand increased with age in both groups (Sartono *et al.*, 1997).

It has also been shown in animal studies that antibodies to inhibitory cytokines have the ability to reverse temporally reverse cellular unresponsiveness and induce a transient increase in microfilariae (Ottesen, 1992. In a human study however by Regunathan *et al.*, (1997) the cellular unresponsiveness was not reversed by the addition of

recombinant IL-1a, IL-1b and interferon γ but it was reversed by the addition of serum from endemic normals.

Some studies however have shown that the correlation between Th and cytokine profiles and the manifestation of filarial disease may not be clear cut and that the presence or absence of infection rather than clinical status is most closely associated with cytokine response (Almeida *et al.*, 1996). Pearlman *et al.*, (1995) have shown that T helper cell responses to filarial antigens are modulated by IL-12. Cytokine profiles are also influenced by age (Sartono *et al.*, 1997). Antigen-specific IL-5 and interferon γ are suppressed in microfilaraemics but IL-4 is normal. In asymptomatic amicrofilaraemic cases IL-4 production was increased in young people but decreased with age, whereas in microfilaraemics IL-4 production showed a significant increase with age. By contrast, interferon γ showed an increase with age in asymptomatic amicrofilaraemics but not in microfilaraemics. IL-5 decreased significantly with age in both asymptomatic microfilaraemic and microfilaraemic groups. This indicates that the cytokine profile change with the length of exposure to, and infection with filarial parasites. Mahanty *et al.*, (1996, 1997) have shown that parasite antigen-induced IL-10 plays an important role in down regulating cellular immune responses in microfilaraemic patients.

CD (Cluster of Differentiation) Cells and Molecules

Studies by Yazdanbakhsh *et al.*, (1993) of soluble CD25 (sCD25) and soluble CD27 (sCD27) showed that levels of sCD27 were significantly increased in microfilaraemics and elephantiasis patients compared with endemic normals. By contrast, sCD25 remained low in microfilaraemics and was only slightly raised in elephantiasis.

Cell Adhesion Molecules

The development of chronic inflammation depends upon the migration of peripheral blood mononuclear cells into tissues. Animal models show that this process is facilitated by vascular cell adhesion molecule (VCAM-1) expression on vascular endothelium (Freedman, 1996). *In-vitro* studies by Freedman *et al.*, (1996) and Plier *et al.*, (1997) that cell culture supernatants from filariasis patients stimulated the migration of mononuclear cells and the effects were greater when cell cultures from individuals with clinical filariasis, as opposed to asymptomatic individuals was used. These studies were conducted using the "traditional" classification of disease status and the findings would need to be revised if the proposed new classification of Freedman (1998) is used (see below).

C-reactive Protein and Filariasis

Both serum and lymphocyte-bound C-reactive protein has shown to be significantly increased in patients with filarial pathology compared with asymptomatic microfilaraemics and endemic normals (Lal *et al.*, 1991).

Attempts to Reverse *in-vitro* Immuno-suppression

In an attempt to reverse the *in-vitro* immuno-suppression against filarial antigens Sartono *et al.*, (1995a,b) cultured peripheral blood mononuclear cells in the presence of agents which had been shown to enhance T cell proliferation in other diseases: interleukin 2 (IL-2), anti-interleukin 4 (IL-4), interleukin 7 (IL-7), anti-interleukin 10, anti-CD2, anti-CD27, anti-CD28, indomethacin, phorbol myristate acetate (PMA) and calcium ionophore. Only IL-2, IL-7, indomethacin and PMA produced a result and then only in some of the cases studied. Unresponsiveness in elephantiasis cases was easier to reverse (50% of cases responded) than microfilaraemics (12.5%) and asymptomatic amicrofilaraemics (20%). Their conclusion was that more than one immunological mechanism was involved in filarial immune suppression.

Humoral Immunity

All filarial infections produce an active polyclonal humoral response with the production of IgG, IgM and IgE antibodies (Ottesen, 1984; Kumar *et al.*, 1987; Nutman *et al.*, 1987a,b; Nanduri and Kazura, 1989) but their titres tend to be lower in people with microfilaraemia than in those with elephantiasis and extremely high in cases of tropical eosinophilia syndrome (Wong and Guest, 1969; Ottesen *et al.*, 1982; Ottesen, 1989; Addis *et al.*, 1995; Ottesen, 1992; Estambale *et al.*, 1995). Antibodies against microfilariae are usually absent in microfilaraemia and lymphocytes isolated from microfilaraemics are unable to produce filaria-specific antibodies *in-vitro* either spontaneously or in response to parasite antigen. This is in contrast to patients with chronic obstructive pathology or TPE (a,b). The failure in antibody production cannot be reversed by removal of adherent cells or T⁸ lymphocytes suggesting that a state of B unresponsiveness to filarial parasites is present (Nutman *et al.*, 1987a,b).

IgG isotype profile also differs in different manifestations of disease. In microfilaraemics, a large proportion of the IgG consists of IgG4 (up to 90%) and there are low levels of IgG1, IgG2 and IgG3. By contrast, patients with elephantiasis have lower titres of IgG4 antifilarial antibody than microfilaraemics and often have increased levels of IgG2 and IgG3 (Ottesen *et al.*, 1985; Hussain *et al.*, 1987; Kwan-Lim, 1990; Ottesen, 1992; Egwang *et al.*, 1993; Kurniawan *et al.*,

1993; Lammie *et al.*, 1993; Yazdanbakhsh *et al.*, 1993a,b; Rahmah *et al.*, 1994; Addiss *et al.*, 1995; Maizels *et al.*, 1995; Wamae *et al.*, 1995; Yazdanbakhsh *et al.*, 1993a,b; Atmadja *et al.*, 1996; Simonsen *et al.*, 1996; Nicolas *et al.*, 1999). Yazdanbakhsh *et al.*, 1995a have shown that in elephantiasis the IgG3 antibody has a particular affinity with the recombinant antigen Bpa-26, a peptide from the C-terminal region of the filarial heat-shock protein and show a striking association between the elevated levels of IgG3 antibody and chronic obstructive filarial pathology. They postulate that an imbalance of isotypes of antibodies to particular filarial antigens may play a part in the pathogenesis of chronic disease. Serum antifilarial IgE is lower in asymptomatic microfilaraemics than in individuals with chronic pathology (Marley *et al.*, 1996) and the ratio of IgG4 to IgE is highest in asymptomatic microfilaraemics (Atmadja *et al.*, 1995). Estamble *et al.*, (1994) showed that the mean level of filarial-specific IgG1 was significantly higher in amicrofilaraemic, symptomatic cases than in microfilaraemic cases whereas the mean level of IgG4 was significantly higher in individuals with microfilaraemia. They also found that in most categories of endemic individuals, and for most antibody isotypes, the mean levels of antibodies tended to be higher in younger than in older age groups. It is likely that the elevation of IgG2 and IgG3 contribute to the pathology of elephantiasis by promoting cellular hypersensitivity reactions or immune complex formation (Maizels *et al.*, 1995). IgG3 is a major mediator of type 3 hypersensitivity and is the IgG isotype for which the Fc receptors on mononuclear cells and granulocytes have the highest affinity (van der Winkel and Capel, 1993; Maizels *et al.*, 1995). Immune complexes formed with IgG1, IgG2 and IgG3 isotype antibodies and filarial antigens are large molecules that can become fixed in tissues. By contrast, IgG4 is monovalent and in lymphatic filariasis targets the lower molecular weight antigens. As a result, forms small immune complexes which do not become fixed in tissue and are less damaging than those formed by IgG1, 2, and 3 (Kazura, 1993a). Studies by Prasad *et al.*, (1983) have shown that patients with chronic filarial pathology have significantly higher levels of immune complexes than microfilaraemics or endemic normals. This finding from this study however is at variance from those of Kobayashi *et al.*, (1997) who found that asymptomatic microfilaraemics have high-level filarial antigen-associated immune complexes that correlate with the level of the microfilaraemia. Lutsch *et al.*, (1988) showed that the antibody complexed to antigen is mainly IgM in asymptomatic microfilaraemics and IgG in symptomatic cases.

IgG4 antibody plays an important part in the aetiology of filariasis. Patients with chronic helminth infection, despite being exposed to parasite antigen seldom exhibit any allergic reactions because IgG4 antibodies "block" the action of IgE. In microfilaraemics the ratio of IgG4 to IgE is high (Maizels *et al.*, 1995, effectively

dampening down the allergic response in the host (Hussain *et al.*, 1992), thus ensuring the survival of microfilariae and preserving the potential to transmit the disease (Nanduri and Kazura, 1989). Mahanty *et al.*, (1994) have shown that antifilarial IgG4 is a good index of the intensity and duration of filarial exposure in endemic populations and the level of IgG4 antibody correlate with microfilariae counts (Maizels *et al.*, 1995). This finding however is not true for all filarial species - Akue *et al.*, (1994) found that in *Loa loa* infections there are high levels of parasite-specific IgG4 even in the absence of microfilaraemia. Simonsen *et al.*, (1996) showed that IgG1 and IgG4 antibodies have high specificity for filarial antigens and that their titres have a relationship to micro filarial status. The differences in IgG isotype profile in groups with different manifestations of filariasis have been consistently demonstrated in a number of countries including: Malaysia (Rahmah *et al.*, 1994), Papua New Guinea (Kwan-Lim, 1990), Gabon (Egwang, 1993), India (Ottesen *et al.*, 1985; Hussain *et al.*, 1987), Haiti (Lammie *et al.*, 1993), Tanzania (Simonsen *et al.*, 1996 and the South Pacific Islands (Chanteau *et al.*, 1991; 1995) but there is one report from Haiti where 10% of a group of microfilaraemics did not have an increase in antifilarial IgG4 antibody.

These persons had very high levels of filarial antigen and it is suggested that in this situation there may be a down regulation of antibody production, which extends to IgG4 (Marley *et al.*, 1995). Most studies of filarial antibodies have utilised crude antigens prepared from adult *Brugia malayi* or *Brugia pahangi*. There have been some studies using antigens prepared from microfilariae and it has been shown that there is an inverse relationship between the anti-micro filarial antibody titre and the level of microfilaraemia leading to the hypothesis that these antibodies control the number of circulating microfilariae (Piessens *et al.*, 1980b) and there is a correlation between the presence of anti-microfilarial sheath antibodies in serum and the absence of detectable circulating microfilariae (McGreevy *et al.*, 1980; Piessens *et al.*, 1980b). It is believed that the microfilariae contribute to the decline of circulating antibody because antibody is bound onto antigen on the surface of the microfilaria (Piessens *et al.*, 1980b).

Infective (L3) larvae elicit a IgG and IgE response (Das *et al.*, 1992) and Day *et al.*, (1991) have shown that the level of anti L3 antibody correlates with the duration of exposure. Kurniawan-Atmadja *et al.*, (1998) however, found that unlike the response to adult or microfilarial antigens, the immune response to L3 surface antigens dominated by IgG1 and IgM rather than IgG4.

Kar *et al.*, 1993 report some interesting changes in the humoral immune response during filarial fever attacks.

Titres of IgG and IgG4 antibody to *W. bancrofti* micro filarial excretory/secretory antigens decreased significantly during fever attacks and remained reduced for up to one month after the fever episode. By contrast, the level of circulating immune complexes rose sharply during the fever episode and in a

percentage cases there was also an increase in circulating filarial antigen suggesting a release of filarial antigen into the circulation during fever which bound to antibody and form immune complexes. Total and differential white cell counts and antistreptolysin O titres did not change during or after febrile episodes strongly suggesting that streptococcal infections do not play a part in filarial fevers. It is therefore postulated (Kar *et al.*, 1993) that antigens released from adult worms into the circulation during parturition may evoke an allergic reaction in the host and triggering a febrile episode.

IgM antibodies: Estambale *et al.*, (1994) found that the levels of IgM filarial antibodies rose in individuals from both filarial-endemic and filarial-non-endemic areas. They believed that this was due to the low specificity of IgM antibodies and the subsequent cross-reaction with other parasite and non-parasite antigens.

Filarial Immunity in the BALB/c Mouse Model

In this model, IgG isotype profiles, IgE production and cytokine profiles have been shown to be stage specific (Lawence *et al.*, 1994). Microfilariae induced production of all IgG isotypes but little IgE. By contrast, adult worms stimulated the production of only IgG1 and IgE. When splenocytes from parasite-implanted mice were stimulated *in-vitro* with parasite antigen or con-A cells from microfilariae-infected mice secreted high levels of interferon- γ but IL-4 production did not peak until 28 days after parasite implantation. On the other hand, adult female worms stimulated high levels of IL-4 and very little interferon γ Male worms stimulated lower levels of IL-4 production than female worms. CD4 lymphocytes were primarily responsible for the IL-4 increase. To quote Lawence *et al.*, (1994):

"These findings demonstrate that adult filarial parasites, and female parasites in particular, exert a rapid polarisation of the immune response in a TH2-like direction, but this effect may be modulated by the microfilaria stage."

The Effect of Filariasis on the Humoral Response to Other Antigens

Whilst it is commonly believed that cellular immunosuppression induced by filariasis is filarial antigen specific (see above), there is evidence that antibody response and delayed skin hypersensitivity to non-filarial antigens is reduced. Grove (1979) found that only 1 out of 35 patients with filariasis responded to tetanus toxoid (3%) compared with 8 out of 31 control subjects (26%)

($p < 0.025$). Cooper *et al.*, (1998) studied the response

of oncocerca-infected patients to tetanus toxoid and found that six months after vaccination, antibody levels lymphocyte proliferative responses and levels of interferon γ were significantly higher in non-infected persons when compared to oncocerca patients. Infected subjects produced IL-10 but un-infected controls did not suggesting a role for IL-2 in this process.

Grove (1979) found that the response to *Salmonella typhi* vaccine is effected with only 15 of 26 patients with filariasis (58%) having an antibody titre of 1:40 2 weeks after immunization compared to 25 of 31 in controls (81%) ($p < 0.05$). Delayed skin hypersensitivity reactions to *Candida*, mumps or streptococcal antigens were significantly reduced in filariasis patients compared with controls ($p < 0.01$). Srivastava *et al.*, (1981) showed that filariasis patients have a significantly impaired antibody response to the H and O antigens of TAB vaccine and reduced skin hypersensitivity to PPD and chlorodinitrobenzene. This immuno-suppression may contribute to secondary cutaneous infections (Grove, 1979). Streptococcal infection has been strongly implicated in the clinical manifestation of filarial disease and the question could be asked, "Does filariasis suppress immunity to streptococci." Lal *et al.*, (1991) however demonstrated *increased* rather than *decreased* streptococcal antibody levels in patients with filariasis and King *et al.*, (1992) found no difference in streptolysin O-induced lymphocyte proliferation between filariasis patients with various manifestations of disease and normal controls.

Srivastava *et al.*, (1981) demonstrated a reduced antibody response to TAB vaccine in filariasis. Antibody titres to *S. typhi* H and O antigens were significantly reduced in asymptomatic microfilaraemics and patients with chronic filaria pathology when compared with normal controls.

Malhotra *et al.*, (1997; 1999) have shown that 2 to 10 year old children born to mothers who did not have filariasis produced a 10-fold increase the production of interferon γ when challenged with PPD than those children born from filarial-infected mothers. This raises two serious questions: is the efficacy of BCG vaccine impaired by filarial infection and does filarial infection influence the course of TB infection? The effect of filarial immuno suppression on the response to other vaccines should be investigated. If filariasis does reduce the efficacy of vaccination it is another reason why the control of filariasis is important.

Does Filarial Immunosuppression Impact Upon Other Tropical Diseases?

It is also possible that filariasis may decrease the

immunity to other tropical diseases and that hypothesis should be actively explored using modern immunological techniques. Sidkey *et al.*, (1987) reported a patient with filariasis who presented with a disseminated *Hymenolepis nana* infection and they postulated that the filarial immuno-suppression may have contributed to the spread of the other parasite. The situation is somewhat clouded however by the fact that the patients had been treated in the past with corticosteroids. Younis *et al.*, 1997 studied the relationship between filariasis and intestinal parasites in a filarial-endemic area of Egypt. The overall prevalence of parasites was 81% in non-filarial-infected people and 92.4% in filariasis patients. On the basis of this data they concluded they suggest that the immune suppression caused by filariasis may increase the host's susceptibility to other parasitic infections. Mohamed *et al.*, (1983) could find no evidence of interaction between filariasis and schistosomiasis in an area endemic for both parasites. This is an area that warrants further study.

Piessens *et al.*, (1983) have shown that malaria and filariasis have opposing effects on human immuno regulatory T lymphocytes and a study by Schmidt and Esslinger (1981) found that *P. falciparum* infections in Owl monkeys (*Aotus tri-virgatus griseimembra*) infected with the filarial parasite *Tetratetalonema barbascalensis* followed a more benign course than in monkeys who were not infected with the filaroid. Yan *et al.*, (1997) showed a down-regulation of murine susceptibility to cerebral malaria by inoculation with third-stage larva of *B. pahangi*. There is no evidence at this time however that filariasis influences the course of human malarial infections. Ravindran *et al.*, (1998) studied patients with both filariasis and malaria and found that there was no evidence to suggest that either of these two diseases impacted upon the other. This subject does warrant further investigation however, as it would not be helpful if the eradication of filariasis led to an upsurge in malaria!

Work by Fiennes (1969) suggested that filarial infection might aid in the pathogenic activation of latent viruses in new world non-human primates. In a Japanese study by Tajima *et al.*, (1983) it was found that the positive rate of antibodies to HTLV-1 virus among people with a high antibody titre to filariasis antigen was high than in those with a low filarial antibody titre and postulated that filariasis may promote the effects of HTLV-1 infection but no correlation could be found by Linhares *et al.*, (1995) in Brazil.

Another important question could be asked in the light of the experimental evidence above. Does filarial infection in children reduce the efficacy of vaccination?


Unravelling the Pathogenesis of Chronic Pathology

Although much progress has been made many questions are yet to be resolved. What causes one individual to progress from asymptomatic microfilaraemia to elephantiasis yet his next door neighbour, or even their spouse who shares the same house (Hunter, 1992) remains asymptomatic or even avoid the disease altogether? What factors cause the "re-awakening" of the suppressed immunity? How is it that microfilaraemia and chronic pathology can co-exist in some people? Why is it that some communities have exceedingly high prevalence of filarial antigenaemia and microfilaraemia and very little chronic pathology? Is it a just a matter of duration and intensity of exposure? Would everyone who has microfilaraemia progress to chronic pathology if the time frame is long enough? Piessens and Partono (1980) hypothesise that the *chances* of developing clinical manifestations of filarial disease increase with increasing worm burdens and the *differences* in the clinical course of filariasis are due to

the individual's immune response to filarial antigens. There is however evidence from animal models which suggests that individuals which are subjected to repeated infections with L3 infective larvae are especially at risk for developing chronic pathology. Repeated infection of cats and ferrets with infective larvae produced increased oedema (Rogers and Denham, 1975; Hines *et al.*, 1985; Maizels and Lawence, 1991) and lymphatic damage is more severe in cats after repeated infection. Lymphatic pathology is more severe in cats that have the immunological capacity to kill adult worms and microfilariae than it cats who are microfilaraemic (Maizels and Lawence, 1991).

A good working model is that proposed by Maizels and Lawence (1991). It is based on the idea that microfilaraemia is dependent upon immune tolerance to filarial antigens developing in the host and that chronic pathology arises when the immune tolerance breaks down (Figure 6).

Figure 13 Maizels and Lawence (1991) working tolerance model (with permission of the authors)



In accepting that model however it needs to be recognised that filariasis is an extremely complex disease and the old classification which separated patients in the basis of microfilaraemia and the presence or absence of chronic pathology have been

seriously challenged in recent years (Michael *et al.*, 1994; Freedman, 1998). What Ottesen (1992) likes to call "overlap" syndromes do exist and other mechanisms besides immunological ones may play a part in the induction of pathology.

More than One Pathway to Chronic Pathology?

In an immuno deficient mouse (nude mouse) model it has been shown that there are pathological mechanisms that lead to lymphodema and elephantiasis. If these immuno deficient mice are infected with *Brugia* the parasite itself will, even in the absence of an immunological reaction in the host, induce lymphatic damage and ultimately elephantiasis.

If infected nude mice are rendered immuno-competent by transplantation of cells from an animal previously sensitised to filarial antigens, inflammatory reactions around the parasites again lead to lymphatic damage and elephantiasis (Suswillo *et al.*, 1981; Vincent *et al.*, 1984; Vickery *et al.*, 1991). Here two different pathways to chronic filarial pathology are demonstrated; one dependent on the parasite alone and one dependant on the reaction of the host. Whether or not a comparable situation occurs in humans remains to be seen, but there is histological evidence that supports the hypothesis. Tissues removed from filariasis cases showed two different patterns of pathological response. Around living parasites the process was that of lymphatic dilation, around dead parasites there was a granulomatous inflammatory reaction (Von Lichtenburg, 1987). Patients with "overlap syndromes" could be explained on the basis that pockets of local immuno-suppression could be developed around adult worms even if the inflammatory process was occurring elsewhere.

Genetic Influences

It is possible that there are genetic variations in antigen processing and immunologic response and there have been a number of attempts to find linkages between genetics and filariasis. Any evidence of such a linkage so far has been weak and sometimes contradictory. Ottesen *et al.*, (1981) noted a family clustering of microfilaraemia but was unable to demonstrate a linkage with human leucocyte antigen (HA) but an association between elephantiasis and HA-B1 has been noted in India and Sri Lanka (Chan *et al.*, 1984; Dasgupta, 1985) and filarial antigens have been shown to be processed with HA-DR (Nutman *et al.*, 1984). Mohamed *et al.*, (1987, 1992) found a significant association between HLA-B7, the susceptibility of retaining circulating microfilariae and the development of clinical filariasis. A study by Yazdanbakhsh *et al.*, (1995b) suggested that HLA class II genes may influence the course of Brugian filariasis by influencing the T-cell dependent antibody repertoire but a later study by the same authors concluded that this evidence is very weak (Yazdanbakhsh *et al.*, 1997). Studies of ABO group linkages have shown varying results (Gyorkos *et al.*, 1983; Srividya and Pani, 1993). Ayers *et al.*, (1976) found a higher incidence of group B in filarial patients, a lack of group A₂ and a higher frequency of non-secretors, Suresh *et al.*, (1982) an increase in A and AB groups and Srikumari Srisailapathy *et al.*, (1990)

showed excess of group B and a deficiency of group AB in filariasis patients compared to controls. This difference however was only statistically significant in males. A positive association between filarial hydrocele and group B (Rh) negative was also noted by Dutta and Diesfeld (1987). Srividya and Pani (1993) tested 1,444 people in south India and found no association between ABO blood groups and filariasis.

Mohamed *et al.*, (1987, 1989) could find no statistical correlation between filariasis and the ABO, Rh and MN blood group systems in 520 Egyptians. There was no association between haemoglobin variants and filariasis in the only published study found to date (Ayers *et al.*, 1976). Because of its rarity, its geographical distribution and distinctive clinical and laboratory features tropical pulmonary eosinophilia would be a likely candidate for a genetic basis but nothing has been found so far (Ottesen, 1992).

The Influence of Maternal Sensitisation

Prenatal experience with filarial antigen determines the subsequent character of the immune response to post natal infection (Ottesen, 1992) and it is believed that neonates exposed to parasite antigens in utero may develop altered foetal immune responses that influence the development of disease in later life (Malhotra *et al.*, 1997). It is now generally accepted that *in vivo* exposure of the foetus to filarial antigens and the development of immune tolerance plays an important role in determining infection status in later life. Animal studies have shown that the offspring of an amicrofilaraemic mother are more likely to develop filarial pathology than the off spring of microfilaraemic mothers (Schrater *et al.*, 1983; Klei *et al.*, 1986; Krishnarao *et al.*, 1989; Miller *et al.*, 1990).

In humans, a child born to (or tolerized by) a microfilaraemic mother is more likely to develop microfilaraemia than a child born to an amicrofilaraemic mother and children born of microfilaraemic mothers are less capable of mounting an immune response to filarial antigens than those child born of amicrofilaraemic mothers. This effect may persist into adulthood (Lammie *et al.*, 1991; Hightower *et al.*, 1993; Steel *et al.*, 1994; Roberts and Janovy 1996). There are probably several mechanisms by which the foetal exposure to filarial antigens occur.

There is evidence of transplacental passage of microfilariae in both animals (Mantovani, 1966; Sucharit and Rongsriyam, 1980) and humans (Neves and Scaff, 1952, 1954; Raghavan, 1958; Bloomfield, 1978; Ramalingeswararao *et al.*, 1984; Pires, 1991; Campbello *et al.*, 1993; Eberhard, 1993) and microfilariae are occasionally seen in the neonatal circulation Bloomfield *et al.*, (1978). Animal studies have shown that transplacental transfer of microfilariae induces immune tolerance in the resulting offspring (Haque *et al.*, 1982, 1988). Antifilarial IgM antibodies

have been detected in cord blood (Dissanayake and de Silva, 1980) and the level of antibody was found to be higher in cord blood from infants of amicrofilaraemic mothers compared to microfilaraemic mothers (Agarwal *et al.*, 1986). Weil *et al.*, (1983) have demonstrated antifilarial IgE in the cord blood of microfilaraemic mothers. IgE does not cross the placenta (Carrier and Truyens, 1995) and there was a quantitative difference between the level of cord blood and maternal antibody which suggests that it was derived from the foetus rather than being the result of the transfer of maternal blood at birth. Not all studies however support the idea of that maternal infection influences subsequent microfilaraemia. Hitch *et al.*, 1997 found that cord blood mononuclear cells showed very little evidence of *in-vivo* sensitisation to filarial antigens. Filaria-specific IgM levels in half the infants were no greater than the background level of pooled cord blood samples from a non-endemic area and the anti-filarial IgM levels did not correlate with the filarial antigen state of the mothers. Seventy one percent of the cord blood samples showed no anti-filarial IgE and there was no correlation between cord blood anti-filarial IgE and maternal filarial infection status. Only 14% of the cord blood mononuclear cells tested responded to filarial antigens in utero. The oral route via breast milk is another possible mechanism for sensitisation in infancy. *Onchocerca volvulus* antigen has been found in breast milk (Petalanda *et al.*, 1988) but there have been no comparable studies in lymphatic filariasis.

Support for the role of maternal sensitisation in influencing the course of filariasis also comes from studies of filariasis in expatriates. As discussed above, exposure to filarial antigens in expatriates produces an inflammatory reaction which is often more severe than that seen in long-term residents of the same endemic area (Ottesen, 1992) and Indonesian studies have shown that newly settled immigrants from an area where filariasis endemicity was low have a greater chance of developing elephantiasis when they re-settle in a highly endemic area than those who had lived there all their lives (Partono *et al.*, 1978). Although the evidence for *in-utero* conditioning as a risk factor for microfilaraemia appears strong, there is also evidence that the intensity of local transmission can influence the outcome. Alexander *et al.*, have shown that local transmission intensity is a confounding variable in determining the propensity of children to develop a patent filarial infection.

The Role of Super-added Bacterial and Fungal Infections

Lymph stasis provides an ideal medium for the growth of microorganisms and skin lesions allow bacteria and fungi to invade and set up chronic infection

(Olszewski *et al.*, 1993, 1994, 1997, 1999; Olszewski and Jamal, 1994; Burri *et al.*, 1996; Roberts and Janovy, 1996). The link between filariasis and bacterial infection, especially streptococci, has been established for a long time (Grace *et al.*, 1932). While some investigators have recovered streptococci directly from patients with acute filarial disease (Grace, 1932, 1934, 1943; Castellani, 1969) others have failed to do so (McKinley, 1931; O'Connor, 1932). Vincent *et al.*, (1998) found strong evidence that the streptococcal disease erysipelas is a significant factor in triggering or amplifying lymphodema or elephantiasis. Anti streptolysin O titres have suggested a relationship between streptococci and filarial pathology in both human (Otero and Lebron, 1936; Rose *et al.*, 1945; Liu *et al.*, 1964; Suma *et al.*, 1997) and animal studies. Bosworth and Ewert (1977) demonstrated that streptococcal infections in cats experimentally infected with *B. malayi* contribute to causation of elephantiasis and developing worms were adversely affected by streptococcal infection of the worm-infected lymphatics. It has been shown that delayed skin hypersensitivity to streptococcal antigens is reduced in filariasis (Grove, 1979) and it would seem that the application of modern methods of immunological research to the interaction of filarial parasites and streptococci might be an area for fruitful research.

It has been shown that careful attention to hygiene, the use of antibiotics and anti-fungals, and the use of physiotherapy and pressure bandages, will modify disease progression and lead to resolution in some cases (Olszewski *et al.*, 1993; Ottesen, 1994; Addiss and Louis-Charles, 1997; Shenoy *et al.*, 1998a, 1999). National inhabitants of the Finschhafen district of Papua New Guinea use the bark of the Lawang tree (*Cinnamomum culilawan*), which contains the essential oils eugenol and safrole, as a poultice to relieve the pain of acute adeno-lymphangitis (McMillan, 1968) and such local treatments warrant more detailed investigation. A fitting closure to this section would be to quote the words of Ottesen (1994).

"No clear cut evidence exists to demonstrate an inevitable progression from one clinical form of lymphatic filariasis to another. Instead there appears to be a definite element of plasticity in the response to the parasite the reflects the balance among a variety of factors including prior sensitisation to parasite antigens and present influence of parasite- and host-derived immuno-modulating molecules, as well as the current nature of exposure to infection by the parasite."

Diagnosis of Lymphatic Filariasis

Lymphatic filariasis may be diagnosed by:

1. Demonstration of microfilariae. This can be subdivided into:
 - Direct techniques, and
 - Concentration techniques
2. Detection of filarial antigens
3. Detection of filarial antibodies
4. Detection of parasite DNA by DNA probes or the polymerase chain reaction (PCR)
5. Detection of adult worms
6. Skin tests with filarial antigen

(WHO, 1984, 1987, 1992, 1993, 1994; Eberhard and Lammie, 1991a; Wame, 1994).

Methods for Direct Detection of Microfilariae

In areas where microfilariae exhibit nocturnal periodicity blood should be taken within two hours either side of midnight when the highest density of microfilariae is expected to occur. Where they show diurnal periodicity blood should be taken two hours either side of midday. Simonsen *et al.*, (1997a) have devised a method to adjust for the effect of sampling time on microfilariae density. Using their technique it is possible to predict what the microfilariae density would be at midnight in blood collected at say, 2200 hours. The simplest technique for direct diagnosis of microfilaraemia is a giemsa-stained thick blood film of capillary blood collected by finger-prick (Khamboonruang *et al.*, 1987; Schultz, 1988; Schuurkamp *et al.*, 1990; Sabry, 1992. If a measured amount of blood is used (say 40 to 60 μl) the number of microfilariae per ml can be calculated (Mouli-Pelat, 1992). A modification (Sasa, 1967), which allows easier counting of microfilariae, is to expel 60 μl of blood onto the slide so it forms three even linear strips. The dried films are de-haemoglobinised in buffered water before staining with Giemsa stain. The disadvantage of thick films, like other direct methods, is that they underestimate the prevalence of microfilaraemia if microfilariae densities are low because theoretical detection limit for such procedures is between 15 and 50 microfilariae per ml (Panicker *et al.*, 1991; Turner 1993; Turner *et al.*, 1996; Faris *et al.*, 1997). Even if 60 μl of blood is used subjects with less than 60 microfilariae per ml will not be reliably detected. Another problem is loss of microfilariae from the film during processing especially if anticoagulated blood is used. Southgate (1973) observed a loss of up to 51% and Denham *et al.*, (1971) a loss of between 10% and 40%. Loss of microfilariae is minimised if un-anticoagulated blood is used and the films are dried over night at room temperature (Goldsmid and Rogers, 1976a,b; Partono and Idris (1977). Youssef *et al.*, (1995) have shown that applying a thin film of agar to the thick film

before staining greatly reduces the loss of microfilariae. The thick film does lack sensitivity when microfilariae density is low but is still a useful and cheap technique for survey work where other more sensitive techniques are too expensive (Mouli-Pelat, 1992) or when it is difficult for cultural or other reasons to obtain to obtain venous blood. It has also been suggested that more microfilariae are present in capillary blood than venous blood and that the use of capillary blood may be an advantage when microfilariae densities are low (Eberhard *et al.*, 1988).

There is no question, however, that direct techniques often fail to identify patients with low parasite densities and concentration techniques should be used wherever possible (Southgate and Hamilton, 1974; Weller *et al.*, 1982). Acridine orange staining and fluorescence microscopy has been used as an alternative to Giemsa staining (Goldsmid and Rogers, 1976a,b; Goldsmid *et al.*, 1976) and various combinations of stains can be used to demonstrate the internal structure of microfilariae (Laurence and Simpson, 1969). Counting chamber techniques using various diluents have been used for counting microfilariae by several investigators (Denham, 1971, 1979; Southgate, 1973; Sucharit and Vutikes (1975) and, although not as sensitive as concentration techniques, are a reasonable compromise if 0.1ml of blood is used (McMahon *et al.* 1979).

Concentration Methods for Microfilariae

The most widely used is the method of Knott (1935). One ml of blood is added to 9ml of a 1% formalin solution in normal saline. After red cell lysis is complete the mixture is centrifuged and the deposit examined for microfilariae. Because the formalin preserves the microfilariae, Knott's tests can be set up in the field and processed when the field worker returns to base. The theoretical detection limit is 1 microfilaria per ml. The accuracy of the Knott's test and its ease of use is compromised when blood is processed from individuals with excessive amounts of plasma gamma globulins, a common finding in tropical populations. The formalin precipitates the protein and makes the examination of the deposit very difficult. The method has been improved by Melrose *et al.*, (2000) who added a small of Triton X-100 to the diluent which dissolves most of the proteinaceous deposit and enhances the visibility of the microfilariae.

If a citrate-saponin solution is substituted for the 1% formalin used in the Knott's method, viable microfilariae can be obtained (McQuay, 1970). Another widely used concentration method is the membrane filter technique whereby 1ml of blood that has been diluted in water is passed through a cellulose filter and the microfilariae are trapped on the cellulose (Bell, 1967; Chulerek and Desowitz, 1970; Desowitz *et al.*, 1973a,b; Nathan *et al.*, 1982; Mouli-Pelat

1992). Viable microfilariae can be recovered from the filter for further experimentation or the filter can be fixed and stained and the microfilariae counted. Again, the theoretical detection level is 1 microfilaria per ml but there are some disadvantages of the method.

Blood must be processed at once, the filter apparatus and filters are relatively expensive and it may be difficult to count the microfilariae if large numbers are present. Dennis *et al.*, (1976) conclude that if the volume of blood is increased from 1ml to 5ml the technique is sufficiently sensitive to allow day time blood to be screened for microfilariae in areas where there is nocturnal periodicity alleviating the necessity of night surveys. Moulia-Pelat *et al.*, (1992) found that less periodicity-induced variation in microfilariae counts was obtained with membrane filtration than with capillary blood films but the microfilariae density at the peak of the microfilaraemia was lower in the membrane filter technique than with capillary blood films.

Other concentration methods include: the examination of the buffy coat in a centrifuged microhaematocrit tube (Goldsmid *et al.*, 1972); buffy coat samples prepared from a cytocentrifuge (Goldsmid and Rogers, 1975); and the Becton Dickinson QBC method (Long *et al.*, 1990; Wang, 1998; El Serougi, 1999). The Shandon cytoconcentration technique ordinarily used to detect malignant cells in cerebral-spinal fluid and organisms in broncho-alveolar lavage fluid can be used to detect microfilariae and other blood parasites in diluted, saponin-lysed blood and it is claimed to 7.5 times more sensitive than a thick film (Petithory *et al.*, 1997).

The Diethylcarbamazine (DEC) Provocation Test

If 1.5 to 2 mg of DEC per kg of body weight is given during the daytime microfilariae are "provoked" into leaving the lungs and entering the peripheral circulation where they can be detected by any of the techniques above (WHO, 1987). The use of this test should be discouraged as it has a lower sensitivity than night blood collection and runs the risk of also provoking a severe reaction (see below) especially in areas where *W. bancrofti* occurs with *O. volvulus* or *L. loa* (WHO, 1987). The other antifilarial drugs ivermectin and albendazole do not induce microfilariae to enter the blood during the day (Dunyo *et al.*, 1999).

Detection of Filarial Antigen

Filarial antigenaemia is associated with active filarial infection (Hamilton, 1985) and several investigators have developed assays for filarial antigen using both polyclonal and monoclonal antibodies raised against various antigens (Dasgupta *et al.*, 1984; Hamilton *et al.*, 1984; Reddy *et al.*, 1984; Forsyth *et al.*, 1985; Hamilton, 1985; Harinath, 1986; Lal *et al.*, 1987; Weil and Liffis, 1987; Weil *et al.*, 1987; Zheng *et al.*, 1987; Santhanam *et al.*, 1989; Cheirmaraj *et al.*, 1990). The

first commercial assay (Trop Bio Og4C3 Antigen Test, produced by Trop Bio Pty Ltd, Townsville, Australia) is based on the assay developed by More and Copeman (1990; 1991). The monoclonal antibody is raised against *Onchocerca gibsoni* antigen and shows very strong specificity for *W. bancrofti* antigen. Since it detects antigen from both adult worms and microfilariae the Og4C3 assay will detect amicrofilaraemic and microfilaraemic infections (More and Copeman, 1990; Turner, 1993) and is a very good marker of active filarial infection with adult worms (Chanteau *et al.*, 1994a,b). Unlike microfilaraemia, antigen levels show no significant nocturnal or diurnal variation and blood can be taken at any time of day or night (Moulia-Pelat *et al.*, 1993b). Blood samples taken onto filter paper strips can be used for the Og4C3 assay and Lalitha *et al.*, (1998) and Itoh *et al.*, 1998 found that the capillary blood collected on filter paper and serum gave comparable results. By contrast, Gyapong *et al.*, (1998b) found that the sensitivity of the filter paper test to be significantly inferior with only 50.3% positive with both tests. More and Copeman (1990) showed no cross-reaction in the Og4C3 test with *Brugia* species or other helminths but there is a recent report by Rocha *et al.*, (1996) of a person from a "non endemic" filarial area who tested positive with the Og4C3 test. This individual was found to be parasitised by *Hymenolepis nana*. Whether this represents a true cross-reaction or not is debatable as she also showed a positive immunoblot with crude *B. malayi* antigen. The reported diagnostic sensitivity of the Og4C3 assay for *W. bancrofti* varies from 73% (Chanteau *et al.*, 1994a) to 100% (Lammie *et al.*, 1994). The variation in sensitivity might be partially explained by the variation in the amount of blood used for the detection of microfilariae, which varied from 20µl to 1ml. A more recent study by Rocha *et al.*, (1996) suggests that the sensitivity of the OG4C3 assay may be reduced when microfilariae density is very low. At micro filarial densities of <1, 1 to 30, and >30 the sensitivity was 72.2, 97.6 and 100% respectively and the log OD of the assay was strongly associated with microfilariae density.

The ICT antigen test (ICT Diagnostics, Sydney, Australia) is rapid immuno-chromatographic technique using specific monoclonal and polyclonal antibodies. It utilises capillary or venous blood and is simple enough for field use by people with a minimum amount of training (Weil *et al.*, 1997). The efficacy of filarial antigen tests has been reviewed by Simonsen and Dunyo (1993), Nguyen *et al.*, (1999) and Phantana *et al.*, (1999). Phantana *et al.*, (1999) compared the ICT to thick blood films and membrane filtration and obtained the following results: Sensitivity 100%, specificity 96.3%, predictive value positive 70.7%, predictive value negative 100%. Pani *et al.*, (2000) however, found it less sensitive than the filtration test for detecting low-level microfilaraemia citing an 88.3% positive predictive value when compared with the latter test.

Detection of Filarial Antigen in Body Fluids Other Than Blood

It has been shown by Needham *et al.*, 1996 that there is potential for parasite diagnosis through the detection of antibodies in saliva. A novel, yet untried approach to filariasis diagnosis would be to develop an antigen detection method similar to that used for blood, which could be used on saliva. This would be a much less invasive test than blood collection and would not expose health workers to the danger of blood-borne pathogens such as HIV and hepatitis B and C. Filarial antigen is also present in urine and may demonstrated there even when it is cannot be detected in serum (Das *et al.*, 1987). This anomaly is thought to be caused by the formation of antigen - antibody complexes in the circulation (Das *et al.*, 1987). These immune complexes should not interfere with detection of filarial antigen by the Trop Bio test however as that assay includes a procedure to disassociate antigen - antibody complexes (More and Copeman, 1991).

Detection of Filarial-specific Enzymes

Filarial-specific enzymes have been characterised and show diagnostic potential either as antigens for antibody assays or by the detection of the enzyme itself. Filarial acetylcholinesterase has been identified in the serum of infected people (Misra *et al.*, 1993), and in *in-vitro* culture fluid (Rathaur *et al.*, 1987) and has inter-species cross-reactivity (Sharma *et al.*, 1998). Filarial Glutathione binding proteins, Glutathione-S-transferase, proteinases and superoxide dismutase have been detected in the serum of filarial-infected cattle and humans (Bal and Das, 1995, 1996; Beuria *et al.*, 1995) and the latter has been shown to be strongly antigenic.

Filarial Antibody Assays

There has already been some discussion about filarial antibody in the section on humoral immunity (above). Lack of readily obtainable adult worms makes has made it very difficult to prepare *W. bancrofti* antigen for immuno-diagnosis but the recent introduction of ultrasonic detect (Amaral *et al.*, 1994; Suresh *et al.*, 1997) should make the task easier. The only animals that can be infected with *W. bancrofti* are the leaf monkeys *Presbytis cristata* (Palmieri *et al.*, 1982; Rajasekariah *et al.*, 1986) and *P. melalophos* (Sucharit *et al.*, 1982) and Taiwan monkeys *Macaca cyclopis* (Cross *et al.*, 1979) and these animals are expensive and difficult to maintain in captivity (Franke *et al.*, 1987). *Wuchereria bancrofti* microfilariae can be maintained in culture (Chen and Howells, 1979; Devaney and Howells, 1979; Kharat *et al.*, 1980) or isolated from blood (Jones *et al.*, 1975; Chandrashekar *et al.*, 1984; Van Hoegaerden and Ivanoff, 1986; Franke *et al.*, 1987; El Bassiouny *et al.*, 1993) and used to prepare antigen for antibody studies. The

viability of separated or cultured microfilariae can be ascertained by means of a tetrazolium formazan assay (Comley and Turner, 1990; Mukherjee *et al.*, 1997).

Fortunately, there is marked cross reactivity between filaroid species (Tandon *et al.*, 1981) and wide range of other crude filarial parasites antigens have been utilised for filarial antibody detection: *Dirofilaria immitis* (Amboise-Thomas, 1980; Ata *et al.*, 1993; Turner, 1993; Turner *et al.*, 1993); *Litomosoides carinii* (Hamilton *et al.*, 1981, 1985; Tandon *et al.*, 1981; Rajasekariah *et al.*, 1986); *Setaria cervi* (Tandon *et al.*, 1981, Almeida *et al.*, 1990); *Setaria digitata* (Dissanayake and Ismail, 1980); *B. malayi* (Hamilton *et al.*, 1981; Ottesen *et al.*, 1985; Paranjape *et al.*, 1985; Rajasekariah *et al.*, 1986; Hussain *et al.*, 1987; 1992; Kwan-Lim *et al.*, 1990; Chanteau *et al.*, 1991, 1994, 1995; Hitch *et al.*, 1991; Kurniawan *et al.*, 1993; Murthy *et al.*, 1993; Yazdanbakhsh *et al.*, 1993a,b; Lawence *et al.*, 1994; Mahanty *et al.*, 1994; Terhell *et al.*, 1996); *B. Phahangi* (Weller, 1978; Maizels *et al.*, 1985; Rajasekariah *et al.*, 1986; Hitch *et al.*, 1991; Lammie *et al.*, 1993; Estambale *et al.*, 1994; Addiss *et al.*, 1995; Marley *et al.*, 1995; Wamae *et al.*, 1995; Simonsen *et al.*, 1996) and *Dipetalonema viteae* (Rajasekariah *et al.*, 1986). There is a down side to this cross reactivity - antifilarial antibodies cannot be used to distinguish between filaroid species (Rajasekariah *et al.*, 1986).

Other antigens under evaluation for filarial antibody detection include a chitinase-like recombinant antigen from *W. bancrofti* (Dissanayake *et al.*, 1994, 1995; Raghavan *et al.*, 1994), a cloned antigen with IgG4 specificity (Dissanayake *et al.*, 1992), recombinant *B. malayi* antigen (Kumari *et al.*, 1994; Ramzy *et al.*, 1995), a highly specific antigen from *W. bancrofti* 3rd stage larvae (Burkot *et al.*, 1996), fractionated urinary antigen (Ramaprasad and Harinath, 1995), fractionated circulating filarial antigen (Cheirmaraj *et al.*, 1992). Antibodies to recombinant paramysin from *B. malayi* can be used as a marker for adult worms of that species (Langy *et al.*, 1998).

Antifilarial IgG antibody reacts with a wide range of other helminths and protozoa due to shared phosphocholine antigens (Ambroise-Thomas 1974; Voller and de Savigny 1981; Taylor and Denham 1985; Lal and Ottesen 1989; Chanteau *et al.*, 1995). Species shown to cross react include *Ascaris suum* (Betschart *et al.*, 1990; Willenbacher *et al.*, 1993), *Ascaris lumbricoides* (Kagan *et al.*, 1963; Paranjape *et al.*, 1985; Gam *et al.*, 1987; Murthy *et al.*, 1993; Rahmah *et al.*, 1994), *Trichuris trichiura* (Rahmah *et al.*, 1994) and hook worm (Gam *et al.*, 1987), *Strongyloides* (Gam *et al.*, 1987, Rahmah *et al.*, 1994), *Fasciola hepatica* and *Shistosoma mansoni* (Rahmah *et al.*, 1994) and *Trichinella spiralis* (Kagar *et al.*, 1963).

The degree of cross-reactivity between filaroids and other parasites vary between different species of filaroids. Saxcena *et al* (1981) showed no cross reactivity between *L. carinii* and *Ancylostoma*, *Ascaris*, *Taenia* or amoeba (species not specified).

Various methods can be used to detect filarial antibody: Complement fixation, indirect haemagglutination, gel diffusion, immunoelectrophoresis, counter current immunoelectrophoresis, indirect immunofluorescence and enzyme-linked immunosorbent assay (ELISA) have all been used (Kagan, 1963; Ambroise-Thomas, 1980). Currently, almost all filarial antibody studies use ELISA and the following discussion deals with that method of detection unless otherwise indicated. Cross reactivity limits the usefulness of IgG antibody in filariasis diagnosis but they may be of some value in communities where parasites other than *W. bancrofti* are absent or rare (Chanteau *et al.*, 1991) and have been used to diagnose occult filariasis in Indian children (Chaturvedi *et al.*, 1995). Because humans are not able to synthesise anti-phosphocholine or anti-carbohydrate antigen IgG4 (Maizels *et al.*, 1987; Scott *et al.*, 1987; Lal *et al.*, 1991) the assay of filarial IgG4 antibodies greatly increases specificity and enhances the diagnostic ability of the test (Lal and Ottesen, 1988; Kwan-Lim, 1990; Rahmah *et al.*, 1994; Haarbrink *et al.*, 1995; Chanteau *et al.*, 1995; Terhell *et al.*, 1996) and Mahanty *et al.*, (1994) have shown that antifilarial IgG4 is a good index of the intensity and duration of filarial exposure in endemic populations, and (Maizels *et al.*, 1995) found that the level of IgG4 antibody correlates with microfilariae counts. Filter paper collection techniques can be used to detect filarial IgG4 antibody (Chanteau *et al.*, 1991; Terhill *et al.*, 1996). IgG4 antibodies are extremely useful for the detection of *Brugia* species because they will not be detected by the current antigen tests because they are specific for *W. bancrofti* (Rahmah *et al.*, 1998; Haarbrink *et al.*, 1999a). There appears to be very little literature on the usefulness of anti-filarial IgM antibody for filariasis diagnosis but Ata *et al.*, (1993) found it to be more specific than IgG. Chanteau *et al.*, (1992) trialed an IgA antibody test but found it unhelpful. It should be borne in mind that sero-positivity does not always indicate active filarial infection. The antibody may have been raised by exposure to infective larvae without an adult worm being present or the antibody may persist after parasites have been cleared and although the sensitivity of antibody tests is high, the specificity, and hence the predicative value, is low (Chanteau *et al.*, 1991).

Demonstration of Parasite DNA by DNA Probes or the Polymerase Chain Reaction (PCR)

Polymerase chain reaction methods have been successfully used in the detection of *W. bancrofti* DNA in sputum (Abbasi *et al.*, 1996, 1999), *B. malayi* DNA in blood (Lizotte *et al.*, 1994; Rahmah *et al.*, 1998). *W. bancrofti* in blood, plasma and paraffin-embedded tissue sections (McCarthy *et al.*, 1996), urine (Lucena *et al.*, 1998) and *W. bancrofti* larvae in infected

mosquitoes (Chanteau *et al.*, 1994c; Nicolas *et al.*, 1996; Furtado *et al.*, 1997). A combination PCR-ELISA method for detection of filarial DNA has been devised by Fischer *et al.*, (1999).

The Demonstration of Adult Worms

The use of ultrasonography to detect adult worms in the scrotum (Amaral *et al.*, 1994; Simonsen *et al.*, 1997a; Faris *et al.*, 1997; Suresh *et al.*, 1997; Dreyer *et al.*, 1999) has already been dealt with above. The same technique can be used to detect adult worms in the breast (Dreyer *et al.*, 1996a,b) and has detected viable worms in children (Dreyer *et al.*, 1999). The need for specialised technology means that this diagnostic tool is not available to most of the at-risk population.

Skin Testing

Skin tests using filarial antigens which were usually prepared from *D. immitis* (Bozicevich and Hutter, 1944; Franks *et al.*, 1947; Sawada *et al.*, 1965) were widely used for filariasis diagnosis from the 1940's to the 1970's and the subject has been extensively reviewed by Kagan (1963) Desowitz *et al.*, (1966) and Ambroise-Thomas (1980). The Sawada antigen skin test was evaluated by Smith *et al.*, (1971). They obtained 22% false positives in non-exposed controls, 53 to 85% false negatives and there was much cross-reaction with other helminth species. Because of this lack of sensitivity and specificity, the skin test has been largely abandoned although it may have a part to play in the monitoring of filarial control programs (Desowitz, 1966; Ambroise-Thomas, 1980; Tada *et al.*, 1982).

Which Test is the Best One to Detect Filarial Infection?

To quote Southgate and Hamilton (1974):

"Clearly, the unexpected prevalence rates determined by the newer techniques will change our ideas on filariasis control; even in areas where it has been recognised as highly prevalent for years, the amount of filariasis obviously been under estimated. Apparently rather small changes in the techniques of epidemiological measurement in estimating the prevalence and densities of microfilaraemia in a human population can produce quite remarkable changes in our views of the epidemiology of the disease. Clearly our changing understanding of epidemiology has very serious implications for control programs."

These comments were made during a study of the value of concentration methods to estimate the prevalence of microfilaraemia but they stress the necessity of utilising the very best test available. The question "which is the best test available" can be best answered by quoting Chanteau *et al.*, (1994). "The detection of microfilariae is very specific but insensitive, the antibody tests are very sensitive but not specific enough, and finally the Og4C3 antigen test seems to be a good compromise of specificity and sensitivity for the diagnosis of Bancroftian filariasis". This sentiment is echoed by Simonsen and Dunyo (1999). The other commercially available antigen test, the ICT rapid card test is now widely accepted as the tool of choice for survey work because of its simplicity and its ability to provide "on the spot" data. While it has been more or less decided what the most appropriate tests for filariasis in patients from endemic areas are, the situation with acute expatriate cases is

more difficult. IgG anti-filarial antibodies can be demonstrated in most exposed expatriates but this is not a useful marker of active infection as it could just be the result of immunological stimuli of injected L3 larvae, infection or contact with an animal filaroid, or could be due to cross reactions with other helminth parasites. IgG4 antibodies are slow to develop and this limits their usefulness. IgM antibodies are too non-specific to be of any value. Expatriates do not seem to become antigen positive (why that is so would be an interesting topic for further research) and microfilaraemia is extremely rare. It has been suggested that testing for low levels of antifilarial IgE may be of value (Nutman 1997, *pers. comm.*) and this should be explored. Currently, the resolution in symptoms and the reduction in eosinophils after a diethylcarbamazine therapy is often the best indicator of infection.

Control of Lymphatic Filariasis

There are several approaches to controlling vector-borne parasitic diseases: vaccination, vector control, breaking of vector-host contact by use of repellents and bed nets, and chemotherapy.

Vaccines

Despite some promising research, a vaccine for any filarial parasite (and indeed any parasite) is not yet a reality (Philip *et al.*, 1988; Selkirk *et al.*, 1992; Chenthamarakshan *et al.*, 1995; Dissanayake *et al.*, 1995; Grieve *et al.*, 1995). There has recently been some progress in this area with cloned filarial antigens, a recombinant chitinase and a protective epitope SXP1, an antigen found in multiple worm stages showing promise as vaccine candidates (Cryz 1991; Wang *et al.*, 1997) and Zang *et al.*, (1999) suggests that a serine protease inhibitor expressed by *B. malayi* microfilariae that inhibits neutrophil serine protease and aids the parasite to evade the immune system, may be a target for vaccine development.

Vector Control

Vector control does have a part to play and can be very successful when malaria and filariasis have common vectors. In the Solomon Islands for instance, where both malaria and filariasis are transmitted by *Anopheles farauti* and *Anopheles koliensis*, the prevalence of microfilaraemia was reduced from around 15% to <2% by vector control carried out during a malaria campaign (Webber, 1977, 1991). Vector control however does take a long time to become effective. For instance, Schuurkamp *et al.*, (1987) estimated that it would take 11 years to reduce the prevalence of microfilaraemia to <2% in the Tabubil area of the Western Province of Papua New Guinea by vector control alone. Such prolonged campaigns are very labour intensive and costly and the development of insecticide resistance makes them become less and less effective. Data collected during a 5-year vector control program in Pondicherry, India, found that vector control alone might have little impact on the overall age-prevalence of infection even when sustained for long periods (Srividya *et al.*, 1996). Toilet pits and the like are prime breeding grounds for *Culex quinquefasciatus* (Nwoke *et al.*, 1993) and the use of polystyrene beads can be a very effective control mechanism for this and other *Culex* species. By use of this method Maxwell *et al.*, (1989) reduced the numbers of mosquitoes entering houses by about 97% Webber and Southgate (1981) have shown that in the Solomon Islands approximately 8 mosquito bites per person per night are required for reliable transmission of filariasis. Maxwell *et al.*, (1999) found that although the treatment of cesspools and pit toilets with polystyrene beads reduced the mosquito population in houses the cost of achieving this could not be justified by its impact on filariasis alone, but the

reduction in nuisance biting may increase the community support for other control programs

Insecticide-treated Bed Nets

Insecticide-treated bed nets have the potential to reduce filariasis transmission (Charlwood and Dagoro, 1987) but they are of only limited effectiveness because there are a large number of mosquito species that transmit filariasis and some of these bite during the time of when people are not under bed nets.

Chemotherapy

There are essentially two regimes used to control filarial infections using chemotherapy: selective and mass treatment. In selective therapy individuals are examined for the presence of disease and those who are found to be infected are treated. There are major problems associated with that approach. If microfilaraemia is used as the indicator of infection many infected people will be overlooked, as not all those who have the disease are microfilaraemic. People with very low micro filarial densities may not be detected and it has been shown that such people are capable of infecting mosquitoes and causing a resurgence of disease (Lowrie *et al.*, 1989). The use of antigen testing will overcome these problems but you are still left with the logistical problem of screening all members of the community. If selective chemotherapy is used the lost infections are balanced by new infections to produce a dynamic equilibrium and there needs to be continual reassessment of the filarial status of the community to identify the newly infected. By studying the transmission potential of a number of diseases, including helminth parasites, Woolhouse *et al.*, (1997) found that, typically, 20% of the infected host population contributes at least 80% of the net transmission potential and that control programs which fail to reach all of that core group, a problem with selective treatment programs, will be less effective in reducing levels of infection in the group as a whole. Using mass chemotherapy regardless of parasite status excludes the problems of selective therapy. Mass treatment aims to treat all members of the endemic community at the same time and will therefore treat pre-patent as well as patent infections. In other words the community becomes the focus of the control program rather than the individual. Originally antimony and arsenic based drugs and naphthalene sulfonic acid (Suromin) were used to treat filariasis but with limited success (James and Gillies, 1985; Campbell and Rew, 1986). These compounds had a moderate amount of anti-filarial activity but were very toxic. Suramin has been shown to be moderately active against adult *W. bancrofti* and *Brugia pahangi* (Howells *et al.*, 1983) but it has no effect on microfilariae and will therefore not interrupt the transmission cycle in the short term. It is also more

toxic than other antifilarial drugs and is not widely used today (James and Gillies, 1985). Levamisole has been shown to have limited activity against filarial parasites (Duke 1974; O'Holohan and Zaman 1974; Rogers and Denham 1976; McMahon 1979).

The Use of Diethylcarbamazine (DEC)

For the last 50 years the piperazine derivative diethylcarbamazine (DEC, proprietary names: Hetrazan, Banocide and Notezine) has been, and still is, the most widely used drug for the treatment of lymphatic filariasis (James and Gillies, 1985; MacKenzie and Kron, 1985; Campbell and Rew, 1986). Diethylcarbamazine was discovered by Hewitt *et al.*, in 1947 and used to treat human filariasis in the same year (Santiago-Stevenson *et al.*, 1947). The pharmacology of DEC has been extensively investigated (for reviews see: Hawking, 1950b; Hawking *et al.*, 1950a; MacKenzie and Kron, 1985; Maizels and Denham, 1992) and the following summary is taken from those publications unless otherwise indicated. Diethylcarbamazine is given orally and is rapidly absorbed reaching peak blood levels one to hours after ingestion. It reaches all parts of the body within 25 minutes after the dose is given. Accurate blood levels of DEC can be determined by ELISA (Mitsui and Aoki, 1998). The plasma half life varies from 6.1 to 8.1 hours. Excretion is mainly renal and the blood concentration reaches zero within 48 hours. Diethylcarbamazine, albeit in high concentrations and in *in-vitro* experiments, has been shown to affect several systems of the body. It can have a depressive effect on myocardial activity, stimulate the production of hepatic enzymes and produce relaxation of intestinal tissue (MacKenzie and Kron, 1985). Joseph and Dixon (1984) found that DEC enhances the ability of oxytocin and acetylcholine to induce contractions of uterine tissue and that led Subbu and Biswas (1971) to recommend that DEC should not be given to pregnant women, but in practice however, there have been no reports of abortifacial or teratogenic effects. Diethylcarbamazine does not have any effect on the white cells of non-filarial infected patients but it does cause a rapid fall in eosinophils which often show loss of granulation and vacuolation suggesting that these cells have a role in the destruction of microfilariae. Diethylcarbamazine causes the release of serotonin from platelets and stimulates the production of platelet adhesion factor and thromboxane. Paradoxically it also appears to inhibit platelet aggregation induced by adenosine diphosphate and collagen.

Hawking *et al.*, (1950a) suggested that DEC acted as an opsonin and enhanced phagocytosis of microfilariae but the exact mechanism by which DEC destroys microfilariae is still a matter of debate (see the comprehensive review by MacKenzie and Kron (1985). Mukhopadhyay and Ravindran (1997) have found evidence that antibodies formed against the drug

itself potentiate the action of antifilarial antibodies but this finding is disputed by Cesbron *et al.*, (1987) whose study suggests that blood platelets play a role in DEC-mediated microfilariae killing possibly involving a free radical process.

In summary, the main modes of action thus far identified are focussed in two main areas: the effect of the drug itself upon the parasite and the facilitation of host-parasite interactions. Diethylcarbamazine has been shown to overstimulate parasite neuromuscular systems and increase motility, to inhibit vital parasite metabolic enzymes, activate complement on the parasite's surface membrane, activate eosinophils and stimulate them to produce eosinophil-derived cationic proteins, enhance eosinophil-dependant antibody-mediated destruction of parasites and increase the adhesion of parasites to phagocytic and antibody-producing cells. It has been demonstrated by Bhattacharya *et al.*, (1997) that one of the ways that filarial parasites ensure their survival in the host is to secrete acetylcholinesterase into the circulation of the host, which degrades circulating acetylcholine. Since acetylcholine stimulates the release of lysosomal enzymes such as β -glucuronidase and acid phosphatase, which are involved in phagocytosis, is depressed during filarial infections. After DEC treatment the level of plasma beta-glucuronidase and acid phosphate rose and this is believed to be due to DEC inhibiting the production of parasite acetylcholinesterase. Another focus in recent years has been the role of nitric oxide in microfilariae clearance. *In-vitro* tests have shown that Microfilariae are killed by exposure to nitric oxide (Taylor *et al.*, 1996) and Winkler *et al.*, (1998) has shown that high levels of nitrite and nitrate were released into the serum during microfilariae killing.

Another contentious issue is whether DEC is macrofilaricidal as well as microfilaricidal. MacKenzie and Kron (1985) concluded that it had a limited amount of macrofilaricidal activity and their opinion is supported by Weil *et al.*, (1988). The degree of activity however, is still questionable (Ottesen, 1985, Weil *et al.*, 1988). There is also histological evidence - Figueredo-Silva *et al.*, (1996) removed nodules after DEC treatment and found almost all contained degenerating adult worms. Experiments by Ismail *et al.*, (1996) suggest that fortnightly 10mg/kg doses of DEC suggested that there was some macrofilaricidal activity but the outcomes were inconsistent and the finding by Weil *et al.*, (1988) that filarial antigenaemia persists for up to 12 months after DEC therapy has eliminated circulating microfilariae suggests that DEC is only partly effective against adult filarial parasites. Eberhard *et al.*, (1997) reported that although filarial antigen fell after treatment, in no case did it fall to zero, even in individuals who remained amicrofilaraemic for several years after treatment. This they suggest was evidence that adult worms were persisting. The absence of microfilariae would suggest however, that if the adult females are still living they are prevented from

producing microfilariae or that DEC in some way, facilitates the rapid removal of microfilariae from the circulation. In any case, the lack of microfilariae means that the transmission cycle has been interrupted. There is no doubt however that DEC is a highly effective microfilaricidal drug for *W. bancrofti*. Jing-Yuan *et al.*, (1991; 1992) when assessing DEC treatment and residual spraying in China, found that Bancroftian filariasis had been eliminated and that subsequent surveillance showed no signs of resurgence. The question of parasite resistance to DEC has been raised since the early days of its use. Hawking (1950a,b) and Otto *et al.*, (1953) both noted that small numbers of microfilariae persisted in some patients after treatment. Zhong-Xing and Zhong-Jun (1989) found that in former filariasis endemic areas where the disease has been largely controlled, a few cases of low-density microfilaraemia remained and could possibly be related to parasite tolerance to DEC.

Kimura *et al.*, (1985) reported that annual low dose treatment with DEC did not reduce microfilariae densities in subjects with high pre-treatment microfilarial levels. Eberhard *et al.*, (1991b) found that there appeared to be evidence for non-susceptibility to DEC in Bancroftian filariasis. They found that treatment with DEC does not always result in total clearance of microfilariae and suggested that filarial parasites may develop partial resistance. The standard DEC treatment regime is 6 mg per Kg body weight per day over a 10 to 20 day period (James and Gillies, 1985; WHO, 1992; 1994). This is suitable for treated isolated cases but it is not a suitable regime for community-wide mass treatment programs because it is labour-intensive and incurs high costs (Michael *et al.*, 1996). Other regimes for mass treatment are: monthly doses, 6 monthly doses and annual doses. All these methods are effective but not all are equal when assessed on a cost-effective basis (Michael *et al.*, 1996)

DEC Medicated Salt

Common cooking salt medicated with DEC in concentrations ranging from 0.1% to 0.6% (Gelbrand 1994; WHO, 1992; 1994) has been effectively used in mass treatment programs for Bancroftian filariasis in China (Jing-yuan *et al.*, 1991; 1992; Liu *et al.*, 1992; Shaoqing *et al.*, 1994), India (Rao *et al.*, 1981; Chandrasekharan *et al.*, 1984; Narasimham *et al.*, 1989; Krishnarao *et al.*, 1991) and Tanzania (Meyrowitsch *et al.*, 1996; Michael *et al.*, 1996) and has shown to be effective in brugian filariasis (Chandrasekharin *et al.*, 1984; Shenoy *et al.*, 1998b, 1999), but it is logistically difficult to organise due to the bulky nature of the salt, and only slightly less expensive than using the standard dose regime.

Mass Drug Treatment with DEC

Current practice is a single annual dose of 300 mg of DEC for adults and 150 mg for children (often combined with Ivermectin, and sometimes

Albendazole - see below). This simplifies the treatment, increases compliance and can be easily accommodate into existing primary health care networks without over-burdening them (Wijers, 1984; Kimura and Mataika, 1996; Taylor and Turner, 1997).

Low dose DEC has been shown to significantly reduce the prevalence and density of microfilariae in treated communities and reduce the prevalence of chronic pathology (Forsyth, 1987; Partono *et al.*, 1984, 1989). Panicker *et al.*, (1991) found that there was a reduction in microfilariae prevalence of 74.9% in the annual treatments and 90% in the biannual treatments.

He also found that attacks of filarial fever and incidence of recent oedema cases were also significantly reduced after DEC treatment. A trial in Tanzania achieved microfilariae clearance rates of 92% (Meyrowitsch *et al.*, 1996). Meyrowitsch *et al.* (1998) have shown that the beneficial effects of DEC treatment persist for at least 4 years and although microfilaraemia does recur in some patients, especially those who had high microfilaria densities before treatment, the level of microfilaraemia does not reach the pretreatment levels. Khan *et al.*, (1998) found that 5 years after a dose of 72 mg/kg of DEC given over a 21 day period, 51% of subjects were still amicrofilaraemic, 36.4% had densities less than pre-treatment levels and 11.8% had increased microfilariae counts. In Samoa, 3 annual treatments of DEC 6 mg/kg, achieved an estimated 80% reduction of microfilaria in the population and lowered the annual transmission potential from 2.18 to 0.67 (Kimura *et al.*, (1992). Mataika *et al.*, (1998) obtained similar results with a five-year annual treatment program in Fiji.

Other Benefits of DEC Therapy

There are several other benefits apart from parasite killing in the treatment of filariasis with DEC. Schuurkamp *et al.*, (1992) found that a significant reduction in splenomegaly within the communities around the Ok Tedi Mine after a successful control programme with DEC. Piessens *et al.*, (1981) stated that *in vitro* lymphocyte proliferative responses to microfilariae antigens increased in patients who became amicrofilaraemic after treatment with DEC. Piessens' group also showed that no changes were observed in amicrofilaraemic individuals who were given DEC or in those who remained microfilariae positive after treatment. Partono *et al.*, (1981) showed that the total number of working days lost due to filariasis was substantially reduced after treatment with DEC in Indonesia with the lymphadenitis rate falling from 46% and the prevalence of elephantiasis falling from 17% to 4%.

There is strong evidence that DEC has some activity against a range of non-filarial infectious agents (Mejia *et al.*, 1996) - probably due to its opsonic activity and enhancement of phagocytosis (Hawking *et al.*, 1950a,b; Mejia *et al.*, 1996). It has been shown to enhance blood microbicidal activity (Kitchen *et al.*, 1995a,b; 1998), to be effective in the treatment of chronic bacterial osteomyelitis (Kitchen *et al.*, 1995b),

and to influence the course of bacterial and fungal infections in mice (Kitchen *et al.*, 1992; 1993) and humans (Mejia *et al.*, 1996). It has been shown that bacterial and fungal infection play a pivotal role in the development of filarial elephantiasis (Addiss *et al.*, 1995) and it has been suggested that DEC's antibacterial and antifungal effects may supplement its antifilarial action when it is given to treat filarial disease (Mejia *et al.*, 1996). It has been known for some time that DEC therapy improves T cell responses to purified protein derivative of *Mycobacterium tuberculosis* (PPD) in filariasis and onchocerciasis patients (Green *et al.*, 1985; Sartono *et al.*, 1995b) but there is also a recent suggestion by Kitchen (1995a, 1998) that the anti-mycobacterial effects of DEC may not be limited to the filarial-infected. If this is proved to be the case it is possible that a major side benefit of mass DEC therapy for filariasis control could be reduction in tuberculosis, leprosy and atypical mycobacterial infections in filarial-endemic areas. Diethylcarbamazine has also been shown to have therapeutic potential in feline and murine leukaemia virus infections (Kitchen *et al.*, 1990; Nelson *et al.*, 1995) and has been shown to increase antibody levels in feline oncornavirus-associated cell membrane antigen (Kitchen and Cotter, 1998).

Use of Ivermectin

Ivermectin has been used very successfully for the treatment of onchocerciasis for a number of years (for reviews see Goa *et al.*, 1991; Townson *et al.*, 1994) and a single annual dose of 400 µg/kg, either alone or in combination with Diethylcarbamazine, has proved to be very effective producing long-term suppression of microfilaraemia in Bancroftian lymphatic filariasis in a number of countries (Diallo *et al.*, 1987; Kumaraswami *et al.*, 1988; Ottesen *et al.*, 1990; Campbell 1991; Ismail *et al.*, 1991; Zheng *et al.*, 1991 a,b; Cartel *et al.*, 1992; Eberhard *et al.*, 1992; Addiss *et al.*, 1993; Kazura, 1993b; Moulia-Pelat *et al.*, 1993a, 1994a,b, Coutinho *et al.*, 1994; Nguyen *et al.*, 1994; Ottesen and Campbell, 1994; Chodakewitz 1995; Dreyer *et al.*, 1995a; Moulia-Pelat *et al.*, 1995; Ottesen and Ramachandran 1995; Nguyen *et al.*, 1996; Cao *et al.*, 1997) and is equally effective against brugian filariasis (Shenoy *et al.*, 1993). Nguyen *et al.*, (1996) found that twice yearly doses of 100 µg/kg of Ivermectin did not reduce the prevalence of microfilaraemia, but when the dose was increased to 400 µg/kg the prevalence dropped from 21% to 7% and the microfilariae density to 0.5% of its initial value. Plaisier *et al.*, (1999) found that Ivermectin removes 100% of microfilariae regardless of dose and at a dose of 400 µg/kg a single treatment irreversibly reduces the microfilariae of the adult parasite by at least 65%. Zheng *et al.*, (1991a,b) found that Ivermectin was especially useful in the treatment of filarial relapses after DEC treatment. Like DEC, Ivermectin treatment does give rise to adverse side effects. Cao *et al.*, (1997) describe them as mild and "flu like", generally mild, and well tolerated by

patients. Mild reactions were also noted by Zheng *et al.*, (1991) who also found that local flare-up of acute filariasis was less likely with Ivermectin than with DEC. The severity of the reaction was strongly associated with the pre-treatment microfilariae density but independent of the dose (Cao *et al.*, 1997). Weil *et al.*, (1991) states that local adverse reactions such as nodule formation, lymphangitis and epididymitis are not seen with Ivermectin therapy because these reactions are caused by the death of adult filarial worms and Ivermectin is not a macrofilaricide. An ELISA for blood levels of Ivermectin is available (Mitsui *et al.*, 1996).

As with DEC, the macrofilaricidal action of Ivermectin is debatable. Ismail *et al.*, (1996) found that multiple, high dose Ivermectin treatment (12 fortnightly doses of 400 µg/kg) does have macrofilaricidal activity but the results are neither predictable nor consistent. Eberhard *et al.*, (1997) reported that although filarial antigen fell after treatment, in no case did it fall to zero, even in individuals who remained amicrofilaraemic for several years after treatment suggesting that some adult worms survived. By contrast however, Dreyer *et al.*, (1995b; 1996) could find no evidence of macrofilaricidal activity. In their study 15 men who had living, adult *W. bancrofti* detected by ultrasound were treated with 400 µg/kg body weight of Ivermectin at 2 week intervals for 6 months (total dose 4.8 mg/kg). Microfilaraemia was rapidly suppressed but no changes in the motility or location of the adult worm were detected. In another study Dreyer *et al.*, (1995a,b) removed live adult worms from several patients 8 months after being treated with 400 µg/kg of Ivermectin.

Effect of Ivermectin on Mosquito Survival

There is strong evidence that Ivermectin ingested from a treated person during a blood meal decreases mosquito survival and fertility (Tesh and Guzman, 1990; Nasr *et al.*, 1996; Bockarie *et al.*, 1999).

Use of Albendazole

Albendazole has been used for the treatment of intestinal helminths for a number of years but it has only recently been trialed as an anti-filarial. Addiss *et al.*, 1997a,b have shown that at a stated dose of 400 mg per person it is an effective microfilaricide. It is even more effective when combined with DEC or Ivermectin (see) below. The problem of adverse reactions with Albendazole is no better or worse than with DEC or Ivermectin (Addiss *et al.*, 1997a,b). The potential for Albendazole to be used as a macrofilaricide for the treatment of individual patients is regarded by Ottesen *et al.*, (1999) as one of the most important questions in filarial research.

Combination Therapy

A combination of DEC and Ivermectin has shown to be very effective in providing rapid and long-term clearance of microfilariae (Moullia-Pelat *et al.*, 1994a,b, 1996). Nicolas *et al.*, 1997 found that DEC and Ivermectin was much more effective than either drug alone for clearing circulating filarial antigen in both amicrofilaraemic and microfilaraemic subjects. Addiss *et al.*, (1997) have shown that a combination of 200-400 µg/kg of Ivermectin plus 400 mg of albendazole is more effective in clearing microfilariae than either drug alone. Ishmail *et al.*, 1998 studied the effects of Albendazole, DEC, and Ivermectin alone and in combination and showed that although all were well tolerated and effective macrofilaricides, a single dose of a combination of 600 mg of albendazole and 400 µg/kg of Ivermectin was the most effective. Decreasing levels of filarial antigens after treatment suggested that all four regimes have significant macrofilaricidal activity. The most effective, however, was a combination of 600 mg of Albendazole with 6 mg/kg of DEC. With this combination, filarial antigen levels decreased by 77% 15 months after therapy.

Other Advantages of Combination Therapy

Another advantage of antifilarial therapy using Albendazole and/or Ivermectin is the potent broad spectrum effects against intestinal helminths (Ismail *et al.*, 1998). Another possibility, if Ivermectin is used, is controlling scabies as well. Beach *et al.*, (1999) have shown in a randomised, placebo-controlled study that a 6 monthly combined treatment with Albendazole and Ivermectin was effective in controlling *W. bancrofti*, *A. lumbricoides*, *T. trichuria* and hookworm in Haitian primary school children with an obvious benefit to their height and weight. The cost effectiveness of controlling more than one parasite with an annual program is obvious and studies to confirm these possibilities are urgently needed.

The Search for New Antifilarial Agents

There is a very active global search for new antifilarial compounds, the ultimate goal being the production of a safe effective macrofilaricide (Lazdins and Kron, 1999). *In-vitro* effects of potential anti-filarial can be studied by use of the colorimetric viability test derived by Comley *et al.*, (1989).

Extracts of a wild Indian herb, *Andrographis*

paniculata and the ginger plant *Zingiber officinale* have been shown to have activity against canine filariasis (*D. immitis*) both *in-vitro* and *in-vivo* (Dutta and Sukul, 1982, 1987) and a major search for new natural anti-parasite drugs is being carried out in the Philippines (Monzon, 1995). Animal trials of three promising new agents were reviewed by Mak *et al.*, (1991). CGP 30376, a derivative of Benzothiazole and has been shown to be both micro and macrofilaricidal but there is concern about its hepatic toxicity. The methyl-piperazine compound 6140 CGP has significant macrofilaricidal activity and CGI 18041 (its chemistry is not revealed) kills both microfilariae and adult worms. Gayral *et al.*, (1995) have identified a group of ethynsulphonamides, epoxyethanesulfonamides and their carboxamide analogues that show some promise as antifilarial agents. Phenoxycyclohexane derivatives are showing promise in *in-vitro* studies, as are thiosemicarbazones in animal studies (Loiseau and Depreux, 1993). Beta-carbolines (Srivastava *et al.*, 1999), bis-cationic hetero-aromatics (Sundberg *et al.*, 1998) and beta-amino substituted 4-amino-3-nitropropionophenones (Varma Rajendra *et al.*, 1992) have shown promise as micro and macrofilaricides in *in-vitro* and animal experiments. An analogue of ubiquinone, 2,3-Dimethoxy-5-methyl-1,4-benzoquinone (QO) has been shown to irreversibly paralyse adults and microfilaria of *S. digitata* and the microfilariae of *W. bancrofti* suggesting that this class of drugs has the potential for anti-filarial therapy (Sivan and Kaleysa, Raj 1999).

Another exciting approach to therapy is to use the lymphatic system rather than the blood for delivery of antifilarial drugs. Adult filarial worms, living as they do in the lymph nodes, are in intimate contact with lymph and this approach may lead to the discovery of a potent macrofilaricide. Derivatives from drugs used in tumour therapy - Closoantel and chlorambucil showed some promise in *in-vitro* experiments but no activity could be detected *in-vivo* (Loiseau *et al.*, 1997).

The search for new antifilarial drugs has moved to the marine environment and the antifilarial activity of a synthetic marine alkaloid, aplysinopsin (CDRI compound 92/138) derived from Australian reef sponges shows *in-vitro* activity (Singh *et al.*, 1997).

Another novel approach suggested by Hoerauf *et al.*, (1999, 2000) is to target the endosymbiont bacteria that live within filaroids and may have a beneficial effect on microfilarial growth and survival. Treatment of onchocerciasis patients with tetracycline resulted in the sterilisation of the female worms.

Adverse Reactions to Anti-Filarial Drugs

Post-treatment reactions are common in lymphatic filariasis and onchocerciasis (where it is called the Mazzotti reaction) and although most cases are relatively mild, they can cause communities to be wary of treatment (Sunny *et al.*, (1986) and lead to failure of control programs (Ottesen, 1980; Forsyth, 1987; Schuurkamp, 1992; Schuurkamp *et al.*, 1990, 1992; Fan, 1992; Turner *et al.*, 1994). Fever is the most common manifestation, but lymphadenitis, arthralgia, chills, drowsiness, headaches, hypotension and urticaria are also common. The reaction is not caused by the drug itself but by an inflammatory reaction induced by dying microfilariae and severity tends to increase with increasing numbers of circulating microfilariae (Piessens *et al.*, 1981b; Francis *et al.*, 1984; Haarbrink *et al.*, 1999b). Laigret (1983) found that reactions to DEC were less frequent as the number of treatments increased. The exact cause of the

treatment reaction is still unknown but it has been shown that cytokines such as Interleukin 6 (IL-6) and tumour necrosis factor (TNF) are implicated (Turner *et al.*, 1994). Anti-inflammatory drugs and steroids can be used to lessen the reaction (Ottesen, 1987; Stingl *et al.*, 1988) but the effects are not consistent and concern has been raised that the use of these drugs may reduce the killing efficacy of antifilarial drugs (Schofield and Rowley, 1961; Adwadi *et al.*, 1982) and Dreyer and de Andrade (1989) are of the opinion that the mild nature of most reactions means that such drugs are seldom warranted. As well as the systemic effects mentioned above, localised reactions such as nodule formation, lymphadenitis and epididymitis are sometimes seen near sites occupied by dying adult worms. This type of reaction is more common when DEC is used because of its at least partial, macrofilaricidal action (Weil *et al.*, 1991).

Effect of Therapy on Microfilaraemia and Immunological Responses

All authors report a rapid decrease in microfilariae levels following therapy. Typically, results are seen within 24 hours and in most cases microfilariae are cleared from the blood within 7 days. There is considerable variation in the length of time that patients remain amicrofilaremic after treatment, varying in different studies from 1 to 12 years (WHO, 1987).

Usually, filarial antibody levels against whole parasite antigen rise immediately after therapy peaking at about 30 days, they fall during the next 12 months to below treatment levels but can remain positive for up to 10 years (Gao *et al.*, 1994). Bal and Das (1999) found that DEC therapy increased the level of IgG and IgM antibodies. Antibodies of different IgG sub-classes and IgE antibody behave differently. After therapy IgG4 antibody levels decreased sharply and were down to 65-78% of the pre-treatment levels within 12 months. IgG1 levels declined in a less predictable manner and IgG2 and IgG3 only declined in elephantiasis patients who have an elevation of these antibodies to start with. IgE showed only a mild reduction (7-28%) over 12 months and 56% over 2 years (Atmadja *et al.*, 1995; Kurniawan *et al.*, 1995). Wamae *et al.*, (1992) found that IgG4 antibody rose markedly after treatment with DEC or Ivermectin to peak at 30 days and then began to decrease. Post treatment microfilariae density was inversely correlated with the decrease in IgG4 antibody and gave a good indication of the success of treatment. Murthy *et al.*, (1997) also found that treatment elevated IgG4 antibody and a reduction in IgG1 antibody. IgG2 and IgG3 antibodies remained unaffected. A rather different pattern is seen if antigen from microfilariae is used. Antibodies to microfilariae are usually absent from the blood of microfilaremic or present only in low titre. Anti-filarial treatment and the clearance of

microfilariae are followed by a rapid rise in antibodies against microfilariae (Grove, 1981).

Sartono *et al.* (1995b) studied a group of patients one year after therapy and found that cellular immune responses to filarial antigens had increased significantly but the response to PPD remained low. Similar results were obtained by (Gopinath *et al.*, 1999). These studies confirm the earlier work of Piessens *et al.*, (1981). Lammie *et al.*, (1992) found that interferon production from filarial-antigen-stimulated mononuclear cells increases but there is no significant change in IL-4 production. Zheng *et al.*, (1991a,b) reported a sustained increase in serum levels of IL-1, TNF and IL-6 in patients treated with either DEC or Ivermectin.

Antigen levels usually rise immediately during, and immediately after treatment peaking but there is considerable individual variation (Ramaprasad *et al.*, 1988; Eberhard *et al.*, 1997). Most authors record a fall in antigen levels in the months following treatment. For instance Weil *et al.*, (1988) found that filarial antigen fell to 72%, 58% 53% and 48% at 1, 3, 6 and 12 months after treatment respectively. A rise in filarial antigen levels is associated with the re-appearance of microfilariae in the patient's blood (Ramaprasad *et al.*, (1988). Nicolas *et al.*, (1997) observed that there was a difference in the clearance of antigen and resurgence of antigenaemia between amicrofilaremic and filaremic subjects when they were treated with an annual single-dose treatment of DEC, Ivermectin or a combination of both. Their results are summarised in Table 7.

Ramaprasad *et al.*, (1988) found that like filarial antigen, filarial antigen associated immune complexes initially increased then fell, reaching a steady state by about day 60 after treatment.

Table 7. Effects of Diethylcarbamazine (DEC) and Ivermectin (IVR) alone and in combination on filarial antigen in microfilaremic (mf+) and microfilaremic (mf-) subjects (from Nicolas *et al.*, 1997)

	DEC 6 mg/kg	IVR 400 µg/kg	DEC 6 mg/kg IVR 400 µg/kg	DEC 3 mg/kg IVR 400µg/kg
% residual antigen level at 12 mths	55.5 mf+	70.0 mf+	60.0 mf+	52.3 mf+
	21.4 mf-	0.0 mf-	19.2 mf-	7.1 mf-
% residual antigen level at 24 mths	37.6 mf+	63.3 mf +	41.0 mf+	27.0 mf+
	25.0 mf-	0.0 mf -	44.4 mf-	17.8 mf-

Conclusion

The last few years has seen a virtual explosion in both basic and applied filariasis research. We now have a better understanding of the immunological basis of infection and the pathogenesis of filarial disease. We now know that opportunistic bacterial and fungal infections play an active part in the progression to chronic obstructive filarial disease and that attention to basic hygiene, the use of antibiotics and antifungals and local physiotherapy can slow, prevent, or in some cases, reverse elephantiasis.

The diagnosis of bancroftian filariasis has been revolutionised by the introduction of filarial antigen tests that are not dependent on the presence of microfilariae, removing the need to take night bloods. The ICT test can be done by finger-stick in the field and results are rapidly available. Ultra-sound is proving to be a valuable tool for identifying adult worms *in-situ*. Annual mass treatment with Diethylcarbamizine (DEC) either alone or in combination with Ivermectin and Albendazole has been proven to be very effective in destroying microfilariae and interrupting transmission. With this knowledge and armed with these tools, the WHO goal of elimination lymphatic filariasis as a health threat by the year 2020 is achievable. Despite these advances there is still work to be done and many questions to be answered. There is still a lot of “traditional dogma” about filariasis that needs to be re-evaluated by modern research methods

A fundamental question is, “What is the burden of disease?”. A conservative estimate is around 160 million cases, but no one really knows because a large part of filarial-endemic areas remain un-surveyed. Geographic Information Systems (GIS) and other spacial analysis tools need to be employed so that the geographic distribution of filariasis can be accurately determined. Ultrasound and antigen tests have shown that many “endemic normals” have low-grade patent infection and many of those with asymptomatic microfilaraemia have lymphatic damage, varying degrees of renal impairment and immuno depression. It makes good sense to define a filariasis case as all those who have evidence of active infection (i.e. those with a positive filarial antigen test regardless of microfilaraemia status or presence or absence of chronic pathology). Clinical diagnosis of acute filariasis is often difficult and can be confused with other febrile inflammatory illnesses. There is thus a need to develop and trial clinical checklist that will improve diagnosis. The role of the filarial parasite *per se* and the contribution of other infectious agents, in particular streptococci, to acute filarial attacks need to be investigated and treatment regimes devised.

There are many other questions. Most filarial research has neglected children. What level of disease and

pathology can be expected in children of different ages? What effect does filarial-induced immuno suppression have on the efficacy of vaccinations? What part does filarial-induced immuno suppression play in host susceptibility to other parasites and other infectious agents especially tuberculosis? There is some anecdotal and some limited experimental evidence of renal disease in filariasis, but there needs to be a detailed assessment of the contribution that filariasis makes to the burden of renal disease in tropical populations and the amount of long-term kidney damage that is prevented by filariasis control programs. There have been some studies on the social effects of disfiguring filarial disease in Africa and India, but very little has been done in other endemic areas. There is an urgent need to develop simple, effective health promotion materials to inform people about the disease and its prevention. Given the link between secondary infection and the development of chronic pathology special education materials on basic limb hygiene must be made available to those who are at risk of developing such complications.

Filarial control can only be achieved if the majority of the population takes the drugs. In many endemic areas the health system is already struggling to cope with programs already in place and are unable to cope with the extra impost of a filariasis control program. There must be research into innovative, cost-effective means of drug delivery. How can non-governmental organisations such as religious organisations and missions, schools and industry be actively involved in filariasis control? In a country like Papua New Guinea, for example, where a large percentage of the population attend church, it would be possible for the church eldership, with local health authority support, to distribute anti-filarial drugs at major church festivals and obtain excellent population coverage. The use of Albendazole raises the possibility of not only controlling filariasis, but a range of intestinal parasites as well, making such a control project attractive to health authorities. Pregnant and lactating women have been excluded from drug treatment because of safety concerns, but hard evidence of risk is lacking and the matter requires investigation. Although the drugs presently available appear to be very effective, there is still a great need to discover other anti-filarial drugs, especially effective macrofilaricides. Although there is at present, no real hard evidence of drug resistance, that does not mean that it will not occur in the future, and techniques to monitor the emergence of drug resistance need to be developed and trialed. As can be seen in the foregoing, much progress has been made in filariasis research and control, but a large amount of work remains before the scourge of filariasis is finally eliminated.

References

1. Abalmasov KG, Egarov YS, Abramov YA, Chatterjee SS, Uvarov DL, Neiman VA. Evaluation of greater omentum in the treatment of experimental lymphedema. *Lymphology* 1994; 27:129-136.
2. Abbasi I, Hamburger J, Githure J, Ochola JJ, Agure R, Koech DK, Ramzy R, Gad A, Williams S. Detection of *Wuchereria bancrofti* DNA in patient's sputum by the polymerase reaction. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1996; 90:531-532.
3. Abbasi I, Githure J, Ochola JJ, Agure R, Koech DK, Ramzy RM, Williams SA, Hamburger J. Diagnosis of *Wuchereria bancrofti* infection by the polymerase chain reaction employing patients' sputum. *Parasitology Research* 1999; 85:844-849.
4. Adamolekun B. The aetiologies of epilepsy in tropical Africa. *Tropical and Geographical Medicine* 1995; 47:115-117.
5. Addiss DG, Eberhard ML, Lammie PL. Comparative efficiency of clearing-dose and single high-dose ivermectin and diethylcarbamazine against *Wuchereria bancrofti* microfilaraemia. *American Journal of Tropical Medicine and Hygiene* 1993; 48:178-185.
6. Addiss DG, Eberhard ML, Lammie PL. "Filarial" adenolymphadenitis without filarial infection. *Lancet* 1994; 343:597.
7. Addiss DG, Dimock KA, Eberhard ML, Lammie PL. Clinical Parasitologic and immunologic observations of patients with hydrocele and elephantiasis in a area with endemic lymphatic filariasis. *The Journal of Infectious Diseases* 1995; 171:755-758.
8. Addiss DG, Louis-Charles J. Feasibility and effectiveness of treatment with filaria-associated lymphedema in Haiti: preliminary results. *American Journal of Tropical Medicine and Hygiene* 1997; 57 (supplement):215 (meeting abstract).
9. Addiss DG, Beach MJ, Streit TG, Lutwick S, LeConte FH, Lafontant JG, Hightower AW, Lammie PJ. Randomised placebo-controlled comparison of ivermectin and albendazole alone and in combination for *Wuchereria bancrofti* microfilaraemia in Haitian children. *Lancet* 1997; 350:480-484.
10. Addiss D. Lymphatic filariasis. *Bulletin of the World Health Organisation* 1998; 76:Supp 145-148.
11. Adebajo AO. Rheumatic manifestations of tropical diseases. *Current Opinions in Rheumatology* 1996; 8:85-89.
12. Adhikari P, Haldar S, Ghosh NR, Mandal MM, Haldar JP. Prevalence of Bancroftian filariasis in Burdwan district, west Bengal: a comparative study between colliery and non-colliery areas. *Journal of Communicable Diseases* 1994; 26:6-13.
13. Agarwal M, Prasad GBKS, Harinath BC, Bhatia BD. Transplacental transfer of filarial infection. *Indian Pediatrics* 1986; 23:169-174.
14. Agarwal R, Bhatnagar SK, Chandra R. Lymphosuction: a new treatment for chronic filarial lymphedema. *European Journal of Plastic Surgery* 1998; 21:113-117.
15. Ahorlu CK, Dunyo SK, Simonsen PE. Scarification as a risk factor for rapid progression of filarial elephantiasis. *Lymphatic Filariasis Research and Control in Africa: report on a workshop held in Tanga, Tanzania. Danish Bilharziasis Laboratory, Denmark & National Institute for Medical Research, Tanzania*, 1997.
16. Ahorlu CK, Dunyo SK, Simonsen PE. Lymphatic filariasis-related perceptions and practices on the coast of Ghana: implications for prevention and control. *Acta Tropica* 1999; 73:251-261.
17. Akue JP, Egwang TG, Devaney E. High levels of parasite-specific IgG4 in the absence of microfilaraemia in *Loa loa* infection. *Tropical Medicine & Parasitology* 1994; 45:246-248.
18. Albuquerque MFM, Marzochi MC, Ximenes RAA, Braga MC, Silva MCM, Furtado AF. Bancroftian filariasis in two urban areas of Recife, Brazil: the role of individual risk factors. *Revista Instituto de Medicina Tropical de Sao Paulo* 1995; 37:225-233.
19. Alexander NDE, Kazura JW, Bockarie MJ, Perry RT, Dimber ZB, Grenfell BT, Alpers MP. Parental infection confounded with local infection intensity as risk factors for childhood microfilaraemia in Bancroftian filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1998; 92:23-24.
20. Alexander NDE (a), Perry RT., Perry RT, Dimber ZB, Hyun PJ, Alpers MP, Kazura JW. Acute disease episodes in a *Wuchereria bancrofti*-endemic area of Papua New Guinea. *American Journal of Tropical Medicine and Hygiene* 1999; 61:319-324.
21. Alexander ND (b), Grenfell BT. The effect of pregnancy on *Wuchereria bancrofti* microfilarial load in humans. *Parasitology* 1999; 119:151-156.
22. Alhadeff R. Clinical aspects of filariasis. *Journal of Tropical Medicine and Hygiene* 1955; 58:173-179.
23. Allen JE, Macdonald AS. Profound suppression of cellular proliferation mediated by the secretions of nematodes. *Parasite Immunology* 1998; 20:241-247.
24. Almeida AJ, Bhopale MK, Renapurkar DM. Evaluation of *Setaria cervi* antigen in the diagnosis of human filariasis. *Folia Parasitologica* 1990; 37:323-329.
25. Amaral F, Dreyer G, Figueredo-Silva J, Noroes J, Cavalcanti A, Samico SC, Santos A, Coutinho A. Live adult worms detected by ultrasonography in human Bancroftian filariasis. *American Journal of Tropical Medicine & Hygiene* 1994; 50:753-757.
26. Amazigo UO, Obikeze DS. Social consequences of onchocercal skin lesions on adolescent girls in rural Nigeria. *World Health Organisation/TDR Discussion Paper, World Health Organisation Geneva* 1992.
27. Ambroise-Thomas P. Immunological diagnosis of human filariasis: present possibilities, difficulties and limitations - review. *Acta Tropica* 1974; 31:108-128.
28. Ambroise-Thomas P. Filariasis. In: Houba V. *Immunological Investigation of Tropical Parasitic Diseases*. Churchill Livingstone, London 1980. American Society of Tropical Medicine and Hygiene. Report of a symposium on helminthic infections. *American Journal of Tropical Medicine and Hygiene* 1957; 6:402.
29. Amuyunzu M. Community perception regarding chronic filarial swellings - A case study of the Duruma of coastal Kenya. *East African Medical Journal* 1997; 74:411-415.
30. Anders RF (a), Price MA, Wilkey I. Geographical distribution of amyloidosis in Papua New Guinea. *Tropical and Geographical Medicine* 1976; 28:273-276.
31. Anders RF (b), Natvig JB, Michaelsen TE, Husby G. Isolation of amyloid-related serum protein SAA as a low molecular weight protein. *Scandinavian Journal of Immunology* 1976; 4:397-342.
32. Anderson RM, May RM. *Infectious diseases of humans, dynamics and control*. pages 576-589. Oxford University Press 1991.

33. Andreano R, Helminiak T. Economics, health & tropical diseases. In: Economics, Health & Tropical Diseases, pages 19-72, published by School of Economics University of the Philippines 1988.
34. Andrews EW. Dzibilchaltun; lost city of the Maya. National Geographic 1959; 115:90-119.
35. Andy JJ, Ogunowo PO, Akpan NA, Odigwe CO, Ekanem IA, Esin RA. Helminth associated hyper-eosinophilia and tropical endomyocardial fibrosis (EMF) in Nigeria. Acta Tropica 1998; 69:127-140.
36. Antani J, Krishnamurthy KR. Filarial haematuria. Journal of the Association of Physicians of India 1971; 19:59-60.
37. Ash LR, Little MD. *Brugia beaveri* sp. n. (Nematoidia: Filarioidea) from the Raccoon (*Procyon lotor*) in Louisiana. Journal of Parasitology 1971; 57:1043-1051.
38. Ashburn PM, Craig CF. A new blood filaria of man: *Filaria philippinensis*. American Journal of Medical Science 1905; 132: 435-443.
39. Ata AHA, El-Bassiouny AEI, Shaker ZA, Attia MM, El-Khashab N, El-Gammal NE, Mourad AMRA, El-Ghoneimy S. Antifilarial IgM versus IgG antibody determination in the diagnosis of *Wuchereria bancrofti* infection in Egyptians. Journal of the Egyptian Society of Parasitology 1993; 23:277-283.
40. Atmadja AK, Atkinson R, Sartono E, Partono F, Yazdanbakhsh M, Maizels RM. Differential decline in filaria-specific IgG1, IgG4 and IgE antibodies in *Brugia malayi*-infected patients after diethylcarbamazine chemotherapy. Journal of Infectious Diseases 1995; 172:1567-1572.
41. Au AC, Denham DA, Steward MW, Draper CC, Ismail MM, Rao CK, Mak JW. Detection of circulating antigens and immune complexes in feline and human lymphatic filariasis. Southeast Asian Journal of Tropical Medicine & Public Health 1981; 12:492-498.
42. Ayers M, Salzano FM, Helena M, Franco LP, De Souza Barros RM. The association of blood groups, ABH secretion, haptoglobins and haemoglobins with filariasis. Human Heredity 1976; 26:105-109.
43. Azimi PH, Grossman M. Your diagnosis please - a thirteen year old with a painful swollen foot and eosinophilia. Pediatric Infectious Disease Journal 1996; 14:820-826.
44. Azoubel E. The clinical manifestations of Bancroftian filariasis. Lymphology 1996; 29 (supplement):12-19
45. Baird JK, Alpert LI, Friedman R, Schraft WC, Connor DH. North American Brugian filariasis: report of nine infections in humans. American Journal of Tropical Medicine and Hygiene 1986; 35:1205-1209.
46. Bal M, Das MK. Antigenicity of filarial superoxide dismutase in human bancroftian filariasis. Journal of Biosciences (Bangalore) 1995; 20:157-166.
47. Bal M, Das MK. Glutathione-binding proteins of *Setaria digitata*: antibody responses in humans infected with *Wuchereria bancrofti*. Parasite Immunology 1996; 18:473-477.
48. Bal M, Das MK. Antibody response to a filarial antigen fraction in individuals exposed to *Wuchereria bancrofti* infection in India. Acta Tropica 1999; 72:259-274.
49. Bandyopadhyay L. Lymphatic filariasis and the women of India. Social Science Medicine 1996; 42:1401-1410.
50. Barsoum RS. Tropical parasitic nephropathies. Nephrology Dialysis and Transplantation 1999; 14 (supp 3):79-81
51. Beach MJ, Streit TG, Addiss DG, Prospere R, Roberts M, Lammie PJ. Assessment of combined ivermectin and albendazole for treatment of intestinal helminth and *Wuchereria bancrofti* infections in Haitian school children. American Journal of Tropical Medicine and Hygiene 1999; 60:479-486.
52. Bean B, Ellman MH, Kagan IG. Acute lymphatic filariasis in an American traveller. Diagnostic Microbiology and Infectious Diseases 1992; 15:345-347.
53. Beaver PC, Orihel TC. Human infection with filariae of animals in the United States. American Journal of Tropical Medicine & Hygiene 1965; 14:1010-1030
54. Beaver PC. Filariasis without microfilaraemia. American Journal of Tropical Medicine & Hygiene 1970; 19:181-189.
55. Beaver PC, Wong MM. *Brugia* species from a cat in California. Proceedings of the Helminthological Society of Washington 1988; 55:111-113.
56. Beckett EB. Some quantitative aspects of flight muscle damage in mosquitoes infected with filarial larvae. Annals of Tropical Medicine and Parasitology 1973; 67:455-466.
57. Behara A, Atiri A, Srinivas C, Ravinda N, Gupta NM, Katariya RN, Jain D. Traumatic rupture of filarial spleen. Indian Journal of Gastroenterology 1995; 14:114.
58. Behm AW, Hayman JM. The course of filariasis after removal from an endemic area. American Journal of the Medical Sciences 1946; 211:386-394.
59. Beisel WR. Synergism and antagonism of parasitic diseases and malnutrition. Review of Infectious Diseases 1982; 4:798-804.
60. Bell D. Membrane filters and microfilariae: a new diagnostic technique. Annals of tropical Medicine and Parasitology 1967; 61:220-223.
61. Bennett IC, Furnival CM, Searle J. Dirofilariasis in Australia: unusual cause of a breast lump. Australia and New Zealand Journal of Surgery 1989; 59:671-673.
62. Binoy C, Govardhana Rao Y, Ananthakrishnan N, Kate V, Yuvarai J, Pani SP. Omentoplasty in the management of filarial lymphodema. Transactions of the Royal Society of Tropical Medicine and Hygiene 1998; 92:317-319.
63. Bernas M, Andrade M, Azoubel E, Witte M. Summary of special course on Bancroftian filariasis, Recife, Brazil, September 25, 1995. Lymphology 1996; 29 (supplement):9-12.
64. Betschart B, Marti S, Glaser M. Antibodies against the cuticle of *Ascaris suum* cross-react with epicuticular structures of filarial parasites. Acta Tropica 1990; 47:331-338.
65. Beuria MK, Bal M, Das MK. Allergic reactivity and IgG subclasses to a proteinase fraction of *Seratia digitata* in filariasis. Journal of Helminthology 1995; 69:181-185.
66. Bhatia B, Misra S. Recurrent Guillain-Barre syndrome following acute filariasis. Journal of Neurology, Neurosurgery and Psychiatry 1993; 56: 1133-1134.
67. Bhattacharya C, Singh RN, Misra S, Rathaur S. Diethylcarbamazine: effect on lysosomal enzymes and acetylcholine in *Wuchereria Bancroftian* infection. Tropical Medicine and International Health 1997; 2:686-690.
68. Bianco AE, Luty A, Whitworth J, Taylor D. Immunity to *Onchocerca volvulus* in mice and the induction of cross-protection with *Onchocerca lienalis*. Tropical Medicine and Parasitology 1991;42:188-190.
69. Blacksin MF, Lin SS, Trofa AF. Filariasis of the ankle: magnetic resonance imaging. Foot and Ankle

International 1999; 20:738-740.

70. Bloch P, Simonsen PE, Weiss N, Nutman TB. The significance of Guinea worm infection in the immunological diagnosis of onchocerciasis and Bancroftian filariasis. Bloomfield RD, Suarez JR, Malangit AC. Transplacental transfer of Bancroftian filariasis. Journal of the National Medical Association 1978; 70:597-598.
71. Bockarie MJ, Paru R. Forest clearing, mosquitoes and malaria. Report of the Jubilee International Colloquium, Papua New Guinea Institute of Medical Research, Madang, 1993.
72. Bockarie MJ. Transmission intensity and its relationship to infection and disease. Lymphatic Filariasis Research and Control in Africa: report on a workshop held in Tanga, Tanzania. Danish Bilharziasis Laboratory, Denmark & National Institute for Medical Research, Tanzania, 1997.
73. Bockarie MJ, Hii JLK, Alexander NDE, Bockarie F, Dagoro JW, Kazura JW, Alpers MP. Mass treatment with ivermectin for filariasis control in Papua New Guinea: impact on mosquito survival. Medical and Veterinary Entomology 1999; 13:120-123.
74. Bolarin DM. Serum ferritin in Nigerian patients with amoebic liver abscess and other tropical infections with liver involvement. Acta Tropica 1983; 40:159-164.
75. Bosshardt SC, Coleman SU, McVay CS, Jones KL, Klei TR. Impact of microfilaraemia on maintenance of a hypo-responsive cellular immune response in *Brugia*-infected Gerbils (*Merionesunguiculatus*). Infection and Immunity 1995; 63:940-945.
76. Bosworth W, Ewert A, Flores Ae. Antistreptolysin titres of cats infected with *Brugia malayi* and streptococci. Southeast Asian Journal of Tropical Medicine & Public Health 1977; 8:255-259.
77. Boudreaux MK, Dillion AR, Spanos JS. Enhanced platelet reactivity in heart worm-infected dogs. American Journal of Veterinary Research 1989; 50:1544-1547.
78. Boudreaux MK, Dillion AR. Platelet function, antithrombin-111 activity, and fibrinogen concentration in heart worm-infected and heart worm-negative dogs treated with thiacetarsamide. American Journal of Veterinary Research 1991; 52:1986-1991.
79. Bourne AG. A note on *Filaria sanguinis hominis*: with a description of the male specimen. British Medical Journal 1888; 1:1050-1051.
80. Bourne AG. A note on *Filaria sanguinis hominis*: with a description of the male specimen. British Medical Journal 1888; 1:1050-1051.
81. Bowers JZ, Carruba RW. The doctoral thesis of Englebert Kaempfer on tropical diseases, oriental medicine and exotic natural phenomena. Journal of the History of Medicine 1970; 25:270-310.
82. Bozicevich J, Hutter AM. Intradermal and serological tests with *Dirofilaria immitis* antigen in cases of human filariasis. American Journal of Tropical Medicine 1944; 24:203-208.
83. Brabin L. Sex differentials in susceptibility to lymphatic filariasis and implications for maternal immunity. Epidemiology and Infection 1990; 105:335-353.
84. Brito AC, Fontes G, Williams P, Rocha EM. Bancroftian filariasis in Maceio, state of Alagoas, Brazil: Observations on *Culex quinquefasciatus* after blood feeding on individuals with different densities of microfilariae in the peripheral blood stream. American Journal of Tropical Medicine and Hygiene 1998; 58:489-494.
85. Brown JD, Armstrong DR. Filariasis in indigenous peoples and American servicemen in Vietnam. Annual Progress Report of United States Army Medical Research Team and Institute Pasteur Vietnam September 1967- June 1968: 276-290.
86. Brug SL. Een nieuwe *Filaria*-soort (*Filaria malayi*) parasiterendi bij den mensch (voorloopige mededeeling). Geneeskundig Tijdschrift voor Nederlandsch-Indie 1927; 67:750-754.
87. Brug SL. Filariasis in Nederlandsch-Indie. Geneeskundig Tijdschrift voor Nederlandsch-Indie 1928; 68:681-704.
88. Brug SL, de Rook H. Filariasis in Nederlandsch-Indie ii. De overbrenging van *Filaria malayi*. Geneeskundig Tijdschrift voor Nederlandsch-Indie 1930; 70:451-472.
89. Brunetiere A. Le filaire de loeil (*filaria loa*) peut-elle determiner des complications cerebrales? Gazette Hebdomadaire des Sceinces Medicales 1913; 30:351-354.
90. Bryan JH, (a) Southgate BA. Factors affecting transmission of *Wuchereria bancrofti* by anopheline mosquitoes 1. Uptake of microfilariae. Transactions of the Royal Society of Tropical Medicine and Hygiene 1988; 82:128-127.
91. Bryan JH, Southgate BA. (b) Factors affecting transmission of *Wuchereria bancrofti* by anopheline mosquitoes 2. Damage to ingested microfilariae by mosquito fore gut armatures and development of filarial larvae in mosquitoes. Transactions of the Royal Society of Tropical Medicine and Hygiene 1988; 82:138-145.
92. Bryan JH, McMahon P, Barnes A. Factors affecting transmission of *Wuchereria bancrofti* by anopheline mosquitoes 3. Uptake and damage to ingested microfilariae by *Anopheles gambiae*, *An arabiensis*, *An merus* and *An funestus* in East Africa. Transactions of the Royal Society of Tropical Medicine and Hygiene 1990; 84:265-268.
93. Buckley JJ. A new genus, *Brugia*, for *Wuchereria* spp. of the *malayi* group. Abstracts of the Papers of the Sixth International Congress of Tropical Medicine and Malaria, Lisbon 1958, page 35.
94. Buckley JJ. On *Brugia* for *Wuchereria* species of the "malayi" group i.e. *W. malayi* Brug 1927, *W. pahangi* Buckley and Edeson 1956, and *W. patei* Buckley, Nelson and Heisch 1958. Annals of Tropical Medicine and Parasitology 1960; 54:75-77.
95. Burhans RA, Camp JD, Butt HR, Cragg RW. Lymphangitis of suspected filarial origin; preliminary report concerning its treatment. United States Naval Medical Bulletin 1944; 42:336-340.
96. Burkot TR, Kwan-Lim GE, Maizels RM. A novel 95-kilodalton antigen of *Wuchereria bancrofti* infective larvae identified by species specific monoclonal antibodies. Infection and Immunity 1996; 64:485-488.
97. Burri H, Loutan L, Kumaraswami V, Vijayasekaran V. Skin changes in chronic lymphatic filariasis. Transactions of the Royal Society of Tropical Medicine & Hygiene 1996; 90:671-674.
98. Burton GJ. Observations on filarial larvae in bedbugs feeding on human carriers of *Wuchereria bancrofti* and *Brugia malayi* in south India. American Journal of Tropical Medicine and Hygiene 1962; 11:69-75.
99. Burton GJ. Natural and experimental infection of bedbugs with *Wuchereria bancrofti* in British Guiana. American Journal of Tropical Medicine and Hygiene 1963; 12:541-547.
100. Cahill KM. Filarial chyluria: a biochemical and radiological study of five patients. Journal of Tropical Medicine and Hygiene 1965; 68:27-31.

101. Campbell WC, Rew RS. Chemotherapy of Parasitic Diseases. Plenum Press, New York 1986.
102. Campbell WC. Ivermectin as an anti-parasite agent for use in humans. Annual Review of Microbiology 1991; 45:445-474.
103. Campbello TR, Santos Ferreira R, Pires ML, Gouveia de Melo P, Albuquerque R, Araujo S, Dreyer G. A study of placentas from *Wuchereria bancrofti* microfilaraemic and amicrofilaraemic mothers. Journal of Tropical Medicine and Hygiene 1993; 96:251-255.
104. Cao W, Van der Ploeg CPB, Plaisier AP, Sivera van der Sluijs IJ, Habbema JDF. Ivermectin for the chemotherapy of Bancroftian filariasis: a meta-analysis of the effect of a single treatment. Tropical Medicine and International Health 1997; 2:393-403.
105. Carlier Y, Truyens C. Influence of maternal infection on offspring resistance towards parasites. Parasitology Today 1995; 11:94-99.
106. Carne B, Utahia A, Tuira E, Teuru T. Filarial elephantiasis in French Polynesia: a study concerning the beliefs of 127 patients about the origins of their disease. Transactions of the Royal Society of Tropical Medicine & Hygiene 1979; 73:424-426.
107. Cartel JL, Spiegel A, Nguyen-Ngoc L, Cardines R, Plichart R, Martin PMV, Roux JF, Moulia-Pelat JP. Compared efficiency of repeated annual and semi-annual doses of ivermectin and diethylcarbamazine for prevention of *Wuchereria bancrofti* filariasis in French Polynesia: final evaluation. Tropical Medicine and Parasitology 1992; 43:91-94.
108. Casey HW, Obeck DK, Splitter GA. Immunopathology studies on canine heart worm disease. In: Bradley RE & Pacheco G. Canine Heart worm Disease: the Current Knowledge. University of Florida Press, Gainesville 1972.
109. Caley-Smith JR, Jamas S, Caley-Smith Judith R. Reduction of filariid lymphoedema and elephantiasis by 5,6 benzO-a-pyrone (coumarin) and the effects of diethylcarbamazine (DEC). Annals of Tropical Medicine and Parasitology 1993; 87:247-258.
110. Castellani A. Researches on elephantiasis nostras and elephantiasis tropica with special regard to their initial stage of recurring lymphangitis. Journal of Tropical Medicine & Hygiene 1969; 72:89-91.
111. Cesbron JY, Capron A, Vargaftig BB, Lagarde M, Pincemail J, Braquet P, Taelman H, Joseph M. Platelets mediate the action of diethylcarbamazine on microfilariae. Nature 1987; 325:533-537.
112. Chakravarty SN, Gupta HB. Filarial pericarditis. Journal of the Indian Medical Association 1966; 46:503-504.
113. Chan MS, Srividya A, Norman RA, Pani SP, Ramaiah KD, Vamamail P, Michael E, Das PK, Bundy DAP. EPIFIL- a dynamic model of infection and disease in lymphatic filariasis. American Journal of Tropical Medicine and Hygiene 1998; 59:606-614.
114. Chan SH, Dissanayake S, Mak JW, Ismail MM, Wee GB, Srinivasan NBH, Zaman V. HLA and filariasis in Sri Lankans and Indians. South East Asian Journal of Tropical Medicine and Public Health 1984; 15:281-286.
115. Chandrasekharan A, Pillai KK, Cherian C, Roy Chowdhury SP, Rao CK. Control of *Brugia malayi* filariasis with common salt fortified with DEC in some hill tribe settlements in Kerala. Indian Journal of Medical Research 1984; 79:600-603.
116. Chandrashekar R, Rao UR, Rajasekariah GR, Subahmanyam D. Separation of viable microfilariae free of blood on percoll gradients. Journal of Helminthology 1984; 58:69-70.
117. Chanteau S, Plichart R, Spiegel A, Martin PMV, Cartel MJL. Diagnostic values of ELISA-IgG4 as compared to ELISA-IgG and indirect immuno-fluorescence for the routine diagnosis of Bancroftian filariasis in the South Pacific: application on capillary blood collected on filter paper. Tropical Medicine & Parasitology 1991; 42:339-342.
118. Chanteau S, Cartel JL, Martin PMV. IgA immuno-assay for the diagnosis of bancroftian filariasis. Tropical Medicine and Parasitology 1992; 43:132-133.
119. Chanteau S (a), Moulia-Pelat JP, Glaziou NL. Og4C3 circulating antigen: a marker of infection and adult worm burden in *Wuchereria bancrofti* filariasis. Journal of Infectious Diseases 1994; 170:247-250.
120. Chanteau S (b), Glaziou P, Luquaud P, Plichart C, Moulia-Pelat JP, Cartel JL. Og4C3 circulating antigen, anti-*Brugia malayi* IgG and IgG4 titres in *Wuchereria bancrofti* infected patients, according to their parasitological status. Tropical Medicine and Parasitology 1994; 45:255-257.
121. Chanteau S (c), Luquaud P, Failloux AB, Williams SA. Detection of *Wuchereria bancrofti* larvae in pools of mosquitoes by the polymerase chain reaction. Transactions of the Royal Society of Tropical Medicine and Hygiene 1994; 88:665-666.
122. Chanteau S, Glaziou P, Plichart C, Luquaud P, Moulia-Pelat JP, Cartel JL. *Wuchereria bancrofti* filariasis in French Polynesia: age-specific patterns of microfilaraemia, circulating antigen, specific IgG and IgG4 responses according to transmission level. International Journal for Parasitology 1995; 25:81-85.
123. Charles SX. Filarial acute abdomen in pregnancy. Journal of the Indian Medical Association 1976; 66:302-303.
124. Charlwood JD, Dagoro H. Impregnated bed nets for the control of filariasis transmitted by *Anopheles punctulatus* in rural Papua New Guinea. Papua New Guinea Medical Journal 1987; 30:199-202.
125. Charters AD. Tropical infections contracted in Papua New Guinea and imported into Western Australia. Papua New Guinea Medical Journal 1981; 24:74-79.
126. Chaturvedi P, Gawdi AM, Parkhe K, Harinath BC, Dey SK. Arthritis in children: an occult manifestation of Bancroftian filariasis. Indian Journal of Paediatrics 1993; 60:803-807.
127. Chaturvedi P, Harinath BC, Reddy MVR, Advani B, Gawdi A, Alikhan A, Rama Rao BV. Filarial antibody detection in suspected occult filariasis in children in an endemic area. Journal of Tropical Paediatrics 1995; 41:243-245.
128. Cheirmaraj K, Reddy MVR, Harinath BC. Diagnostic use of polyclonal antibodies raised in mouse ascitic fluid in Bancroftian filariasis. Journal of Immuno-assay 1990; 11:429-444.
129. Cheirmaraj K, Parkhe KA, Harinath BC. Evaluation of fractionated circulating filarial antigen in diagnosis of Bancroftian filariasis. Journal of Tropical Medicine & Hygiene 1992; 95:47-51.
130. Chen SN, Howells RE. The *in-vitro* cultivation of the infective larvae and the early mammalian stages of the filarial worm *Brugia pahangi*. Annals of Tropical medicine and Parasitology 1979; 73:473-485.
131. Chenthamarakshan V, Reddy MVR, Harinath BC. Immuno-prophylactic potential of a 120kDa *Brugia malayi* adult antigen fraction, BMa-2, in lymphatic filariasis. Parasite Immunology 1995; 17:277-285.
132. Cheong WH. Vectors of Filariasis in Malaysia. In Mak J.W. Filariasis, Bulletin No.19, Institute for Medical Research Malaysia, Kuala Lumpur, 1983.

133. Chodakewitz J. Ivermectin and lymphatic filariasis: a clinical update. *Parasitology Today* 1995; 11:233-235.
134. Chugh KS, Singhal PC, Tewari SC, Nath IVS, Data BN. Acute glomerulonephritis associated with filariasis. *American Journal of Tropical Medicine & Hygiene* 1978; 27:630-631.
135. Chulerek P, Desowitz RS. A simplified membrane filtration technique for the diagnosis of microfilaraemia. *Journal of Parasitology* 1970; 56:23-26.
136. Clodius L. Lymphosuction: a new treatment modality for chronic filarial lymphedema. *European Journal of Plastic Surgery* 1998; 21:111-112.
137. Cobbold TS. Discovery of the adult representative of microscopic filaria. *Lancet* 1877; 2:70-71.
138. Coggeshall LT. Problems of filariasis. *Southern Medical Bulletin* 1945; 38:186-188.
139. Coggeshall LT. Filariasis in the serviceman: retrospect and prospect. *Journal of the American Medical Association* 1946; 131:8-12.
140. Colwell EJ, Armstrong DR, Brown JD, Duxbury RE, Sadun EH, Legters LJ. Epidemiologic and serologic investigations of filariasis in indigenous populations and American soldiers in South Vietnam. *American Journal of Tropical Medicine and Hygiene* 1970; 19:227-231.
141. Comley JCW, Turner CH. Potential of a soluble tetrazolium/formazan assay for the evaluation of filarial viability. *International Journal for Parasitology* 1990; 20:251-255.
142. Conn HC, Greenslit FS. Filariasis residuals in veterans with a report of microfilaraemia. *American Journal of Tropical Medicine and Hygiene* 1952; 474-476.
143. Cooper MJ, Espinel I, Paredes W, Guderian RH, Nutman TB. Impaired tetanus-specific cellular and humoral responses following tetanus vaccination in human oncocerciasis - a possible role for interleukin - 10. *Journal of Infectious Diseases* 1998; 178:1133-1138.
144. Cooke RA, Champness LT. Amyloidosis in Papua New Guinea. *Medical Journal of Australia* 1970; 2:1177.
145. Coreil J, Mayard G, Louis-Charles J, Addiss D. Filarial elephantiasis among Haitian women: social context and behavioural factors in treatment. *Tropical Medicine and International Health* 1998; 3:467-473.
146. Coutinho AD, Dreyer G, Medetros Z. Ivermectin treatment of Bancroftian filariasis in Recife, Brazil: a two-year comparative study of the efficiency of single treatments with ivermectin or diethylcarbamazine. *American Journal of Tropical Medicine and Hygiene* 1994; 50:339-348.
147. Crandall R, Crandall C, Doyle T, Nayar J. Immune responses and chronic lymphedema in experimental filariasis. *Lymphology* 1993; 26:120-127.
148. Cross JH, Partono F, Hsu MYK, Ash LR, Oemijati S. Experimental transmission of *Wuchereria bancrofti* to monkeys. *American Journal of Tropical Medicine & Hygiene* 1979; 28:56-66.
149. Crowell WA, Malone JB, Chapman WL, Thompson PE. Amyloidosis in *Litomosoides carinii*-infected Mongolian gerbils (*Meriones unguiculatus*). *Journal of the American Veterinary Association* 1973; 163:596-598.
150. Cryz SJ Jr. Molecular parasitology progress towards the development of vaccines for malaria, filariasis and schistosomiasis. *Experientia* 1991; 47:146-151.
151. Das Bidyut K, Sahoo PK, Ravindran B. A role for tumour necrosis factor- α in acute lymphatic filariasis. *Parasite Immunology* 1996; 18:421-424.
152. Das MK, Subramanyam VR, Ravindran B, Pattnaik NM. A study of the antigen, antibody and immuno-complex levels in *Wuchereria bancrofti* filariasis with reference to clinical status. *Journal of Tropical Medicine & Hygiene* 1987; 90:135-141.
153. Das MK, Beuria MK, Dash AP. Immunoglobulin E and G4 antibodies to infective larvae in a *Wuchereria bancrofti* endemic area. *International Archives of Allergy and Immunology* 1992; 99:118-122.
154. Das Bidyut K, Sahoo Prakask K, Ravindran B. A role for tumour necrosis factor in acute lymphatic filariasis. *Parasite Immunology* 1996; 18:421-424.
155. Dasgupta A. Cursory overview of lymphatic filariasis. *Indian Journal of Pathology and Microbiology* 1984; 27:273-280.
156. Dasgupta A. Genetic aspects in filariasis. *Journal of Communicable Diseases* 1985; 17 (suppl 1):117-124.
157. Dasgupta A, Bala S, Data SN. Lymphatic filariasis in man: demonstration of circulation antigens in *Wuchereria bancrofti* infection. *Parasite Immunology* 1984; 6:341-348.
158. Dasgupta A, Bala S, Data SN. Lymphatic filariasis: lymphocyte sub-populations in *Wuchereria bancrofti* infection. *Journal of Communicable Diseases* 1986; 18:295-301.
159. Dasgupta A, Bala S, Data SN. Cellular unresponsiveness in patients with soluble circulating antigens in Bancroftian filariasis. *Indian Journal of Medical Research* 1987; 85-136-139.
160. da Silva Araujo AJ. Caso de chyluria, elefantia do escroto, escroto lymphatico, craw-craw e erisipela em um mesmo individuo, descobrimento da *Wuchereria filaria* na lympho do escroto. Tratamento pela electricidade com excelentes resultados. *Gazeta Medica da Bahia*, second series, 1877; 2:492-504.
161. Date A, Gunasekaran V, Kirubakaran MG, Shastry JCM. Acute eosinophilic glomerulonephritis with Bancroftian filariasis. *Postgraduate Medical Journal* 1979; 55:905-907.
162. Date A, Neela P, Shastry JCM. Membrano-proliferative glomerulonephritis in a tropical environment. *Annals of Tropical Medicine & Parasitology* 1983; 77:279-285.
163. de Almeida AB, de Silva MCM, Maciel MA, Freedman DO. The presence or absence of active infection, not clinical status, is most closely associated with cytokine responses in lymphatic filariasis. *Journal of Infectious Diseases* 1996; 173:1453-1459.
164. de Almeida AB, Freedman DO. Epidemiology and immunopathology of bancroftian filariasis. *Microbes and Infection* 1999; 1:1015-1022.
165. David HL, Edeson JF. Filariasis in Portuguese Timor, with observations of a new microfilariae found in man. *Annals of Tropical Medicine and Parasitology* 1965; 59:193-204.
166. de Sylva DS, Ismail MM, de Silva ID, de Silva P, Gooneratne D, Goonawardana S. Is filariasis a trigger for bronchial asthma? *Acta Paediatrica Japan* 1990; 32:164-168.
167. Day KP. The endemic normal in lymphatic filariasis: a static concept. *Parasitology Today* 1991; 7:341-343.
168. Day KP, Gregory WF, Maizels RM. Age-specific acquisition of immunity to infective larva in a Bancroftian filariasis endemic area of Papua New Guinea. *Parasite Immunology* 1991; 13:277-283.
169. Day KP, Grenfell B, Spark R, Kazura JW, Alpers MP. Age specific patterns of change in the dynamics of *Wuchereria bancrofti* infection in Papua New Guinea. *American Journal of Tropical Medicine and Hygiene*

- 1991; 44:518-527.
170. Deehan MR, Frame MJ, Parkhouse RME, Seatter SD, Reid SD, Harnett MM, Harnett W. A phosphorylcholine-containing filarial nematode-secreted product disrupts B lymphocyte activation by targeting key proliferative signalling pathways. *Journal of Immunology* 1998; 160:2692-2699.
171. Denham DA, Dennis DT, Ponnudurai T, Nelson GS, Guy F. Comparison of a counting chamber and thick smear method of counting microfilariae. *Transactions of the Royal Society of Tropical Medicine & Hygiene* 1971; 65:521-523.
172. Denham DA, McGreevy PB. Brugian filariasis. *Advances in Parasitology* 1977; 17:243-308.
173. Denham DA. Counting and identifying microfilariae: a laboratory manual. London School of Hygiene and Tropical Medicine, London 1979.
174. Denham DA, McGreevy PB, Suswillo RR, Rogers R. The resistance to re-infection of cats repeatedly inoculated with infective larvae of *Brugia phangi*. *Parasitology* 1983; 86:11-18.
175. Demarquay JN. Note sur une tumeur des bourses contenant un liquide laiteux (galacoele de Vidal) et renferment des petits etres vermiformes que l'on peut considerer comme des helminthes hematoides a l etat d'embryon. *Gazette Medicale de Paris* 1863; 18:665-667.
176. Dennis DT, McConnell E, White GB. Bancroftian filariasis and membrane filters: are night surveys necessary? *American Journal of Tropical Medicine and Hygiene* 1976; 25:257-262.
177. Desowitz R.S, Saave J.J, Sawada T. Studies of the immuno-epidemiology of parasite infection in New Guinea. I. Population studies in the relationship of a skin test to microfilaraemia. *Annals of Tropical Medicine and Parasitology* 1966; 60:257-264.
178. Desowitz RS (a), Southgate BA. Studies on filariasis in the Pacific: 2. The persistence of microfilaria in diethylcabamazine treated populations of Fiji and Western Samoa: diagnostic application of the membrane filter technique. *Southeast Asian Journal of Tropical Medicine and Public Health* 1973; 4:179-183.
179. Desowitz RS (b), Southgate BA, Mataika JU. Studies on filariasis in the Pacific: 3. Comparison of the stained blood film, counting chamber and membrane-filtration techniques for the diagnosis of *Wuchereria bancrofti* microfilaraemia in untreated patients in areas of low endemicity. *Southeast Asian Journal of Tropical Medicine and Public Health* 1973; 4:329-335.
180. Desowitz RS, Jenkins C, Anian G. Bancroftian filariasis in an isolated hunter-gather shifting horticulturist group in Papua New Guinea. *Bulletin of the World Health Organisation* 1993;71:55-58.
181. Devaney E, Howells RE. Culture systems for the maintenance and developement of microfilariae. *Annals of Tropical Medicine and Parasitology* 1979; 73:139-143.
182. Devasia A, Ponnaiya J, Gopalakrishnan G. Filarial cystitis with urethral obstruction. *Journal of Urology* 1998; 159:1637-1638.
183. Dhanda V, Das PK, Lal R, Srinivasan F, Ramaiah KD. Spread of lymphatic filariasis, re-emergence of leishmaniasis and the threat of babesiosis in India. *Indian Journal of Medical Research* 1996; 103:46-54.
184. Diallo S, Aziz MA, Ndir O, Badiane S, Bah IB, Gaye O. Dose-ranging study of ivermectin in the treatment of filariasis due to *Wuchereria bancrofti*. *Lancet* 1987; i:1030-1031.
185. Dickson JG, Huntington RW, Eichold S. Filariasis in defense force, Samoan Group. *United States Navy Medical Bulletin* 1943; 41:1240-1251.
186. Dimock KA, Eberhard ML, Lammie PJ. Th1-like antifilarial immune responses predominate in antigen-negative persons. *Infection and Immunity* 1996; 64:2962-2967.
187. Dissanayake S, Ismail MM. ELISA in the diagnosis of *Wuchereria bancrofti* infection in man: a technique for reducing cross reactivity. *Bulletin of the World Health Organisation* 1980; 58:655-657.
188. Dissanayake S, de Silva LVK. IgM antibody to filarial antigens in human cord blood: possibility of transplacental infection. *Transactions of the Royal Society for Tropical Medicine & Hygiene* 1980; 74:542-544.
189. Dissanayake S, Galahitiyawa SC, Ismail MM. Immune complexes in *Wuchereria bancrofti* infection in man. *Bulletin of the World Health Organisation* 1982; 60:919-927.
190. Dissanayake S, Xu M, Piessens WF. A cloned antigen for serological diagnosis of *Wuchereria bancrofti* microfilaraemia with day time blood samples. *Molecular and Biochemical Parasitology* 1992; 56:269-277.
191. Dissanayake S, Zheng H, Dreyer G, Xu M, Watawana L, Cheng G, Wang S, Morin P, Deng B, Kurniawan L, Vincent A, Piessens WF. *American Journal of Tropical Medicine & Hygiene* 1994; 50:727-734.
192. Dissanayake S (a), Watawana L, Piessens WF. Lymphatic pathology in *Wuchereria bancrofti* microfilaraemic infections. *Transactions of the Royal Society of Tropical Medicine & Hygiene* 1995; 89:517-521.
193. Dissanayake S (b), Perler FB, Xu M, Southworth MW, Yee CK, Wang S, Dreyer G, Watawana L, Kurniawan A, Furhrman JA, Piessens WF. Differential recognition of microfilarial chitinase, a transmission-blocking vaccine candidate antigen by sera from patients with brugian and Bancroftian filariasis. *American Journal of Tropical Medicine and Hygiene* 1995; 53:289-294.
194. Doherty JF, Grant AD, Bryceson ADM. Fever as the presenting complaint of travellers returning from the tropics. *Quarterly Journal of Medicine* 1995; 88:277-281.
195. dos Santos F. *Filaria bancrofti*, verificacao no Brazil dadescoberto de Bancroft, no Australia. *Progresso Medico Rio de Janeiro* 1877; 2:95-100.
196. Dreyer G, de Andrade L. Inappropriateness of the association of diphenhydramine with diethylcarbamazine for the treatment of lymphatic filariasis. *Journal of Tropical Medicine and Hygiene* 1989; 92:32-34.
197. Dreyer G, Ottesen EA, Galdino E, Andrade L, Rocha A, Medeiros Z, Moura I, Casimiro I, Beliz F, Coutinho A. Renal abnormalities in microfilaraemic patients with Bancroftian filariasis. *American Journal of Tropical Medicine and Hygiene* 1992; 46:745-751.
198. Dreyer G, Figueredo-Silva j, Andrade LD, Marchetti F, Coutinho A. Diagnosis, treatment and control of filariasis. *Parasite* 1994 1 (supplement 1):22-23.
199. Dreyer G, Coutinho AD, Miranda D. Treatment of Bancroftian filariasis in Recife, Brazil: a two-year comparative study of the efficiency of single treatments with ivermectin or diethylcarbamazine. *American Journal of Tropical Medicine and Hygiene* 1995; 89:98-102.
200. Dreyer G (a), Amaral F, Nor-oes J Nen A, Medeiros Z, Coutinho A, Addiss D. Direct assesment of the aduldicidal efficincy of a single dose of ivermectin in

- lymphatic filariasis. Transactions of the Royal Society of Tropical Medicine and Hygiene 1995; 89:441-443.
201. Dreyer G (b), Nor-oes J, Amaral F, Medeiros Z, Addiss D. A new tool to assess the adulticidal efficacy in vivo of antifilarial drugs for Bancroftian filariasis. Transactions of the Royal Society for Tropical Medicine and Hygiene 1995; 89:225-226.
202. Dreyer G (a), Addiss D, Nor-oes J, Amaral F, Rocha A, Coutinho A. Ultrasonic assessment of the adulticidal efficacy of repeat high-dose ivermectin in Bancroftian filariasis. Tropical Medicine and International Health 1996; 1:427-432.
203. Dreyer G (b), Brand AC, Amaral F, Medeiros Z, Addiss D. Detection by ultrasound of living adult *Wuchereria bancrofti* in the female breast. Memoirs of the Institute of Oswaldo Cruz 1996; 91:95-96.
204. Dreyer G (c), Santos A, Noroes J, Rocha A, Addiss D. Amicrofilaraemic carriers of *Wuchereria bancrofti*. Transactions of the Royal Society for Tropical Medicine and Hygiene 1996; 90:288-289.
205. Dreyer G (d), Noroes J, Rocha A, Addiss D. Detection of living adult *Wuchereria bancrofti* in a patient with tropical pulmonary eosinophilia. Brazilian Journal of Medical and Biological Research 1996; 29:1005-1008.
206. Dreyer G, Noroes J, Addiss D. The silent burden of sexual disability associated with lymphatic filariasis. Acta Tropica 1997; 63:57-60.
207. Dreyer G, Noroes J, Addiss D. Proposed panel of diagnostic criteria, including the use of ultrasound, to refine the concept of "endemic normals" in lymphatic filariasis. Tropical Medicine and International Health 1999; 4:575-579.
208. Dreyer G (a), Medeiros Z, Nett MJ, Leal NC, de Castro LG, Piessens WF. Acute attacks in the extremities of persons living in an area endemic for bancroftian filariasis: differentiation of two syndromes. Transactions of the Royal Society of Tropical Medicine and Hygiene 1999; 93:413-417.
209. Dreyer G (b), Noroes J, Addiss D, Santos A, Medeiros Z, Figueredo-Silva J. Bancroftian filariasis in a paediatric population: an ultrasonographic study. Transactions of the Royal Society for Tropical Medicine and Hygiene 1999; 93:633-636.
210. Daffier B, Lepage T. Une cause rare d'hématurie: la filariose. Journal of Urology and Nephrology (Paris) 1978; 84:231-235.
211. Duke BOL. The effect of levamisole on *Onchocerca volvulus*. Transactions of the Royal Society of Tropical Medicine and Hygiene 1974; 68:1-4.
212. Dunn FL. Human behavioural factors in the epidemiology and control of *Wuchereria* and *Brugia* infections. Bulletin of the Public Health Society of Malaysia 1976;
213. Dunn FL. Behavioural aspects of the control of parasitic diseases. Bulletin of the World Health Organisation 1979; 57; 499-506.
214. Dunyo SK, Nkrumah FK, Collins K, Simonsen P. Exfoliative skin manifestations in acute lymphatic filariasis. Transactions of the Royal Society of Tropical Medicine and Hygiene 1998; 92:539-540.
215. Dunyo SK, Nkrumah FK, Simonsen P. A study on the procative day test effect of ivermectin and albendazole on nocturnal periodic *Wuchereria bancrofti* microfilaraemia. Transactions of the Royal Society of Tropical Medicine and Hygiene 1999; 93:608-610.
216. Dutta A, Sukul N. Filariacidal properties of a wild herb *Andrographis paniculata*. Journal of Helminthology 1982; 56:81-84.
217. Dutta A, Sukul N. Antifilarial effect of *Zingiber officinale* on *Dirofilaria immitis*. Journal of Helminthology 1987; 61:268-270.
218. Dutta SN, Diesfeld HJ. Laboratory parameters in early and late cases of Bancroftian filariasis in an endemic area. Journal of Communicable Diseases 1987; 4:15-416.
219. Dutta SN, Diesfeld HJ. Evidence of sex variations in microfilaraemia and fluorescent antibody titre level at puberty in a bancroftian filariasis area. Journal of Communicable Diseases 1994; 26:43-51.
220. Dzodzomenyo M, Dunyo SK, Ahorlu CK, Coker WZ, Appawu MA, Pedersen EM, Simonsen PE. Bancroftian filariasis in an irrigation project community in southern Ghana. Tropical Medicine and International Health 1999; 4:13-18.
221. Eberhard ML, Roberts IM, Lammie PJ, Lowrie RC. Comparative densities of *Wuchereria bancrofti* microfilaria in paired samples of capillary and venous blood. Tropical Medicine & Parasitology 1988; 39:295-298.
222. Eberhard ML, Lammie PJ, Dickinson CM, Roberts M. Evidence of non-susceptibility to diethylcarbamazine. Journal of Infectious Diseases 1991; 163:1157-1160.
223. Eberhard (a) ML, Dickinson CM, Hightower AW, Lammie MJ. Bancroftian filariasis: long term effects of treatment with diethylcarbamazine in a Haitian population. American Journal of Tropical Medicine and Hygiene 1991; 45:728-733.
224. Eberhard ML (b), Lammie MJ. Laboratory diagnosis of filariasis. Clinics in Laboratory Medicine 1991; 11:977-1010.
225. Eberhard ML, Hightower AW, McNeeley DF, Lammie MJ. Long-term suppression of microfilaraemia following ivermectin therapy. Transactions of the Royal Society of Tropical Medicine and Hygiene 1992; 86:287-288.
226. Eberhard ML, Hitch WL, McNeeley DF, Lammie MJ. Transplacental transmission of *Wuchereria bancrofti* in Haitian women. Journal of Parasitology 1993; 79:62-66.
227. Eberhard MJ, Walker EM, Addiss DG, Lammie MJ. A survey of knowledge, attitudes and perceptions (KAPs) of lymphatic filariasis, elephantiasis, and hydrocele among residents in an endemic area of Haiti. American Journal of Tropical Medicine and Hygiene 1996; 54:299-303.
228. Eberhard ML, Hightower AW, Addiss DG, Lammie MJ. Clearance of *Wuchereria bancrofti* antigen after treatment with diethylcarbamazine or ivermectin. American Journal of Tropical Medicine and Hygiene 1997; 57:483-486.
229. Edeson JFB. Clinical diagnosis of filariasis (letter). Transactions of the Royal Society of Tropical Medicine and Hygiene 1955; 49:488-489.
230. Edeson JF, Wharton RH. The experimental transmission of *W. malayi* to various animals in Malaya. Transactions of the Royal Society of Tropical Medicine & Hygiene 1958; 52:25-38.
231. Egwang TG, Nguiri C, Kombila M, Duong TH, Richard-Lenoble D. Elevated antifilarial IgG4 antibody levels in microfilaraemic and microfiladermic Gabonese adults and children. American Journal of Tropical Medicine & Hygiene 1993; 49:135-142.
232. El Bassiouny AEI, El Gammal NE, Mahmoud AM. Isolation and concentration of microfilariae from peripheral blood of *Wuchereria bancrofti*-infected patients by density gradient centrifugation. Journal of the Egyptian Society of Parasitology 1993; 23:255-262.
233. Elson LH, Calvopina MH, Paredes W, Araujo E, Bradley JE, Guderian RH, Nutman TB. Immunity to

- onchocerciasis: putative immune persons produce a Th1-like response to *Onchocerca volvulus*. The Journal of Infectious Diseases 1995; 171:652-658.
234. El Serougi Ahmed OM.. Value of quantitative buffy coat capillary tube test in the microscopic diagnosis of bancroftian filariasis. Journal of the Egyptian Society of Parasitology 1999; 29:223-228.
 235. El-Tamami MY. Effect of changed habits of life during the fasting month of ramazan and of treatment on the periodicity of Bancroftian filaria in Rosetta, Egypt. Journal of the Egyptian Medical Association 1960; 43:125-163.
 236. Englehorn TD, Wellman WE. Filariasis in soldiers on an island in the South Pacific. American Journal of the Medical Sciences 1945; 209:141-152.
 237. Estambale BBA (a), Simonsen PE, Knight R, Bwayo JJ. Bancroftian filariasis in Kwale district of Kenya 1. Clinical & parasitological survey in an endemic community. Annals of Tropical Medicine & Parasitology 1994; 88:145-151.
 238. Estambale BBA (b), Simonsen PE, Vennervald BJ, Knight R, Bwayo JJ. Bancroftian filariasis in Kwale district of Kenya 2. Humoral immune responses to filarial antigens in selected individuals from an endemic community. Annals of Tropical Medicine & Parasitology 1994; 88:153-161.
 239. Estambale BBA, Simonsen PE, Vennervald BJ, Knight R, Bwayo JJ. Bancroftian filariasis in Kwale district of Kenya 3. Quantitation of the IgE response in selected individuals from an endemic community. Annals of Tropical Medicine & Parasitology 1995; 89:287-295.
 240. Evans DB, Gelband H, Vlassoff C. Social and economic factors and the control of lymphatic filariasis: a review. Acta Tropica 1993; 53:1-26.
 241. Ewert A, Reitmeyer JC, Folse D. Chronic infection of cats with *Brugia malayi* and streptococcus. Southeast Asian Journal of Tropical Medicine & Public Health 1980; 11:32-39.
 242. Eyles DE, Hunter GW, Warren VG. The periodicity of microfilariae in two patients with filariasis acquired in the South Pacific. American Journal of Tropical Medicine 1947; 27:203-209.
 243. Fan PC. Diethylcarbamazine treatment of Bancroftian and malayan filariasis with emphasis on side effects. Annals of Tropical Medicine and Parasitology 1992; 86:399-402.
 244. Farid HA, Morsy ZS, Gad AM, Ramzy RM, Faris R, Weil GJ. Filariasis transmission potential of mosquitoes to humans of different age groups. Journal of the Egyptian society of parasitology 1997; 27:355-364.
 245. Faris R, Hussein O, El Setouhy M, Weil GJ. Bancroftian filariasis in Egypt: visualisation of adult worms and subclinical lymphatic pathology by scrotal ultrasound. American Journal of Tropical Medicine & Hygiene 1997; 57 (supplement):214 (abstract).
 246. Fiennes R.N.T.W. The possible influence of filaria on the development of viral infections in new world primates. Annals of the New York Academy of Sciences 1969; 162:600-603.
 247. Figueredo- Silva, Jungman P, Noroes J, Piessens WF, Coutinho A, Brito C, Rocha A, Dreyer G. Histological evidence for adulticidal effect of low doses of diethylcarbamazine in Bancroftian filariasis. Transactions of the Royal Society of Tropical Medicine and Hygiene 1996; 90:192-194.
 248. Fischer P, Liu X, Lizotte-Waniewski M, Kamal Ibrahim H, Ramzy Reda MR, Williams SA. Development of a quantitative, competitive polymerase chain reaction-enzyme linked immunosorbent assay for the detection of *Wuchereria bancrofti* DNA. Parasitology Research 1999; 85:176-183.
 249. Fonticiella M, L'opez-Negrete L, Prieto A, Garcia-Hernandez JB, Orense M, Fernandez-Diego J, Gomez JL. Congenital intra cranial filariasis: a case report. Paediatric radiology 1995; 25:171-172.
 250. Flynn PD. Filariasis suspects: a review of cases admitted. Naval Medical Bulletin 1944; 42:1075-1079.
 251. Fogel RH, Huntingdon JR. Genital manifestations of early filariasis. United States Navy Medical Bulletin 1945; 43:263-270.
 252. Forsyth KP, Spark R, Kazura J. A monoclonal antibody-based assay for detection of circulating antigen in Bancroftian filariasis. Journal of Immunology 1985; 134:1172-1177.
 253. Forsyth K. New approaches to the control of lymphatic filariasis using diethylcarbamazine. Papua New Guinea Medical Journal 1987; 30:189-191.
 254. Foster DG. Filariasis - a rare cause of pericarditis. Journal of tropical Medicine and hygiene 1956; 59:213-214.
 255. Foster R, Pringle G, King DF, Paris J. The therapeutic activity of some nitrofurans in experimental filariasis and trypanosomiasis. Annals of Tropical Medicine and Parasitology 1969; 63:95-107.
 256. Francis H, Awadzi K, Ottesen EA. The Mazotti reaction following treatment of onchocerciasis with diethylcarbamazine: clinical severity as a function of infection intensity. American Journal of Tropical Medicine and Hygiene 1985; 34:529-536.
 257. Franke ED, Räuber W, Wadded I. In-vitro cultivation of third stage larvae of *Wuchereria bancrofti* to the fourth stage. American Journal of Tropical Medicine & Hygiene 1987; 37:370-375.
 258. Franks MB, Kenneth BM, STOL NR. Reactions of natives of Okinawa and of American personnel to skin tests with antigen prepared from microfilariae of *Dirofilaria immitis*. American Journal of Tropical Medicine 1947; 27:617-632.
 259. Freedman DO, de Almeida Filo MJ, Besh S, de Silva MC, Braga C, Maciel A. Lymphoscintigraphic analysis of lymphatic abnormalities in symptomatic and asymptomatic human filariasis. Journal of Infectious Diseases 1994; 170:927-933.
 260. Freedman DO, de Almeida Filo MJ, Besh S, de Silva MC, Braga C, Maciel A, Furtado AF. Abnormal lymphatic function in presymptomatic bancroftian filariasis. Journal of Infectious Diseases 1995; 171:997-1001.
 261. Freedman DO. The VLA-4/VCAM-1 pathway is involved in the trans-endothelial migration of lymphocytes in Bancroftian filariasis. Journal of Immunology 1996; 156:2901-2908.
 262. Freedman DO. Immune dynamics in the pathogenesis of human lymphatic filariasis. Parasitology Today 1998; 14:229-234.
 263. Furtado AF, Abath FG, Regis L, Gomes YM, Lucena WA, Furtado PB, Dhalia R, Miranda JC, Nicolas L. Improvement and application of a polymerase chain reaction system for the detection of *Wuchereria bancrofti* in *Culex quinquefasciatus* and human blood samples. Memoirs of the Instituto Oswaldo Cruz 1997; 92:85-86.
 264. Gad AM, Feinsod FM, Soliman BA, Nelson GO, Gibbs PH, Shoukry A. Exposure variables in Bancroftian filariasis in the Nile delta. Journal of the Egyptian Society of Parasitology 1994; 24:439-455.
 265. Gajanana A, Bheema Rao US, Manonmani LM.

- Preliminary study on circulating immune complexes in Bancroftian filariasis. Indian Journal of Medical Research 1982; 76:146-150.
266. Galliard H. Outbreak of filariasis (*Wuchereria malayi*) among French & North African service men in North Vietnam. Bulletin of the World Health Organisation 1957; 16:601-608.
 267. Gam AA, Neva FA, Krotoski WA. Comparative sensitivity and specificity of ELISA and IHA for serodiagnosis of strongyloidiasis with larval antigens. American Journal of Tropical Medicine and Hygiene 1987; 37:157-161.
 268. Gao CL, Cao WC, Chen XX. Changes in anti-filarial antibody after control of filariasis in Shandong province. Chinese Medical Journal 1994; 107:360-363.
 269. Garg RK, Jain S, Kar AM. Pott's paraplegia-like presentation: a neurological complication of lymphatic filariasis. Journal of the Physician's Association of India 1996; 44:741-742.
 270. Gayral P, Loiseau PM, Bories C, Brienne MJ, Leclercq M, Jacques J. *In-vitro* and *in-vivo* antifilarial activity of ethylenesulfonamides, epoxyethanesulfonamides and carboxamide analogues. Arzneimittelforschung 1995; 45:1122-1127.
 271. Gelbrand H. Diethylcarbamazine salt in the control of lymphatic filariasis. American Journal of Tropical Medicine and Hygiene 1994; 50:655-662.
 272. Gilbert HM, Hartman BJ. Short report: a case of fibrosing mediastinitis caused by *Wuchereria bancrofti*. American Journal of Tropical Medicine and Hygiene 1996; 54:596-599.
 273. Glauser F. Filariasis in returning marines. Naval Medical Bulletin 1945; 44:21-36.
 274. Goa KL, McTavish D, Clissold SP. Ivermectin. A review of its antifilarial activity, pharmacokinetic properties and clinical efficiency in Onchocerciasis. Drugs 1991; 42:640-658.
 275. Goel TC, Misra G. Omental transposition in the treatment of the treatment of tropical elephantiasis of the lower limb. British Journal of Surgery 1968; 55:440-446.
 276. Goldsmid JM, Mahomed K, Makanji H, Muir M. Microhaematocrit centrifuge technique for the laboratory diagnosis of filarial infections. South African Medical Journal 1972; 46:171-174.
 277. Goldsmid JM, Rogers S. The cytocentrifuge for the isolation of parasites from the blood. Central African Journal of Medicine 1975; 21:160-161.
 278. Goldsmid JM (a), Rogers S. Studies on the diagnosis and treatment of human filariasis in Rhodesia. South African Medical Journal 1976; 50:1129-1132.
 279. Goldsmid JM (b), Rogers S, Cookson J. The use of acridine orange O in the diagnosis of human filariasis. Central African Journal of Medicine 1976; 22:4-6.
 280. Goldsmith HS, De Los Santos R, Beattie EJ. Relief of chronic lymphedema by omental transposition. Annals of Surgery 1967; 166:847-859.
 281. Goldsmith HS. Long term evaluation of omental transposition for chronic lymphedema. Annals of Surgery 1974; 180:847-859.
 282. Goodman AA, Weinberger EM, Lippincott SW, Marble A, Wright WH. Studies of filariasis in soldiers evacuated from the South Pacific. Annals of Internal Medicine 1945; 23:823-836.
 283. Gopinath R, Hanna LE, Kumaraswami V, Pillai SVP, Kavitha V, Vijayasekaran V, Rajaskharan A, Nutman TB. Long-term persistence of cellular hyporesponsiveness to filarial antigens after clearance of microfilaraemia. American Journal of Tropical Medicine and Hygiene 1999; 60:848-853.
 284. Gourlay WA, Chiu A, Montessori VC, Dunn IJ. Urinary filariasis presenting as bladder pseudotumours. Journal of Urology 1999; 161:603-604.
 285. Grace AW. The parallel incidence of *Filaria bancrofti* and the α -haemolytic streptococcus in certain tropical countries. American Journal of Tropical Medicine 1932; 12:493-508.
 286. Grace AW. Filarial lymphangitis, considered as a mild erysipelas resulting from hypersensitivity to a β -haemolytic streptococcus of a particular type. Transactions of the Royal Society of Tropical Medicine and Hygiene 1934; 28:259.
 287. Grace AW. Tropical lymphangitis and abscesses. Journal of the American Medical Association 1943; 13:462-463.
 288. Greene BM, Otto GF, Greenough WB III.. Circulating non-human microfilariae in a patient with lupus erythematosus.
 289. American Journal of Tropical Medicine and Hygiene 1978; 27:905-909.
 290. Greene BM, Fanning MM, Ellner JJ. Non-specific suppression of antigen-induced lymphocyte blastogenesis in *Onchocerca volvulus* infection in man. Clinical and Experimental Immunology 1983; 52:259-263.
 291. Greene BM, Gbakima AA, Albiez EJ, Taylor HR. Humoral and cellular immune responses to *Oncocerca volvulus*. Review of Infectious Diseases 1985; 7:789-795.
 292. Greenberg JH. Public health problems relating to the Vietnam returnee. Journal of the American Medical Association 1969; 207:697-702.
 293. Grieve RB, Wisniewski N, Frank GR, Tripp CA. Vaccine research and development for the prevention of filarial nematode infections. Pharmaceutical Biotechnology 1995; 6:737-768.
 294. Grove DI. Immunity in filariasis: a review. Papua New Guinea Medical Journal 1978; 21:32-42.
 295. Grove DI. Immuno-suppression in Bancroftian filariasis. Transactions of the Royal Society of Tropical Medicine & Hygiene 1979; 73:23-26.
 296. Grove DI. Serum microfilarial antibody titres before and after treatment of Bancroftian microfilaraemia with diethylcarbamazine. Southeast Asian Journal of Tropical Medicine and Public Health 1981; 12:487-491.
 297. Grove DI. *Wuchereria bancrofti*, *Brugia* species and filariasis. In: History of Human Helminthology chapter 23. Published by C.A.B. International 1990.
 298. Gupta A, Agarwal A, Dogra Mangat R. Retinal involvement in bancrofti filariasis. Acta Ophthalmology 1992; 70:832-835.
 299. Gyapong JO, Magnussen P, Binka FN. Parasitological and clinical aspects of Bancroftian filariasis in Kassena-Nankana district, upper east region, Ghana. Transactions of the Royal Society of Tropical Medicine and Hygiene 1994; 88:555-557.
 300. Gyapong JO, Dollimore N, Binka FN, Ross DA. Lay reporting of elephantiasis of the leg in northern Ghana. Transactions of the Royal Society of Tropical medicine and Hygiene 1995; 89:616-618.
 301. Gyapong JO (a), Gyapong M, Evans DB, Aitkins MK, Adjei S. The economic burden of lymphatic filariasis in northern Ghana. Annals of Tropical Medicine & Parasitology 1996; 90:39-48.

302. Gyapong M, Gyapong JO (b), Adjei S, Vlassoff C, Weiss M. Filariasis in Northern Ghana: some cultural beliefs & practices & their implication for disease control. *Social Science and Medicine* 1996; 43:235-242.
303. Gyapong JO (c), Gyapong M, Adjei S. The epidemiology of adenolymphangitis due to lymphatic filariasis in Northern Ghana. *American Journal of Tropical Medicine and Hygiene* 1996; 54:591-595
304. Gyapong JO (a), Webber RH, Morris J, Bennett S. Prevalence of hydrocele as a rapid diagnostic index for lymphatic filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1998; 92:40-43.
305. Gyapong JO (b), Omane-Badu, Webber RH. Evaluation of the filter paper blood collection method for detection Og4C3 circulating antigen in Bancroftian filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1998; 92:407-410.
306. Gyapong JO (c), Webber RH, Bennett S. The potential role of peripheral health workers and community key informants in the rapid assessment of community burden of disease: the example of lymphatic filariasis. *Tropical Medicine and International Health* 1998; 3:522-528.
307. Gyorkos TW, Sukul NC, Dasdal A. Filariasis and ABO blood groups: a critical appraisal. *Transactions of the Royal Society of Tropical Medicine and Parasitology* 1983; 77:564-565.
308. Haarbrink M, Terhell A, Abadi K, van Beers S, Asri M, Yazdanbakhsh M. IgG4 antibody assay in the detection of filariasis. *Lancet* 1995; 346:853-853.
309. Haarbrink M (b), Terhell A, Abadi GK, Asri M, de Medeiros F, Yazdanbakhsh M. Anti-filarial IgG4 in men and women living in a *Brugia malayi*-endemic area. *Tropical Medicine and International Health* 1999; 4:93-97.
310. Haarbrink M (b), Terhell A, Abadi GK, Mitsui Y, Yazdanbakhsh M. Adverse reactions following intake in "endemic normals", microfilaraemics and elephantiasis patients. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1999; 93:91-96.
311. Haliza B, Mohd R. Comparison of knowledge on filariasis and epidemiological factors between infected and uninfected respondents in a Malay community. *Southeast Asian Journal of Tropical Medicine & Public Health* 1986; 17:457-463.
312. Hamilton RG, Hussain R, Ottesen EA, Franklin N. The quantitation of parasite-specific human IgG and IgE in sera: evaluation of solid-phase RIA and ELISA methodology. *Journal of Immunological Methods* 1981; 44:101-114.
313. Hamilton RG, Hussain R, Ottesen EA. Immuno-radiometric assay for the detection of filarial antigen in human serum. *Journal of Immunology* 1984; 133:2237-2242.
314. Hamilton RG. Application of immuno-assay methods in the sero-diagnosis of human filariasis. *Review of Infectious Diseases* 1985; 7:837-843.
315. Handa R, Gupta R, Wig N, Agarwal P, Wali JP. Microfilariae of *Wuchereria bancrofti* in ascitic fluid. *Indian Journal of Medical Science* 1996; 50:368-369.
316. Haque A, Capron A. Transplacental transfer of rodent microfilariae induces antigen-specific tolerance in rats. *Nature* 1982; 299:361-363.
317. Haque A, Capron A, Ouaisi A, Kouemini L, Lejeune JP, Bonnel B, Pirce R. Immune unresponsiveness and its possible role to filarial disease. *Contributions to Microbiology and Immunology* 1983; 7:9-21.
318. Haque A, Cuna W, Pestel J, Capron A, Bonnel B. Tolerance in rats by transplacental transfer of *Dipetalonema viteae* microfilariae: recognition of putative toleragen(s) by antibodies that inhibit antigen-specific lymphocyte proliferation. *European Journal of Immunology* 1988; 18:1167-1172.
319. Harinath BC. Detection and diagnostic ability of *in-vitro* and *in-vivo* released antigens in Bancroftian filariasis. *Journal of Communicable Diseases* 1986; 18:261-266.
320. Haviland JW. Recent experiences with filariasis. *Northwest Medicine* 1944; 371-373.
321. Hawking F (a). Some recent work on filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1950; 44:153-193
322. Hawking F (b), Sewell P, Thurston JP. The mode of action of hetrazan on filarial worms. *British Journal of Pharmacology* 1950; 5:217-238.
323. Hawking F (a). Microfilaria infestation as an instance of periodic phenomenon seen host parasite relationships. *Annals of the New York Academy of Science* 1962; 98:940-953.
324. Hawking F (b). The role of the spleen in controlling the number of microfilariae (*Dirofilaria immitis*, *D. Repens*, *Litomosoides carini* and *Dipetalonema vitei*) in the blood. *Annals of Tropical Medicine and Parasitology* 1962; 56:168-172.
325. Hawking F, Denham D. The distribution of human filariasis throughout the world: Part 1, the Pacific region including New Guinea. *Tropical Diseases Bulletin* 1976; 73:347-371.
326. Hayashi Y, Noda K, Shirasaka A, Nogami S, Nakamura M. Vaccination of BALB/c mice against *Brugia malayi* and *Brugia pahangi* with larvae attenuated by gamma irradiation. *Japanese Journal of Experimental Medicine* 1984; 52:177-189.
327. Hewitt RI, White E, Wallace WS, Stewart HW, Kushner S, Subbarow Y. Experimental Chemotherapy of filariasis 2. Effect of piperrazine derivatives against naturally acquired filarial infections in cotton rats and dogs. *Journal of Laboratory and Clinical Medicine* 1947; 32:1304-1313.
328. Hightower AW, Lammie MJ, Eberhard ML. Maternal filarial infection - a persistent risk factor for microfilaraemia in offspring? *Parasitology Today* 1993; 9:418-421.
329. Hines SA, Williams JL, Doyle TJ, Crandall RB, Crandall CA, Nayar JK. Lymphangiography in ferrets infected with *Brugia malayi*. *Lymphology* 1985; 18:173-174.
330. Hirst WR, McCann WJ. Tropical eosinophilia. Report of a case. *United States Navy Medical Bulletin* 1945; 44:1277-1281.
331. Hitch WL, Lammie MJ, Walker EM, Hightower AW, Eberhard ML. Antifilarial cellular responses detected in a Haitian pediatric population by use of a microblastogenesis assay. *Journal of Infectious Diseases* 1991; 164:811-813
332. Hitch WL, Eberhard ML, Lammie MJ. Investigation of the influence of maternal infection with *Wuchereria bancrofti* on the humoral and cellular responses of neonates to filarial antigens. *Annals of Tropical Medicine & Parasitology* 1997; 91:461-469.
333. Hodge IG, Denhoff E, Van der Veer JB. Early filariasis (Bancrofti) in American soldiers. *American Journal of the Medical Sciences* 1945; 210:207-223.
334. Hoeppli R. Parasitic diseases in Africa & the western hemisphere. *Acta Tropical* 1969; 10 supplement:240-246.
335. Hoeppli R. Early navigation and the spread of parasitic diseases. *Episteme* 1971; 5:201-217.

336. Hoerauf A, Nissen-Pahle K, Schmetz C, Henkle-Duhrsen K, Blaxter ML, Buttner DW, Gallin MY, Al-Qaoud KM, Lucious R, Fleischer B. Tetracycline therapy targets intracellular bacteria in the filarial nematode *Litomosoides sigmodontis* and results in filarial infertility. *Journal of Clinical Investigation* 1999; 103:11-18.
337. Hoerauf A, Volkmann L, Hamelmann C, Adjei O, Autenrieth I, Fleischer B, Buttner DW. Endosymbiotic bacteria in worms as targets for a novel chemotherapy for filariasis. *Lancet* 2000; 355:211-213.
338. Howells RE, Medis AM, Bray PG. The mode of action of suramin on the filarial worm *Brugia pahangi*. *Parasitology* 1983; 87:29-48.
339. Hughes CC, Hunter JM. Disease and development in Africa. *Social Science & Medicine* 1970; 3:443-493.
340. Hunter JM, Rey L, Scott D. Man-made lakes and man-made diseases; towards a policy resolution. *Social Science & Medicine* 1982; 16:1127-1145.
341. Hunter JM, Rey L, Chu KY, Adekolu-John EO, Mot KE. Parasitic diseases in water resource development: the need for intersectoral negotiation. WHO TDR/SCH, Geneva 1991.
342. Hunter JM. Elephantiasis: a disease of development in north east Ghana. *Social Science Medicine* 1992; 35:627-649.
343. Huntington RW, Fogel RH, Eichold A, Dickson JG. Filariasis among American Troops in a South Pacific island group. *Yale Journal of Biology and Medicine* 1944; 16:529-536.
344. Huntington RW, Eichold S, Scott OK. *American Journal of Tropical Medicine and Hygiene* 1950; 30:873-880.
345. Hussain R, Grogil M, Ottesen EA. IgG antibody subclasses in human filariasis. *Journal of Immunology* 1987; 139:2794-2798.
346. Hussain R, Poindexter RW, Ottesen EA. Control of allergic reactivity in human filariasis: predominant localisation of blocking antibody to IgG4 subclass. *Journal of Immunology* 1992; 148:2731-2737.
347. Ismail MM, Premaratne UN, Abeyewickreme W. Treatment of Bancroftian filariasis with ivermectin in Sri Lanka: evaluation of efficacy and adverse reactions. *Tropical Biomedicine* 1991; 8:71-75.
348. Ismail MM, Weil GJ, Jayasinghe KS, Premaratne UN, Abeyewickreme W, Rajaratnam HN, Sheriff MH, Perera CS, Dissanaik AS. Prolonged clearance of microfilaraemia in patients with Bancroftian filariasis after multiple high doses of ivermectin of diethylcarbamazine. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1996; 90:684-688.
349. Ismail MM, Jayakody RL, Weil GJ, Nirmalan N, Jayasinghe KSA, Abeyewickreme W, Rezvi Sheriff MH, Rajaratnam HN, Amarasekera N, de Silva DCL, Michalski ML, Dissanaik AS. Efficacy of single dose combinations of albendazole, ivermectin and diethylcarbamazine for the treatment of Bancroftian filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1998; 92:94-97.
350. Itoh M, Gunawardena NK, Xu-Guang Q, Weerasooriya MV, Kimura E. The use of whole blood absorbed on filter paper to detect *Wuchereria bancrofti* circulating antigen. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1998; 92:513-515.
351. Iyengar MOT. Filariasis in Trivandrum. *Indian Journal of Medical Research Memoir* 1938, No. 30.
352. Jamal S (a) Summary of symposium on elephantiasis and management of swollen limb. *Lymphology* 1989; 22:103-114.
353. Jamal S (b), Casley-Smith JR, Casley-Smith Judith R. The effects of 5,6 benzo-(a)pyrone (coumarin) and DEC on filaritic lymphoedema and elephantiasis. Preliminary results. *Tropical Medicine and Parasitology* 1989; 83:287-290.
354. James DM, Gillies HM In: *Human Antiparasite Drugs: Pharmacology and Usage*. John Wiley and Sons, Brisbane 1985.
355. Jing-yuan L, Xin-ji L, Zi C, Zhao-ping T, Guo-bin Z, Yue-nan C, Ying-zhen Z, Shao-peng W, Xian-hong H, Fa-zhu Y. Filariasis and its control in Fujian, China. *Southeast Asian Journal of Tropical Medicine and Public Health* 1991; 22:147-154.
356. Jing-yuan L, Zi C, Xiao-hong H, Zhao-ping T. Mass treatment of filariasis using DEC-medicated salt. *Journal of Tropical Medicine and Hygiene* 1992; 95:132-135.
357. Johnson PAG. Filariasis: clinical findings in 189 cases. *United States Navy Medical Bulletin* 1944; 43:950-954.
358. Jones TC, Mott K, Cavalcante Pedrosa L. A technique for isolating and concentrating microfilariae from peripheral blood by gradient centrifugation. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1975; 69:243-246.
359. Jordan P. Notes on elephantiasis and hydrocele due to *Wuchereria bancrofti*. *Journal of Tropical Medicine and Hygiene* 1955; 58:113-118.
360. Jordan P. The treatment of Bancroftian filariasis and observations on hyaluronidase. *Tropical Medicine and Hygiene* 1959; 62:287-288.
361. Joseph CA, Dixon PA. Possible prostaglandin mediated effect of diethylcarbamazine on rat uterine contractility. *Journal of Pharmacy and Pharmacology* 1984; 36:281-282.
362. Kagan I.G. A review of the immunologic methods for the diagnosis of filariasis. *Journal of Parasitology* 1963; 49:773-798.
363. Kar SK, Mania J, Kar PK. Humoral immune response during filarial fever in Bancroftian filariasis. *Transactions of the Royal Society of Tropical Medicine & Hygiene* 1993; 87:230-233.
364. Kaur V. Tropical diseases and women. *Clinics in Dermatology* 1997; 15:171-178.
365. Kazura JW, Davis RS. Soluble *Brugia malayi* microfilarial antigens protect mice against challenge by an antibody-dependant mechanism. *Journal of Immunology* 1982; 128:1792-1796.
366. Kazura JW, Spark R, Forsyth K, Brown G, Heywood P, Alpers M. Parasitologic and clinical features of Bancroftian filariasis in the Sepik Province of Papua New Guinea. *American Journal of Tropical Medicine and Hygiene* 1984; 33:1119-1123.
367. Kazura JW (a), Cicirello H, McCall JW. Induction of protection against *Brugia malayi* infection in jirds by microfilaria antigens. *Journal of Immunology* 1986; 136:1422-1426.
368. Kazura JW (b), Cicirello H, Forsyth KP. Differential recognition of a protective filarial antigen by antibodies from humans with Bancroftian filariasis. *Journal of Clinical Investigation* 1986; 77:1985-1992.
369. Kazura JW. Ivermectin and lymphatic filariasis. *Microbiological Pathogenesis* 1993; 14:337-342.
370. Kazura JW (a), Nutman TB, Greene BM. Filariasis. In: *Immunology and Molecular Biology of Parasitic Infections*, 3rd edition, chapter 18 Edited by Warren KS, Blackwell Scientific Publications, Boston 1993.
371. Kazura J (b), Greenberg J, Perry R, Weil G, Day K,

- Alpers M. Comparison of single-dose diethylcarbamazine and ivermectin for the treatment of Bancroftian filariasis in Papua New Guinea. *American Journal of Tropical Medicine and Hygiene* 1993; 49:804-811.
372. Kazura JW, Bockarie M, Alexander N, Perry R, Bockarie F, Dagoro H, Dimber Z, Hyun P, Alpers MP. Transmission intensity and its relationship to infection and disease due to *Wuchereria bancrofti* in Papua New Guinea. *Journal of Infectious Diseases* 1997; 176:242-246.
373. Kessel JF. Disabling effects and control of filariasis. *American Journal of Tropical Medicine & Hygiene* 1957; 6:402-414.
374. Khamboonruang C, Thitasut P, Pan-In S, Morakote N, Choochote W, Somboon P, Keha P. Filariasis in Tak province, Northwest Thailand: the presence of sub-periodic variant *Wuchereria bancrofti*. *Southeast Asian Journal of Tropical Medicine & Public Health* 1987; 18:218-222.
375. Khan Am, Dutta P, Khan SA, Baruah NK, Sarma CK, Mahanta J. Long term effect of diethylcarbamazine citrate on microfilaraemia status in treated individuals. *Indian Journal of Medical Research* 1998; 108:134-138.
376. Kharat I, Satyanarayana U, Ghirmikar SN, Harinath BC. *In vitro* cultivation of *Wuchereria bancrofti* microfilariae. *Indian Journal of Experimental Biology* 1980; 18:1245-1247.
377. Kimata H, Yoshida A, Ishioka C, Mikawa H. Disodium cromoglycate (DSCG) selectively inhibits IgE production and enhances IgG4 production by human B cells *in vitro*. *Clinical and Experimental Immunology* 1991; 84:395-399.
378. Kimura E, Penania L, Spears GFS. The efficacy of annual doses of diethylcarbamazine citrate against diurnally sub-periodic Bancroftian filariasis in Samoa. *Bulletin of the World Health Organisation* 1985; 63:1097-1106.
379. Kimura E, Mataika JU. Control of lymphatic filariasis by annual single dose diethylcarbamazine treatments. *Parasitology Today* 1996; 12:240-244.
380. King BG. Early filariasis diagnosis and clinical findings: report of 268 cases in American troops. *American Journal of Tropical Medicine* 1944; 24:285-296.
381. King CL, Kumaraswami V, Poindexter RW, Kumari S, Jayaraman K, Alling DW, Ottesen EA, Nutman TB. Immunologic tolerance in lymphatic filariasis: diminished parasite-specific T and B lymphocyte precursor frequency in the microfilaraemic state. *Journal of Clinical Investigation* 1992; 89:1403-1410.
382. Kinnamon KE, Engle RR, Poon BT, McCall JW, Dzimianski MT. A new class of anti-filarial compounds - a preliminary look. *Military Medicine* 1994; 159:368-372.
383. Kip W, Burham G, Kamugisha J. Improvements in seizures after ivermectin. *Lancet* 1992; 340:789-790.
384. Kitchen LW, Cotter SM. Effect of diethylcarbamazine on serum antibody to feline oncornavirus-associated cell membrane antigen in feline leukaemia virus cats. *Journal of Clinical and Laboratory Immunology* 1988; 25:101-103.
385. Kitchen LW, Mather FJ, Chapple FE, Biello JA. Effect of administration of diethylcarbamazine on murine leukaemia virus (Cas-Br-M) infected mice. *Journal of Clinical and Laboratory Immunology* 1990; 33:97-105.
386. Kitchen LW, Ross JA, Hernandez JE, Zarraga AL, Mather FJ. Effect of administration of diethylcarbamazine on experimental bacterial and fungal infections in mice. *International Journal of Antimicrobial Agents* 1992; 1:259-267.
387. Kitchen LW, Ross JA, Hernandez JE, Zarraga AL, Mather FJ. Effect of administration of diethylcarbamazine with and without fluconazole on experimental fungal infections in mice. *International Journal of Antimicrobial Agents* 1993; 2:237-246.
388. Kitchen LW (a), Ross JA, Turner GJ, Hernandez JE, Mather FJ. Diethylcarbamazine enhances blood microbicidal activity. *Advances in Therapy* 1995; 12:22-29.
389. Kitchen LW (b) Hernandez JE, Zarraga AL. Diethylcarbamazine treatment of chronic osteomyelitis Complications in Orthopedics 1995; 10:47-49.
390. Kitchen LW, Weston CM, Day SP. Diethylcarbamazine-related antimicrobial activity in Mycobacterium tuberculosis-infected blood. *Journal of Antimicrobial Therapy* 1998; 42:241-243.
391. Klei TR, Crowell WA, Thompson PE. Ultrastructural glomerular changes associated with filariasis. *American Journal of Tropical Medicine & Hygiene* 1974; 23:609-618.
392. Klei TR, Blanchard DP, Coleman SU. Development of *Brugia pahangi* infections and lymphatic lesions in male offspring of female jirds with homologous infections. *Transactions of the Royal Society of Tropical Medicine & Hygiene* 1986; 80:214-216.
393. Knight R, McAdam KPWJ, Matola YG, Kirkham V. Bancroftian filariasis and other parasitic infections in the middle fly region of western Papua New Guinea. *Annals of Tropical Medicine and Parasitology* 1979; 76:565-576.
394. Knott J. The periodicity of the microfilaria of *Wuchereria bancrofti*. Preliminary report of some injection experiments. *Transactions of the Royal Society for tropical Medicine and Hygiene* 1935; 29:59-64.
395. Kobayashi M, Niimura M, Kanazawa T, Husky MK, Malagueno S, Santana JV. Detection of microfilariae antigen in circulating immune complexes from sera of *Wuchereria bancrofti*-infected individuals. *American Journal of Tropical Medicine and Hygiene* 1997; 57:200-204.
396. Koya SM, Kumar S, Regu K, Sharma RC. Early development of acute filariasis in a migrant from non-endemic to hyper-endemic area: a case report. *Journal of Communicable Diseases* 1998; 30:117-120.
397. Krishnarao P, Suryakala G, Raina VK, Raina VK, Ghosh TK. Study of prenatal sensitization in relation to susceptibility or resistance to filarial infection or disease. *Journal of Communicable Diseases* 1989; 21:142-144.
398. Krishnarao P, Kaur R, Ghosh TK. Long term effect of diethylcarbamazine medicated common salt on bancroftian filariasis. *Journal of Communicable Diseases* 1991; 23:128-130.
399. Kumar H, Malhotra A, Santhanam S, Jain DC, Ghosh TK. Evaluation of specific humoral immune responses in human filariasis. *Folia Parasitologica* 1987; 34:249-254.
400. Kumar A. Human filariasis: infection rate as the uniform measurable criterion for filarial endemicity. *Journal of Communicable Diseases* 1996; 28:163-167.
401. Kumaraswami V, Ottesen EA, Vijayasekaran V. Ivermectin for the treatment of *Wuchereria bancrofti* filariasis: efficacy and adverse reactions. *Journal of the American Medical Association* 1988; 259:3150-3153.
402. Kumari S, Lillibridge CD, Bakeer M, Lowrie RC, Jayaraman K, Philip MT. *Brugia malayi*: the diagnostic potential of recombinant excretory/secretory antigens. *Experimental Parasitology* 1994; 79:489-505.
403. Kurniawan A, Yazdanbakhsh M, van Ree R, Aalberse R, Selkirk ME, Partono F, Maizels RM. Differential

- expression of IgE and IgG4 specific antibody responses in asymptomatic and chronic human filariasis. *Journal of Immunology* 1993; 150:3941-3950.
404. Kurniawan-Atmadja A, Atkinson R, Sartono E, Partono F, Yazdanbakhsh M, Maizels RM. Differential decline in filaria-specific IgG1, IgG4 and IgE antibodies in *Brugia malayi*-infected patients after diethylcarbamazine chemotherapy. *Journal of Infectious Diseases* 1995; 172:1567-1572.
 405. Kurniawan-Atmadja A, Sartono E, Partono F, Yazdanbakhsh M, Maizels RM. Antibody responses to filarial infected larvae are not dominated by the IgG4 isotype. *Parasite Immunology* 1998; 20:9-17.
 406. Kwa BH, Mak JW. Specific depression of cell-mediated immunity in Malayan filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1980; 74:522-527.
 407. Kwan-Lim GE, Forsyth KP, Maizels RM. Filarial specific IgG4 response correlates with active *Wuchereria bancrofti* infection. *Journal of Immunology* 1990; 145:4298-4305.
 408. Laing AB, Edeson JF, Wharton RH. Studies on filariasis in Malaya: the vertebrate hosts of *Brugia malayi* and *Brugia pahangi*. *Annals of Tropical Medicine & Parasitology* 1960; 54:92-99.
 409. Lal RB, Paranjape RS, Briles DE, Nutman TB, Ottesen EA. Circulating parasite antigens in lymphatic filariasis: use of monoclonal antibodies to phosphocholine for immuno diagnosis. *Journal of Immunology* 1987; 138:3454-3460.
 410. Lal RB, Ottesen EA. Enhanced diagnostic specificity in human filariasis by IgG4 antibody assessment. *Journal of Infectious Diseases* 1988; 158:1043-1038.
 411. Lal RB, Ottesen EA. Phosphocholine epitope on helminth and protozoal parasites and their presence in the circulation of infected human patients. *Transactions of the Royal Society of Tropical medicine & Hygiene* 1989; 83:652-655.
 412. Lal RB, Kumaraswami V, Steel C, Nutman TB. Phosphocholine-containing antigens of *Brugia malayi* non-specifically suppress lymphocyte function. *American Journal of Tropical Medicine & Hygiene* 1990; 42:56-64.
 413. Lal RB, Dhawan RR, Tarrand JJ, Ayoub EM, Ottesen EA. Lack of IgG4 antibody response to carbohydrate antigens in patients with lymphatic filariasis. *Immunology* 1991; 74:333-337.
 414. Lalitha P, Ravichandran M, Suba S, Kaliraj P, Narayanan Jayaraman K. Quantitative assessment of circulating antigens in human lymphatic filariasis: a field evaluation of monoclonal antibody-based ELISA using blood collected on filter paper strips. *Tropical Medicine and International Health* 1998; 41-45.
 415. Lammie PJ, Katz SP. Immuno regulation in experimental filariasis: *in-vitro* suppression of mitogen-induced blastogenesis by adherent cells from jirds chronically infected with *Brugia pahangi*. *Journal of Immunology* 1983; 130:1381-1385.
 416. Lammie PJ, Hitch WL, Walker Allen EM, Hightower W, Eberhard ML. Maternal infection as risk factor for infection in children. *Lancet* 1991; 337:1005-1006.
 417. Lammie PL, Hightower AW, Richards FO, Bryan RT, Spencer HC, McNeeley, McNeeley MB, Eberhard ML. Alterations in filarial antigen-specific immunologic reactivity following treatment with ivermectin and diethylcarbamazine. *American Journal of Tropical Medicine and Hygiene* 1992; 46:292-295.
 418. Lammie PJ, Addiss DG, Leonard G, Hightower AW, Eberhard ML. Heterogeneity in filarial-specific immune responsiveness among patients with lymphatic obstruction. *Journal of Infectious Diseases* 1993; 167:1178-1183.
 419. Lammie PJ, Hightower AW, Eberhard ML. The age-specific prevalence of antigenaemia in a *Wuchereria bancrofti*-exposed population. *American Journal of Tropical Medicine and Hygiene* 1994; 51:348-355.
 420. Lane C. Bancroftian filariasis. Biological mechanisms that underlie its periodicity and other of its clinical manifestations. *Transactions of the Royal Society of Tropical Medicine & Hygiene* 1948; 41:717-784.
 421. Langhammer J, Birk HW, Zahner H. Renal disease in lymphatic filariasis: evidence for tubular and glomerular disorders at various stages of infection. *Tropical Medicine and International Health* 1997; 2:875-884.
 422. Langy S, Plichart C, Luquaud P, Williams SA, Nicolas L. The immunodominant *Brugia malayi* paramyosin as a marker of current infection with *Wuchereria bancrofti* adult worms. *Infection and Immunity* 1998; 66:2854-2858.
 423. Lawrence BR. Elephantiasis in Greece and Rome and the Queen of Punt. *Transactions of the Royal Society for Tropical Medicine & Hygiene* 1967; 61:234-245.
 424. Lawrence BR. The curse of St. Thomas. *Medical History* 1970; 14:352-363.
 425. Lawrence BR. Sciapodes in tropical medicine. *Transactions of the Royal Society for Tropical Medicine & Hygiene* 1990; 84:174-175.
 426. Lawrence RA, Allen JE, Osbourne J, Maizels RM. Adult and microfilarial stages of the filarial parasite *Brugia malayi* stimulate contrasting cytokine and Ig responses in BALB/c mice. *Journal of Immunology* 1994; 153:1216-1224.
 427. Lazdins J, Kron M. New molecular targets for filariasis drug discovery. *Parasitology Today* 1999; 15:305-306.
 428. Leede WE, Josey. Al. The early diagnosis of filariasis and certain suggestions relative to cause of symptoms. *Annals of Internal Medicine* 1945; 23:816-823.
 429. Levin M, Pras M, Franklin EC. Immunologic studies of the major non-immunoglobulin protein of amyloid. 1: Identification and partial characterisation of a related serum component. *Journal of Experimental Medicine* 1973; 138:373-378.
 430. Lewis TR. On a haematozoon inhabiting human blood, its relation to chyluria and other diseases. Eighth Annual Report of the Sanitary Commissioner of the Government of India, Appendix B, pages 111-113, 1872.
 431. Lewis TR. Letter to P. Manson In: Manson-Bahr PH, Alcock A. *The Life & Works of Sir Patrick Manson*, published by Cassell & Company, London, 1927, page 273.
 432. Libman MD, Maclean JD, Gyorkos TW. Screening for schistosomiasis, filariasis and strongyloidiasis among expatriates returning from the tropics. *Clinical Infectious Diseases* 1993; 17:353-359.
 433. Lichtenstein A. Filaria-onderzoek te Bireuen. *Geneeskundig Tijdschrift voor Nederlandsch-Indie* 1927; 67:742-749.
 434. Linhares MI, Malagueno E, Carvalho LB, de Oliveira VF, Minamishima Y. No relationship between HTLV-1 infection and filariasis- serological study on patients with filariasis in Recife, Brazil. *Microbiology and Immunology* 1995; 39:917-919.
 435. Liu JY, Chen Z, Huang JH, Tu SP. Mass treatment of filariasis using DEC-medicated salt. *Journal of Tropical Medicine and Hygiene* 1992; 95:132-135.

436. Lai LX, Welder PF. Intra vascular filarial parasites inhibit platelet aggregation. Role of parasite-derived prostanoids. *Journal of Clinical Investigation* 1992; 89:1113-1120.
437. Lai LX, Serhan CN, Welder PF. Intra vascular filarial parasites elaborate cyclooxygenase-derived eicosanoids. *Journal of Experimental Medicine* 1990; 172:993-996.
438. Liu YK, Hsieh SC, Tai T. The role of streptococcal infections in filariasis. *Chinese Medical Journal* 1964; 83:17-20.
439. Lizotte MR, Taniawati S, Partono F, Williams SA. A polymerase chain reaction assay for the detection of *Brugia malayi* in blood. *American Journal of Tropical Medicine and Hygiene* 1994; 51:314-321.
440. Lobos E, Ondo A, Ottesen EA, Nutman TB. Biochemical and immunologic characterisation of a major IgE-inducing filarial antigen of *Brugia malayi* and implications for the pathogenesis of tropical pulmonary eosinophilia. *Journal of Immunology* 1992; 149:3029-3034.
441. Loiseau PM, Depreux P. *In-vitro* antifilarial evaluation of phenoxycyclohexane derivatives. *Annals of Tropical Medicine and Parasitology* 1993; 87:469-476.
442. Loiseau PM, Bourass J, Letourneux Y. Lymphotropic antifilarial agents derived from closantel and chlorambucil. *International Journal for Parasitology* 1997; 27:443-447.
443. Loke YW. Transfer of parasites across the placenta. *Advances in Parasitology* 1982; 21:155-228.
444. Lombange CK. Trends in sexually transmitted disease incidence in Papua New Guinea. *Papua New Guinea Medical Journal* 1984; 27:145-157.
445. Long GW, Rickman LS, Cross JH. Rapid diagnosis of *Brugia malayi* and *Wuchereria bancrofti* filariasis by an acridine orange/microhaemocrit tube technique. *Journal of Parasitology* 1990; 76:278-281.
446. Low K, Barker AJ, Vincent JM. A thirteen-year old with a painful swollen foot and eosinophilia. *Pediatric Infectious Diseases* 1995; 14:820, 825-826.
447. Lowman EW. Incidence of filariasis in children. *United States Navy Medical Bulletin* 1944; 52:341-343.
448. Lowrie RC, Eberhard ML, Lammie PJ, Raccurt CP, Katz SP, Duverseau YT. Uptake and development of *Wuchereria bancrofti* in *Culex quinquefasciatus* that fed on Haitian carriers with different microfilaria densities. *American Journal of Tropical Medicine and Hygiene* 1989; 41:429-434.
449. Lu AG, Valencia LB, Llagas L, Aballa L, Postrado L. Filariasis: A study of knowledge, attitudes and practices of the people of Sorsogon. *Social & Economic Research Project Reports No.1, TDR/WHO, Geneva* 1988.
450. Lucena WA, Dhalia R, Abath FGC, Licolas L, Regis LN, Furtado AF. Diagnosis of *Wuchereria bancrofti* infection by the polymerase chain reaction using urine and day blood from microfilaraemic patients. *Transactions of the Royal Society for Tropical Medicine and Hygiene* 1998; 92:290-293.
451. Lunde MN, Paranjape R, Lawley TJ, Ottesen EA. Filarial antigen in circulating immune complexes from patients with *Wuchereria bancrofti* filariasis. *American Journal of Tropical Medicine & Hygiene* 1988; 38:366-371.
452. Lutsch C, Cesbron JY, Henry D, Dessaint JP, Wandji K, Ismail M, Capron A. Lymphatic filariasis: detection of circulating and urinary antigen and differences in antibody isotypes complexed with circulating antigen between symptomatic and asymptomatic subjects. *Clinical & Experimental Immunology* 1988; 71:253-260.
453. Laurence BR, Simpson MG. Staining microfilariae for structures other than nuclei. *Transactions of the Royal Society of Tropical Medicine & Hygiene* 1969; 63:801-809.
454. Lynch GW. A note on the occurrence of filaria in Fijians. *Lancet* 1905; i:21.
455. MacKenzie S. A case of filarial haematochyluria. *Transactions of the Pathological Society of London* 1882; 33:394-410.
456. MacKenzie CD, Kron MA. Diethylcarbamazine: a review of its action in oncocerciasis, lymphatic filariasis and inflammation. *Tropical Diseases Bulletin* 1985; 82 (review): R1-R36.
457. Magnussen P, Makunde W, Simonsen PE, Meyrowitsch D, Jakubowski K. Chronic pulmonary disorders, including tropical pulmonary eosinophilia, in villages with endemic lymphatic filariasis in Tanga region and in Tanga town, Tanzania. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1995; 89:406-409.
458. Mahanty S, Luke HE, Kumaraswami V, Narayanan PR, Vijayshekar V, Nutman TB, Maizels RM, Denham DA, Sutanto I. Secreted and circulating antigens of the filarial parasite *Brugia pahangi*: analysis of in-vitro released components and detection of parasite products in-vivo. *Molecular and Biochemical Parasitology* 1985; 17:277-288.
459. Mahanty S, Day KP, Alpers MP, Kazura JW. Antifilarial IgG4 antibodies in children from filarial-endemic areas correlate with the duration of infection and are dissociated from antifilarial IgE antibodies. *Journal of Infectious Diseases* 1994; 170:1399-1343.
460. Mahanty S (a), Luke HE, Kumaraswami V, Narayanan PR, Vijayshekar V, Nutman TB. Stage-specific induction of cytokines regulates the immune response in lymphatic filariasis. *Experimental Parasitology* 1996; 84:282-290.
461. Mahanty S (b), Mollis S, Ravichandran M, Abrams JS, Kumaraswami V, Yaraman K, Ottesen EA, Nutman TB. High levels of spontaneous and parasite-driven interleukin-10 production are associated with antigen-specific hypo responsiveness in human lymphatic filariasis. *Journal of Infectious Diseases* 1996; 173:769-773.
462. Mahanty S, Ravichandran M, Raman U, Jayaraman K, Kumaraswami V, Nutman TB. Regulation of parasite antigen-driven immune responses by interleukin 10 (IL-10) and IL-12 in lymphatic filariasis. *Infection and Immunity* 1997; 65:1742-1747.
463. Maizels RM, Burke JA, Denham DA. Phosphoryl choline-bearing antigens in filarial nematode parasites: analysis of somatic extracts in *in-vivo* secretions and infection sera from *Brugia malayi* and *Brugia pahangi*. *Parasite Immunology* 1987; 9:49.
464. Maizels RM, Lawrence RA. Immunological Tolerance: the key feature in human filariasis. *Parasitology Today* 1991; 7:272-276.
465. Maizels RM, Denham DA. diethylcarbamazine (DEC): immuno pharmacological interactions if an antifilarial drug. *Parasitology* 1992; 105 (suppl):S49-S60.
466. Maizels RM, Sartono E, Kurniawan A, Partono F, Selkirk ME, Yazdanbakhsh M. T cell activation and the balance of antibody isotypes in human lymphatic filariasis. *Parasitology Today* 1995; 11:50-56.
467. Mak JW. Problems in filariasis control and the need for human behaviour and socio-economic research. *Southeast Asian Journal of Tropical Medicine & Public*

Health 1986; 17:479-485.

468. Mak JW, Navaratnam V, Ramachandran CP. Experimental chemotherapy of lymphatic filariasis. A review. *Annals of Tropical Medicine and Parasitology* 1991; 85:131-137.
469. Malhotra I, Ouma J, Wamachi A, Kioko J, Mungai P, Omollo A, Elson L, Koech D, Kazura JW, King CL. *In-utero* exposure to helminth and mycobacterial antigens generates cytokine responses similar to that observed in adults. *Journal of Clinical Investigation* 1997; 99:1759-1766.
470. Malhotra I, Mungai P, Wamachi A, Kioko J, Ouma J, Kazura JW, King CL. Helminth and Calmette-Guerin-induced immunity in children sensitised in utero to filariasis and schistosomiasis. *Journal of Immunology* 1999; 162:6843-6848.
471. Manfredi R, Mastroianni A, Coronado OV, Chiodo F. Imported tropical diseases in subjects with HIV infection. *Panminerva Medicine* 1998; 40:72-74.
472. Manson P (a). Further observations of *Filaria sanguinis hominis*. China Imperial Maritime Customs. Half Yearly Medical Report for 1877 pages 1-26, published 1878.
473. Manson P (b). On the development of *Filaria sanguinis hominis* and on the mosquito considered as a nurse. *Journal of the Linnean Society (Zoology)* 1878; 14:14:304-311.
474. Manson P (a) Additional notes on *Filaria sanuinis hominis* and filarial disease. China Imperial Customs. Half Yearly Medical Report for 1879 pages 31-51, published 1880
475. Manson P (b). Further observations on microfilariae, with descriptions of new species communicated (with a prefatory note) by the president. *Journal of the Quekett Microscopical Club* 1880; 6:130-140.
476. Manson P(a). Additional notes on *Filaria sanguinis hominis* and filaria disease. Discovery of adult *F. bancrofti*. China Imperial Maritime Customs, Half Yearly Medical Report for 1880 pages 13-15, published 1881.
477. Manson P(b). Lymph scrotum showing filaria *in situ*. *Transactions of the Royal Pathological Society of London* 1881; 32:285-302.
478. Manson P. Notes on filaria disease. China Imperial Maritime Customs, Medical Reports for the Half Year 1882; 23:1-16.
479. Manson P. The *Filaria sanguinis hominis* and certain new forms of parasitic diseases in India, China and certain warm countries. 1883, page 186, publisher unknown.
480. Manson P. The metamorphosis of *Filaria sanguinis hominis* in the mosquito. *Transactions of the Linnean Society of London* 1884; series 2,3:367-388.
481. Manson P. On filarial periodicity. *British Medical Journal* 1899; ii:644-646.
482. Manson-Bahr PH, Alcock A. The life and works of Sir Patrick Manson, Cassell and Company, London 1927, page 273. Manson-Bahr P. The story of *Filaria bancrofti*. *Journal of Tropical Medicine and Hygiene* 1959; 52:61; 84-94; 106-117; 138-144; 160-173.
483. Manson-Bahr PH. Patrick Manson as a parasitologist: a critical review. In; *International Review of Tropical Medicine*, edited by Linicombe DR, Academic Press New York, 1961, pages 77-129.
484. Mantovani A. Transplacental transmission of *Dirofilaria immitis* in the dog. *Journal of Parasitology* 1966; 52:116.
485. Marley SE, Lammie PJ, Eberhard ML, Hightower AW. Reduced antifilarial IgG4 responsiveness in a sub-population of microfilaraemic persons. *Journal of Infectious Diseases* 1995; 172:1630-1633.
486. Marley SE, Lammie PJ, Eberhard ML. Immuno purification and measurement of IgE in serum samples from bancroftian filariasis. *Journal of Parasitology* 1996; 82:178-181.
487. Mataika JU, Kimura E, Koroivuetu J, Shimada M. Efficacy of five annual single doses of diethylcarbamazine for treatment of lymphatic filariasis in Fiji. *Bulletin of the World Health Organisation* 1998; 76:575-579.
488. Mathai E, David S. Intra ocular filariasis due to *Wuchereria bancrofti*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2000; 94:317-318
489. Mavoungou D, Lansoud-Soukate J, Dupont A. Steroid and gonadotropin hormone levels in young African women with filarial infection. *Journal of Steroid Biochemistry* 1989; 43:577-580.
490. Maxwell CA, Curtis CF, Kisumku S, Ali Yahya, Issa Thalib A, Haji H. Integrated control of filariasis including supression of the vector using expanded polystyrene beads. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1989; 428.
491. Maxwell CA, Mohammed K, Kismuku U, Curtis CF. Can vector control play a useful supplementary role against bancroftian filariasis. *Bulletin of the World Health Organisation* 1999; 77:138-143
492. McAdam KPWJ. Leprosy, filariasis and malaria as causes of secondary amyloidosis in the tropics. *Papua New Guinea Medical Journal* 1978; 69-78.
493. McCarthy JS, Zhong M, Gopinath R, Ottesen EA, Williams SA, Nutman TB. Evaluation of a polymerase chain reaction-based assay for diagnosis of *Wuchereria bancrofti*. *Journal of Infectious Diseases* 1996; 173:1510-1514.
494. McGreevy PB, Ratiwayato S, Tuti S, McGreevy MM, Dennis DT. *Brugia malayi*: relationship between anti-sheath antibodies and amicrofilaraemic in natives living in an endemic area of south Kalimantan, Borneo. *American Journal of Tropical Medicine & Hygiene* 1980; 29:553-562.
495. McKinley EB. The role of bacteria in acute filarial lymphangitis. *Puerto Rico Journal of Public Health* 1931; 6:419.
496. McMahon JE. Preliminary screening of antifilarial activity of levamisole and amodiaquine on *Wuchereria bancrofti*. *Annals of Tropical Medicine and Parasitology* 1979; 73: 465-472.
497. McMahon JE, Simonsen PE. Filariasis. In Manson's tropical diseases, 20th edition edited by Cook GC. W.B. Saunders Company London 1996. Chapter 70.
498. McMillan B. Lawang bark as a rubefacient in the treatment of filarial lymphangitis is Papua New Guinea. *Medical Journal of Australia* 1968; 2:63-64.
499. McQuay RM. Parasitological studies in a group of furloughed missionaries. *American Journal of Tropical Medicine and Hygiene* 1967; 16:161-166.
500. McQuay RM. Citrate-saponin-acid method for the recovery of microfilariae from blood. *American Journal of Clinical Pathology* 1970; 54:743-746.
501. Mejia CR, Hernandez JE, Kitchen LW. DEC treatment of non-nematodal infections. *Tropical Doctor* 1996; 26:55-57.
502. Meyrowitsch D, Simonsen PE, Makunde WH.. Mass diethylcarbamazine therapy for the control of Bancroftian filariasis through community participation: comparative efficiency of a low monthly dose and medicated salt.

- Transactions of the Royal Society of Tropical Medicine and Hygiene 1996; 90:74-79.
503. Meyrowitsch D, Simonsen PE. Long-term effect of mass diethylcarbamazine chemotherapy on bancroftian filariasis: results at four years after start of treatment. Transactions of the Royal Society of Tropical Medicine and Hygiene 1998; 92:98-103.
 504. Michael E, Grenfell BT, Bundy DAP. The association between microfilaraemia and disease in lymphatic filariasis. Proceedings of the Royal society of London. Series B. Biological Sciences 1994; 256:33-40.
 505. Michael E (a), Bundy DAP, Grenfell BT. Re-assessing the global prevalence and distribution of lymphatic filariasis. Parasitology 1996; 112:409-428.
 506. Michael E (b), Meyrowitsch DW Simonsen PE. Cost and cost effectiveness of mass diethylcarbamazine chemotherapy for the control of Bancroftian filariasis: comparison of four strategies in Tanzania. Tropical Medicine and International Health 1996; 1:414-426.
 507. Michael E, Bundy DAP. Global mapping of lymphatic filariasis. Parasitology Today 1997; 13:472-476.
 508. Michael E Bundy DA. Herd immunity to filarial infection is a function of vector biting rate. Proceedings of the Royal society of London. Series B. Biological Sciences 1998; 265:855-860.
 509. Michael E. The control of human filariasis. Current Opinion in Infectious Diseases 1999; 12:565-578.
 510. Michael P. Filariasis Among Navy and Marine Personnel: Report on Laboratory Investigations. United States Naval Medical Bulletin 1944; 42:1059-1074.
 511. Miller S, Snowden K, Schreuer D, Hammerburg B. Selective Breeding of Dogs for Segregation of Limb Edema From Microfilaria As Clinical Manifestations of Brugia Infections. American Journal of Tropical Medicine and Hygiene 1990; 43:489-497.
 512. Misra S, Mohapatra TM, Rathaur S. Wuchereria bancrofti: identification of parasitic acetylcholinesterase in microfilariae-infected human sera. Tropical Medicine & Parasitology 1993; 44:75-78.
 513. Mistry SS, Subrahmanyam D. Cellular Immune Competence in Bancroftian filariasis. Journal of Tropical Medicine and Hygiene 1985; 88:383-388.
 514. Mitsui Y, Tanamori H, Kitagawa T, Fujimaki Y, Aoki Y. Simple and sensitive enzyme-linked immunosorbent assay for ivermectin. American Journal of Tropical Medicine and Hygiene 1996; 54:243-248.
 515. Mitsui Y, Aoki Y. A direct competitive ELISA for the simple, sensitive and accurate determination of diethylcarbamazine concentration in serum. International Journal of Parasitology 1998; 28:623-626.
 516. Mohamed NH, Rifaat MA, Morsy TA, Bebars MA, Faiad MH. Concomitant infection with filariasis and schistosomiasis in an endemic area in Sharkya Governorate, Egypt. Journal of the Egyptian Society of Parasitology 1983; 13:403-406.
 517. Mohamed NH, el Okby LM, Abdel Baky MH, el Sibai MM, Fawzy AF, Sarwat MA, Fayad ME, Kamal AM. Study of the relationship between tissue typing and occurrence of filariasis and its complications in Egypt. Journal of the Egyptian Society of Parasitology 1987; 17:751-758.
 518. Mohamed NH (a), el Okby LM, Abdel Baky MH, el Sibai MM, Fawzy AF, Sarwat MA, Fayad ME, Kamal AM. The relation between red cell antigens and the occurrence of filariasis and its complications. Journal of the Egyptian Society of Parasitology 1989; 19:75-80.
 519. Mohamed NH, Hamadto HHA, El-Fakahany AF, Farrag AMK, El-Hamshary AMS. Study of human and parasitic factors in relation to bancroftian filariasis in Egypt. Journal of the Egyptian Society of Parasitology 1992; 22:719-729.
 520. Monzon RB. Traditional medicine in the treatment of parasitic diseases in the Philippines. Southeast Asian Journal of Tropical Medicine and Hygiene 1995; 26:421-428.
 521. More SJ, Copeman DB. A highly specific and sensitive monoclonal antibody-based ELISA for the detection of circulating antigen in Bancroftian filariasis. Tropical Medicine & Parasitology 1990; 41:403-406.
 522. More SJ, Copeman DB. Antigen detection ELISA: pre-treatment of serum to reduce interference by specific host antibodies. Tropical Medicine & Parasitology 1991; 42:91-94.
 523. Moore TA, Reynolds JC, Kenney RT, Johnston W, Nutman TB. Diethylcarbamazine-induced reversal of early lymphatic dysfunction in a patient with Bancroftian filariasis - assessment with lymphoscintigraphy. Clinical Infectious Diseases 1996; 23:1007-1011.
 524. Moore TA, Nutman TB. Eosinophilia in the returning traveller. Infectious Disease Clinics of North America 1998; 12:503-508.
 525. Moulia-Pelat JP, Glaziou P, Nguyen-Ngoc L, Cardines R, Spiegel A, Cartel JL. A comparative study of detection methods for evaluation of microfilaraemia in lymphatic filariasis control programmes. Tropical Medicine & Parasitology 1992; 43:146-148.
 526. Moulia-Pelat JP (a), Nguyen LN, Glaziou P, Chanteau S, Gay VM, Martin PMV, Cartel JL. Safety trial of single-dose treatments with a combination of ivermectin and diethylcarbamazine in Bancroftian filariasis. Tropical Medicine and Parasitology 1993; 44:79-82.
 527. Moulia-Pelat JP (b), Glaziou P, Chanteau S, Nguyen-Ngoc L, Marcet Y, Cardines R, Martin PMV, Cartel JL. Periodicity of Wuchereria bancrofti var Pacifica filariasis in French Polynesia. Tropical Medicine and Parasitology 1993; 44:83-85.
 528. Melrose WD, Turner PF, Pisters P, Turner B. An improved Knott's concentration test for the detection of microfilariae. Transactions of the Royal Society of Tropical Medicine and Hygiene 2000; 94:176.
 529. Moulia-Pelat JP (a), Nguyen LN; Glaziou P, Chanteau S, Plichart R, Beylier I, Martin PMV, Cartel JL. Ivermectin 400 µg/kg: long-term suppression of microfilaraemia in Bancroftian filariasis. Transactions of the Royal Society of Tropical Medicine and Hygiene 1994; 88:107-109.
 530. Moulia-Pelat JP (b), Nguyen LN, Glaziou P, Chanteau S, Ottesen EA, Cardines R, Martin PMV, Cartel J. Ivermectin plus diethylcarbamazine: an additive effect on early microfilariae clearance. American Journal of Tropical Medicine and Hygiene 1994; 50:206-209.
 531. Moulia-Pelat JP, Glaziou PH, Weil GJ, Nguyen LN, Gaxotte PH, Nicolas L. Combination ivermectin plus diethylcarbamazine, a new effective tool for control of lymphatic filariasis. Tropical Medicine and Parasitology 1995; 46:9-12.
 532. Moulia-Pelat JP, Nguyen LN, Hascoet H, Nicolas L. Combinations of ivermectin and diethylcarbamazine for improved control of lymphatic filariasis. Parasite 1996; 3:45-48.
 533. Muhondwa EPY. Community involvement in filariasis control: the Tanzania experiment. World Health Organisation Report, Geneva 1983.
 534. Mujinja PGM, Gasarasi DB, Premji ZG, Nguma J. Social and economic impact of lymphatic filariasis in Rufiji district, Southeast Tanzania. Lymphatic Filariasis Research and Control in Africa: report on a workshop

- held in Tanga, Tanzania. Danish Bilharziasis Laboratory, Denmark & National Institute for Medical Research, Tanzania, 1997.
535. Mukherjee SK, Mukherjee AK, Sen JK. Filariasis with recurrent haematuria. *Journal of the American Medical Association* 1974; 230:1254-1255.
 536. Mukherjee M, Misra S, Chatterjee RK. Optimization of test conditions for the development of MTT as an in-vitro screen. *Indian Journal of Experimental Biology* 1997;35:73-76.
 537. Mukhopadhyay S, Ravindran B. Antibodies to diethylcarbamazine potentiate the antifilarial activity of the drug. *Parasite immunology* 1997; 19:191-195.
 538. Murthy PK, Kapil A, Sircar AR, Katiyar JC, Saxcena KC. Avidin-biotin enzyme-linked immunosorbent assay (AB_ELISA) for the detection of Bancroftian filariasis. *Annals of Tropical Medicine & Parasitology* 1993; 87:299-302.
 539. Murthy PK, Mehrotra M, Tyagi K, Chaturvedi UC, Chatterjee RK, Response of IgG sub-classes to diethylcarbamazine therapy in bancroftian filarial patients. *Current Science (Bangalore)* 1997; 72:265-268.
 540. Myers WW. Further observations of *Filaria sanguinis hominis* in South Formosa. *China Imperial Maritime Customs. Medical Reports for the Half Year 1881*; 21:1-21.
 541. Myung K, Massougoudji A, Ekoue S, Atchade P, Kikifagla V, Klion AD. Lymphatic filariasis in a hyper endemic region- a ten-year follow-up panel survey. *American Journal of Tropical Medicine and Hygiene* 1998; 59:222-226.
 542. Nguyen NL, Moulia-Pelat JP, Glaziou P, Martin PMV, Cartel JL. Advantages of ivermectin at a single dose of 400ug/kg compared with 100ug/kg compared with 100ug/kg for community treatment of lymphatic filariasis in Polynesia. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1994; 88:461-464.
 543. Nanduri J, Kazura JW. Clinical and laboratory aspects of filariasis. *Clinical Microbiology* 1989; 2:39-50.
 544. Napier LE. Filariasis in U.S. marines (Editorial). *Journal of Tropical Medicine and Hygiene* 1947; 50:189.
 545. Narasimham MVVL, Sharma SP, Sundram RM, Reddy GS, Raina VK, Sambasivam V, Das M. Control of Bancroftian filariasis by diethylcarbamazine-medicated common salt in Karaikal, Pondicherry, India. *Journal of Communicable Diseases* 1989; 21:157-170.
 546. Narayanan PR, Vanamala CR, Alamelu R, Kumaraswamy V, Tripathy SP, Prabhakar R. Reduced lymphocyte response to mitogens in patients with Bancroftian filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1986; 80:78-84.
 547. Nar NT, Bodghdadi AM, Allam KA, El-Adawi AI, Soliman MI. Effect of ivermectin on survival and fecundity of *Culex pipiens* the vector of *Wuchereria bancrofti*. *Journal of the Egyptian Parasitology Society* 1996; 26:161-168.
 548. Nathan MB, Lambourne A, Monteil S. Evaluation of a membrane (Nucleopore) filtration method using capillary blood for the detection of microfilariae. *Annals of Tropical Medicine and Parasitology* 1982; 76:339-345.
 549. Negesse Y, Lanoie OO, Nearsh RO, Conner DH. Loasis: Calabar swellings and involvement of deep organs. *American Journal of Tropical Medicine and Hygiene* 1985; 34:537-546.
 550. Needham CS, Lilywhite JE, Beasley NMR, Didier JM, Kihamia CM, Bundy PA. Potential for diagnosis of intestinal nematode infections through antibody detection in saliva. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1996; 90:526-530.
 551. Neisius B. Inaugural medical debate on elephantiasis or lepra arabum (filarial elephantiasis), Strassburg 1673. *Annals of Medical History* 1927; 9:267-276.
 552. Nelson GS. Filarial infections as Zoonosis. *Journal of Helminthology* 1965; 39:229-250.
 553. Nelson P, Sellon R, Novotney C. Therapeutic effects of diethylcarbamazine and 3'-azido-3'-deoxythymidine on feline leukaemia virus lymphoma formation. *Veterinary Immunology and Immunopathology* 1995; 46:181-194.
 554. Neves HA, Scaff LM. *Microfilaria congenita*. *Revista Brasileira de Malariologia e Doencas Tropicais* 1952; 4:415-421.
 555. Neves HA, Scaff LM. *Comprovacao da Microfilaria congenita de Wuchereria bancrofti*. *Revista Brasileira de Malariologia e Doencas Tropicais* 1954; 6:415-421.
 556. Ngu JL, Chatelanat F, Leke R, Ndunde P, Youmbissi J. Nephropathy in Cameroon: evidence for filarial derived immune-complex pathogenesis in some cases. *Clinical Nephrology* 1985; 24:128-134.
 557. Nguyen NL, Moulia-Pelat JP, Cartel JL. Control of lymphatic filariasis in an endemic area of Polynesia by ivermectin 400 micrograms/kg. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1996; 90:689-691.
 558. Nguyen NL, Plichart C, Esterre P. Assessment of immuno chromatographic test for rapid lymphatic filariasis diagnosis. *Parasite-Journal de la Societe Francaise do Parasitologie* 1999; 6:355-358.
 559. Nicolas L, Luquiaud P, Lardeux F, Mercer DR. A polymerase chain reaction assay to determine infection of *Aedes polynesiensis* by *Wuchereria bancrofti*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1996;136-139..
 560. Nicolas L, Plichart C, Nguyen NL, Moulia-Pelat JP. Reduction of *Wuchereria bancrofti* adult worm circulating antigen after annual treatments of diethylcarbamazine combined with ivermectin in French Polynesia. *Journal of Infectious Diseases* 1997; 175:489-492.
 561. Nicolas L, Lang S, Plichart C, Deparia X. Filarial antibody responses in *Wuchereria bancrofti* transmission area are related to parasitological but not clinical status. *Parasite Immunology* 1999; 21; 73-80.
 562. Nilsen TW, Maroney PA, Goodwin RG, Perrine KG, Denker AJ, Nanduri J, Kazura JW. Cloning and characterisation of a potentially protective antigen in lymphatic filariasis. *Proceedings of the National Academy of Science USA* 1988; 85:3606-3607.
 563. Noroes J, Addiss D, Santos A, Medeiros Z, Coutinho A, Dreyer G. Ultrasonic evidence of abnormal lymphatic vessels in young men with adult *Wuchereria bancrofti* infection in the scrotal area. *Journal of Urology* 1996; 156: 409-412.
 564. Nooruddin, Ahmed SS. The effects of splenectomy on parasitic infections, and the role of the spleen in filariasis - a brief appraisal of our current knowledge. *Journal of Tropical Medicine and Hygiene* 1967; 70:229-231.
 565. Nussenzweig R. Parasitic disease as a cause of immuno suppression. *New England Journal of Medicine* 1982; 306:423-426.
 566. Nutman TB, Ottesen EA, Fauci AS, Volkman DJ. Parasite antigen-specific human T cell lines and clones. Major histocompatibility complex restriction and B cell helper function. *Journal of Clinical Investigation* 1984; 73:1754-1762.

567. Nutman TB (a), Kumaraswami V, Pao L, Narayanan PR, Ottesen EA. Analysis of *in-vitro* B cell immune responsiveness in human lymphatic filariasis. The Journal of Immunology 1987; 138:3954-3959.
568. Nutman TB (b), Kumaraswami V, Ottesen EA. Parasite-anergy in human filariasis: insights after analysis of parasite antigen-driven lymphokine production. Journal of Clinical Investigation 1987; 79:1516-1523.
569. Nwoke BEB, Nduka FO, Okereke OM, Ehighibe OC. Sustainable urban development and human health: septic tank as a major breeding habitat of mosquito vectors of human disease. Applied Parasitology 1993; 34:1-10.
570. Olszewski W, Jamal S, Manokaran G, Lukomska B, Kubicka U. Skin changes in filarial and non-filarial lymphodema of the lower extremities. Tropical Medicine and Parasitology 1993; 44:40-41.
571. Olszewski W, Jamal S. Skin bacterial factor in progression of filarial lymphodema. Lymphology 1994; 27:148-149.
572. Olszewski W, Jamal S, Manokaran G, Pani S, Kumaraswami V, Kubicka U, Lukomska B, Dworzczynski A, Swoboda E, Meisel-Mikolajczyk F. Bacteriologic studies of skin, tissue fluid, lymph and lymph nodes in patient lymphodema. American Journal of Tropical Medicine and Hygiene 1997; 57:7-15.
573. Olszewski W, Jamal S, Manokaran G, Pani S, Kumaraswami V, Kubicka U, Lukomska B, Swoboda E, Meisel-Mikolajczyk F, Stelmach E, Zaleska M. Bacteriological studies of blood, tissue fluid, lymph and lymph nodes in patients with dermatolymphangioadenitis (DLA) in course of filarial lymphedema. Acta Tropica 1999; 73:217-214.
574. O'Connor FW. The aetiology of the disease syndrome in *Wuchereria bancrofti* infections. Transactions of the Royal Society of Tropical Medicine & Hygiene 1932; 26:13-16.
575. O'Holohan DR, Zaman V. Treatment of *Brugia malayi* infection with levamisole. Journal of Tropical Medicine & Hygiene 1974; 77:113-115.
576. Oliveira DBG. Membranous nephropathy: an IgG4-mediated disease. Lancet 1998; 351:670-672.
577. Oothuman P, Denham DA, McGreevy PB, Nelson GS, Rogers R. Successful vaccination of cats against *Brugia malayi* with larva attenuated by irradiation with 10 rad of Cobalt 60. Parasite Immunity 1979; 1:209-215.
578. Orihel TC. Cerebral filariasis in Rhodesia - a zoonotic infection? American Journal of Tropical Medicine and Hygiene 1973; 22:596-599.
579. Orihel TC, Eberhard ML. Zoonotic filariasis. Clinical Microbiology Reviews 1998; 11:366-381.
580. Ormerod AD, Petersen J, Husey JK, Weir J, Edward N. Immune complex glomerulonephritis and chronic anaerobic urinary infection - complications of filariasis. Postgraduate Medical Journal 1983; 59:730-733.
581. Otero PM, Lebron AP. Immunological responses of cases of recurrent tropical lymphangitis to haemolytic streptococci and their products. Puerto Rico Journal of Public Health 1936; 12:43-45.
582. Ottesen EA, Weller PF, Heck L. Specific cellular unresponsiveness in human filariasis. Immunology 1977; 33:413-421.
583. Ottesen EA. Immunopathology of lymphatic filariasis in man. Springer Seminars in Immunopathology 1980; 2:373-385.
584. Ottesen EA, Mendell NR, MacQueen JM, Weller PF, Amos DB, Ward FE. Familial predisposition to filarial infection - not linked to HLA-A or -B locus specificities. Acta Tropica 1981; 38:205-216.
585. Ottesen EA, Weller PF, Lunde MN, Hussain R. Endemic filariasis on a Pacific island 2. Immunologic aspects: Immunoglobulin, complement, and specific antifilarial IgG, IgM, and IgE antibodies. American Journal of Tropical Medicine and Hygiene 1982; 31:953-961.
586. Ottesen EA. Immunological aspects of lymphatic filariasis and onchocerciasis in man. Transactions of the Royal Society of Tropical Medicine & Hygiene 1984; 78 (supplement):9-18.
587. Ottesen EA. Efficacy of diethylcarbamazine in eradicating infection with lymphatic-dwelling filariae in humans. Review of Infectious Diseases 1985; 7:341-356.
588. Ottesen EA (b), Skvaril F, Tripathy SP, Poindexter RW, Hussain R. Prominence of IgG4 in the antibody response to human filariasis. Journal of Immunology 1985; 134:2707-2712.
589. Ottesen EA. Filariasis now. American Journal of Tropical Medicine & Hygiene 1989; 41 (supplement): 9-17).
590. Ottesen EA, Kumaraswami V, Vijayasekaran V. Controlled trial of ivermectin and diethylcarbamazine in lymphatic filariasis. New England Journal of Medicine 1990; 322; 1113-1117.
591. Ottesen EA. Infection and disease in lymphatic filariasis: an immunologic perspective. Parasitology 1992; 104:S71-S79.
592. Ottesen EA. Filarial infections. Infectious Disease Clinics of North America 1993; 7:619-633.
593. Ottesen EA. The human filariases: new understandings, new therapeutic strategies. Current Opinion in Infectious Diseases 1994; 7:550-558.
594. Ottesen EA, Campbell WC. Ivermectin in human medicine. Journal of Antimicrobial Chemotherapy 1994; 34:195-203.
595. Ottesen EA. Immune responsiveness and the pathogenesis of human onchocerciasis. The Journal of Infectious Diseases 1995; 171:659-671.
596. Ottesen EA, Ramachandran CP. Lymphatic filariasis infection and disease: control strategies. Parasitology Today 1995; 11:129-131.
597. Ottesen EA, Ismail MM, Horton J. The role of Albendazole in programmes to eliminate lymphatic filariasis. Parasitology Today 1999; 15:382-386.
598. Otto GF, Jachowski LA, Wharton JD. Filariasis in American Samoa. III. Studies on chemotherapy against the non-periodic form of *Wuchereria bancrofti*. American Journal of Tropical Medicine and Hygiene 1953; 2:495-516.
599. Paily KP, Gomathi K, Hoti SL, Balaraman K, Dhanda V. A report on the demonstration of microfilariae of *Brugia malayi* in the brain of an experimental animal host *Mastomys natalensis*. International Journal of Parasitology 1995; 1123-1126.
600. Palmieri JR, Conner DH, Purnomo DT, Marwoto HA. Experimental infection of *Wuchereria bancrofti* in the silvered leaf monkey *Presbytis cristatus* Eschscholtz 1821. Journal of Helminthology 1982; 56:243-245.
601. Pampiglione S, Canestri Trotti G, Rivasi F. Human dirofilariasis due to *Dirofilaria (Noctiella) repens*: a review of the world literature. Parassitologia 1995; 37:149-193.
602. Pancre V, Cesbron JY, Aurialt C, Joseph M, Chandenier J, Capron A. IgE-dependent killing of *Brugia malayi* microfilaraemia by human platelets and

- its modification by T cell products. International Archives of Allergy and Applied Immunology 1988; 85:483-486.
603. Pani SP, Balakrishnan N, Srividya A, Bundy DA, Grenfell BT. Clinical epidemiology of bancroftian filariasis: effect of age and gender. Transactions of the Royal Society of Tropical Medicine & Hygiene 1991; 85:260-264.
 604. Pani SP, Srividya A. Clinical manifestations of Bancroftian filariasis with special reference to lymphodema grading. Indian Journal of Medical research 1995; 102:114-118.
 605. Pani SP, Hoti SL, Elango A, Yuvaraj J, Lall R, Ramaiah KD. Evaluation of the ICT whole blood antigen card test to detect infection due to nocturnally periodic *Wuchereria bancrofti* in South India. Tropical Medicine and International Health 2000; 5:359-363.
 606. Panicker KN, Sabesan S. Economic effects of filariasis. TDR News No. 34, TDR Geneva 1990.
 607. Panicker KN, Krishnamoorthy K, Sabesan S, Prathiba J, Abidha K. Comparison of effects of mass annual and biannual single dose therapy with diethylcarbamazine for the control of Malayan filariasis. Southeast Asian Journal of Tropical Medicine & Public Health 1991; 22:402-411.
 608. Paranjape RS, Kumaraswami V, Prabhakar R, Narayanan PR. Anti-filarial IgG antibodies in patients with Bancroftian filariasis and tropical pulmonary eosinophilia. Indian Journal of Experimental Biology 1985; 23:676-678.
 609. Partono F, Purnomo A, Dennis DT, Atmosoedjono S, Cross JH. *Brugia timori* sp. n. (Nematoda: Filarioidea) from Flores island, Indonesia. Journal of Parasitology 1977; 63:540-546.
 610. Partono F, Idris KN. Some factors influencing the loss of microfilariae from stained thick blood films. Southeast Asian Journal of Tropical Medicine and Public Health 1977; 8:158-163.
 611. Partono F, Purnomo W, Pribadi W, Soewarta A. Epidemiological features of *Brugia timori* in a newly established village, Karakuak, West Flores, Indonesia. American Journal of Tropical Medicine & Hygiene 1978; 27:910-915.
 612. Partono F, Purnomo A, Oemijati S, Soewarta A. The long term effects of repeated diethylcarbamazine administration with special reference to microfilaraemia and elephantiasis. Acta Tropica 1984; 38:217-225.
 613. Partono F. The spectrum of disease in lymphatic filariasis. In: Filariasis: Ciba Foundation Symposium No. 127. Edited by Ottesen EA. John Wiley and Sons, New York 1987.
 614. Partono F, Maizels RM, Puromo S. Towards a filarial-free community: evaluation of filariasis control over an eleven year period in Flores, Indonesia. Transactions of the Royal Society of Tropical Medicine and Hygiene 1989; 83:821-826.
 615. Pearlman E, Heinzel FP, Hazlett FE, Kazura JW. IL-12 modulation of T helper responses to the filarial helminth *Brugia malayi*. Journal of Immunology 1995; 154:4658-4664.
 616. Peruzzi M. Cerebral filariasis in five monkeys infected with trypanosomal meningo-encephalitis. Final report of the League of Nations International Commission on Human Trypanosomiasis 1928; pages 309-313.
 617. Petithory JC, Ardoin F, Ash LR, Vandemeulebroucke E, Galeazzi G, Dufour M, Paugam A. Microscopic diagnosis of blood parasites following a cytoconcentration technique. American Journal of Tropical Medicine and Hygiene 1997; 57:637-642.
 618. Petralanda I, Yarzabel L, Piessens WF. Parasite antigens are present in breast milk of women infected with *Onchocerca volvulus*. American Journal of Tropical Medicine and Hygiene 1988; 38:372-379.
 619. Petrocheilou V, Theodorakis M, Williams J, Prifti H, Geogilas K, Apostolopoulou I, Mavrikakis M. Microfilaraemia from a *Dirofilaria*-like parasite in Greece. Acta Pathologica, Microbiologica et Immunologica Scandinavica 1998; 106:315-318.
 620. Phantana S, Sensathien S, Songtrus J, Klagrathoke S, Phongnin. ICT filariasis test: a new screening test for bancroftian filariasis. Southeast Asian Journal of Tropical Medicine and Public Health 1999; 30:47-51.
 621. Philip M, Davis TB, Storey N, Callow C.S. Immunity in filariasis: perspectives for vaccine development. Annual Review of Microbiology 1988; 42:685-716.
 622. Piessens WF (a), McGreevy PB, Radiant S, McGreevy M, Piessens PW, Caiman I, Ceras S, Dennis DT. Immune responses in human infections with *Brugia malayi*: specific cellular unresponsiveness to filarial antigens. Journal of Clinical Investigation 1980; 65:172-179.
 623. Piessens WF (b), McGreevy PB, Radiant S, McGreevy M, Piessens PW, Caiman I, Ceras S, Dennis DT. Immune responses in human infections with *Brugia malayi*: correlation of cellular and humoral reactions to microfilarial antigens with clinical status. American Journal of Tropical Medicine & Hygiene 1980; 31:953-957.
 624. Piessens W, Partono F. Host-vector-parasite relationships in human filariasis. Seminars in Infectious Disease 1980; 3:131-152.
 625. Piessens WF (d), McGreevy PB, Radiant S, McGreevy M, Piessens PW, Caiman I, Ceras JS, Dennis DT. Immune responses in human infections with *Brugia malayi*: correlation of cellular and humoral reactions to microfilarial antigens with clinical status. American Journal of Tropical Medicine and Hygiene 1980; 29:563-570.
 626. Piessens WF. Lymphatic filariasis in humans: an immunologic maze. Annals of Internal Medicine 1981; 95:779-779.
 627. Piessens WF, Ratiwayato S, Piessens PW, Tuti S, McGreevy PB, Darwis F, Palmieri JR, Caiman I, Dennis DT. Effect of treatment with diethylcarbamazine on immune responses to filarial antigens in patients infected with *Brugia malayi*. Acta Tropica 1981; 38:227-234.
 628. Piessens WF, Partono F, Hoffman SL, Ratiwayato S, Piessens PW, Palmieri JR, Caiman I, Dennis DT, Carney WP. Antigen-specific suppressor T lymphocytes in human lymphatic filariasis. New England Journal of Medicine 1982; 307:144-148.
 629. Piessens WF, Hoffman SL, Radiant S, Piessens PW, Partono F, Kurniawan L, Marwoto HA. Opposing effects of filariasis and chronic malaria on immunoregulatory T lymphocytes. Diagnostic Immunology 1983; 1:157-160.
 630. Pinkston P, Vijayan VK, Nutman TB, Rom WN, O'Donnell K, Cornelius MJ, Kumaraswami V, Ferrans VJ, Takemura T, Yenokida G, Thiruvengadam KV, Tripathy SP, Ottesen EA, Crystal RG. Acute tropical eosinophilia: characterisation of the lower respiratory tract inflammation and its response to therapy. Journal of Clinical Investigation 1987; 80:216-225.
 631. Pires ML, Santos Ferreira R, Araujo S. Investigacao sobre a passagem transplacentaria de microfilarias de *Wuchereria bancrofti* em recém-nascidos de maes microfilaemicas. Revisto do Instituto de Medicina Tropical de Sao Paulo 1991; 33:91-95.
 632. Piyaratn P, Pradatsundarasar A. Microfilarial

- granuloma of the spleen. American Journal of Tropical Medicine & Hygiene 1961; 10:838-842.
633. Plaisier AP, Subramanian S, Das PK, Souza W, Lapa T, Furtado AF, Vanderploeg CPB, Habbema JDF, Vanoortmarssen GJ. The LYMFASIM simulation program for modelling lymphatic filariasis and its control. Methods of Information in Medicine 1998; 37:97-108.
 634. Plaisier AP, Cao WC, Van Oortmarssen GJ, Habbema JDE. Efficacy of ivermectin in the treatment of *Wuchereria bancrofti* infection: a model-based analysis of trial results. Parasitology 1999; 119:385-394.
 635. Plier DA, Mai e Silva MC, Maciel MA, Freedman DO. T-lymphocytes from individuals with filarial inflammatory disease have increased transendothelial migration *in vitro*. Clinical Immunology and Immunopathology 1997; 82:216-220.
 636. Poddar SK, Misra S, Singh NK. Acute polymyositis associated with *Wuchereria bancrofti*. Acta Neurologica Scandinavica 1994; 89:225-226.
 637. Portaro JK, Britten S, Ash LR. *Brugia pahangi*: depressed mitogen reactivity in filarial infections in the jird *Meriones unguiculatus*. Experimental Parasitology 1976; 40:438-446.
 638. Poynton JO, Hodgkin EP. Two microfilaria of the Kra monkey (*Macaca irus*). Transactions of the Royal Society of Tropical Medicine and Hygiene 1939; 32:555-556.
 639. Prasad GB (a), Reddy MV, Harinath BC. Detection of filarial antigen in immune complexes in Bancroftian filariasis by ELISA. Indian Journal of Medical Research 1983; 78:780-783.
 640. Prasad GB (b), Kharat I, Harinath BC. Detection of anti microfilarial ES antigen-antibody in immune complexes in Bancroftian filariasis by enzyme immuno assay. Transactions of the Royal Society of Tropical Medicine & Hygiene 1983; 77:771-772.
 641. Prasad GB (c), Reddy MRV, Harinath BC. Immune complexes and immunoglobulins involved in human filariasis. Indian Journal of Medical Research 1983; 77:813-816.
 642. Prasad GB, Harinath BC. Immuno suppression in clinical filariasis. Indian Journal of Medical Research 1987; 85:531-535.
 643. Prasad GB, Harinath BC. Analysis, characterisation and diagnostic utility of filarial antigen fractions isolated from immune complexes in Bancroftian filariasis. Diagnostic & Clinical Immunology 1988; 5:269-275.
 644. Prasad GB, Kharat I, Harinath BC. Suppression of Con A induced lymphocyte transformation by sera from filarial patients with clinical manifestations. Indian Journal of Experimental Biology 1991; 29:352-354.
 645. Prescott NM. On the benefits of tropical disease control. In: Health policies in developing countries Edited by Wood C, Rue Y. Royal Society of Medicine International Congress and Symposium Series. No. 24, 1980.
 646. Price EW. Mappa mundi and tropical medicine. Transactions of the Royal Society for Tropical Medicine and Hygiene 1989; 83:574.
 647. Rachou RG. Daily variation in microfilaraemia due to *Wuchereria bancrofti* in 27 persons for 28 consecutive days. Tropical Diseases Bulletin 1956; 53:470.
 648. Raghavan NGS. Congenital filariasis. Bulletin of the National Society of India for Malaria and Mosquito Communicable disease 1958; 6:147-154.
 649. Raghavan N, Freedman DO, Fitzgerald PC, Unnasch TR, Ottesen EA, Nutman TB. Cloning and characterization of a potentially protective chitinase-like recombinant antigen from *Wuchereria bancrofti*. Infection and Immunity 1994; 62:1901-1908.
 650. Rahmah N, Khairul Anuar A, Karim R, Mehdi R, Sinniah B, Wan Omar A. Potential use of IgG2 ELISA in the diagnosis of chronic elephantiasis and IgG4 ELISA in the follow-up of microfilaraemic patients infected with *Brugia malayi*. Biochemical & Biophysical Research Communications 1994; 205:202-207.
 651. Rahmah N, Khairul Anuar A, Tengu Ariff, Zurainee MN, Noor A'shikin A, Fadzillah A, Maimunah A, Ashrafu Haq J. Use of antifilarial IgG4 ELISA to detect *Brugia malayi* infection in an endemic area of Malaysia. Tropical Medicine and International Health 1998; 3:184-188.
 652. Rahmah N, Noor Ashikin A, Khairil Anuar A, Tengku Ariff RH, Abdullah B, Chan GT, Williams SA. PCR-ELISA for the detection of *Brugia malayi* infection using finger prick blood. Transactions of the Royal Society of Tropical Medicine and Hygiene 1998; 92:404-406.
 653. Rajasekariah GR, Parab PB, Sreedhara Swamy KH, Subrahmanyam D. Relevance of species-specific filarial antigens for diagnosis. Journal of Communicable diseases 1986; 18:223-233.
 654. Ramachandran CP. The control of lymphatic filariasis: despair to hope. Filarial Links 1997; 2:4-6. Published by the WHO Collaborating Centre for the Control of Lymphatic Filariasis, James Cook University, Townsville Australia.
 655. Ramaiah KD (a), Vijay, Kumar KN, Ramu K. Knowledge and beliefs about transmission, prevention and control of lymphatic filariasis in rural areas of South India. Tropical Medicine and International Health 1996; 1:433-438.
 656. Ramaiah KD (b), Ramu K, Kumar KN, Guyatt H. Epidemiology of acute filarial episodes caused by *Wuchereria bancrofti* infection in two rural villages in Tamil Nadu, South India. Transactions of the Royal Society of Tropical Medicine and Hygiene 1996; 90:639-643.
 657. Ramaiah KD, Kumar KN, Ramu K, Pani S, Das PK. Functional impairment caused by lymphatic filariasis in rural areas of south India. Tropical Medicine and International Health 1997; 2:832-838.
 658. Ramaiah KD, Ramu K, Guyatt, Vijar Kumar KN, Pani SP. Direct and indirect costs of the acute form of lymphatic filariasis to households in rural areas of Tamil Nadu, south India. Tropical Medicine and International Health 1998; 3:1008-115.
 659. Ramaiah KD, Vijay Kumar KN. Effect of lymphatic filariasis on school children. Acta Tropica 2000; 76:197-199.
 660. Ramakrishnan SP. National filaria control programme-a review. Indian Journal of Medical Research 1960; 14:457-494.
 661. Ramalingeswararao K; Venkatanarayana M, Dhanvantari Naidu Y, Narasimham MVVL, Viswanatham R, Rao CK. Transplacental transmission in Bancroftian filariasis. Indian Journal of Medical Research 1984; 79:495-496.
 662. Ramaprasad P, Prasad GB, Harinath BC. Microfilaraemia, filarial antibody, antigen & immune complex levels in human filariasis before, during & after DEC therapy: a two year follow up. Acta Tropica (Basel) 1988; 45:245-255.
 663. Ramaprasad P, Harinath BC. Use of fractionated urinary filarial antigen in the diagnosis of human filariasis. Journal of Tropical Medicine & Hygiene 1995; 98:35-40.
 664. Ramu K, Ramaiah KD, Guyatt H, Evans D. Impact of lymphatic filariasis on the productivity of male weavers

- in a south Indian village. Transactions of the Royal Society of Tropical Medicine & Hygiene 1996; 90:669-670.
665. Ramzy RMR, Helmy H, Faris R, Gad AM, Chandrashekar R, Weil GJ. Evaluation of a recombinant antigen-based antibody ELISA for the diagnosis of Bancroftian filariasis in Egypt. Annals of Tropical Medicine & Parasitology 1995; 89:443-446.
 666. Rao CK, Panduranga Rao P, Russel S, Hamzakoya KK, Chandrasekaran A, Roychoudhury SP, Krishna Rao C. Control of Bancroftian filariasis with common salt medicated with diethylcarbamazine in Lakshadweep. Indian Journal of Medical Research 1981; 73:865-873.
 667. Rao CK. Filariasis control in primary health care. Journal of Communicable Diseases 1982; 14:207-209.
 668. Rao SS, Maplestone PA. The adult of *Microfilaria malayi* Brug. Indian Medical Gazette 1940; 75:159-160.
 669. Rathaur S, Robertson BD, Selkirk ME, Maizels RM. Secretory acetylcholinesterases from *Brugia malayi* adult and microfilarial parasites. Molecular and Biochemical Parasitology 1987; 26:257-265.
 670. Rauyajin O, Kamthornwachara B, Yablo P. Socio-cultural and behavioural aspects of mosquito-borne lymphatic filariasis in Thailand: a qualitative analysis. Social Science and Medicine 1995; 41:1705-1713.
 671. Ravindran B, Sahoo PK, Dash AP. Lymphatic filariasis & malaria: concomitant parasitism in Orissa, India. Transactions of the Royal Society of Tropical Medicine & Hygiene 1998; 92:21-23.
 672. Ravindranathan TC, Sethumadhavan KPV, Roychowdhury SP, Babu CS, Kanoujia KH, Narasimham MVVL, Rao CK. Seasonal variation in microfilaraemia (Bancrofti) density & filarial disease. Journal of Communicable Diseases 1980; 12:73-77.
 673. Ray D, Abel R, Selvarai KG. Epidemiology of pulmonary eosinophilia in rural south India - a prospective study 1981-86. Journal of Epidemiology and Community Health 1993; 47:469-474.
 674. Reddy MVR, Malhotra A, Harinath BC. Detection of circulating antigen in Bancroftian filariasis by sandwich ELISA using filarial serum IgG. Journal of Helminthology 1984; 58:259-262.
 675. Regunathan J, Jayaraman K, Kaliraj P. Cellular immune response studies in Bancroftian filariasis. Journal of Helminthology 1997; 71:265-267.
 676. Remme JHF, Raadt P, Godal T. The burden of tropical disease. Medical Journal of Australia 1993; 158:465.
 677. Reynouard F, Barrabes A, Lacroix R, Combescot C. Effect of 17 beta-estradiol, progesterone and testosterone on *Dipetalonema vitae* parasitosis in the castrated female golden hamster *Cricetus auratus*. Annales de Parasitologie Humaine et Comparee 1984; 59:237-244.
 678. Rifkin H, Thompson KJ. Structural changes in early filariasis. Archives of Pathology 1945; 40:220-229.
 679. Riji HM. Comparison of knowledge on filariasis and epidemiological factors between infected and uninfected respondents in a Malay community. Southeast Asian Journal of Tropical Medicine & Public Health 1986; 17:457-463.
 680. Riley EM, MacLennan C, Kwiatkowski DK, Greenwood BM. Suppression on *in-vitro* lymphoproliferative responses can be partially reversed by indomethacin. Parasite Immunology 1989; 11:509-517.
 681. Roberts LS, Janovy J. Nematodes: Filarioidea, the filarial worms. In: Schmidt GD, Robers LS. Foundations of Parasitology 5th. Edition, chapter 29. WCB publishers Boston 1996.
 682. Rocha A, Addiss D, Ribeiro ME, Noroes J, Baliza M, Medeiros Z, Dreyer G. Evaluation of the Og4C3 ELISA in *Wuchereria bancrofti* infection: infected persons with undetectable or ultra-low microfilarial densities. Tropical Medicine & International Health 1996; 1:859-864.
 683. Rodrigues-Silva R, Moura H, Dreyer G, Rey L. Human pulmonary dirofilariasis: a review. Revista Instituto de Medicina Tropical de Sao Paulo 1995; 37:225-233.
 684. Rogers R, Denham DA. Studies with *Brugia pahangi*. Measurement of lymph flow in infected cats. Southeast Asian Journal of Tropical Medicine and Public Health 1975; 6:199-205.
 685. Rogers R, Denham DA. Studies with *Brugia malayi*: The activity of levamisole. Journal of Helminthology 1976; 50:21-28.
 686. Rome HP, Fogel RH. The psychosomatic manifestations of filariasis. Journal of the American Medical Association 1943; 123:944-947.
 687. Rose HM, Culbertson JT, Lipman MO. Antistreptolysin titres in cases of filariasis with recurrent lymphangitis among military personnel. Journal of Clinical Investigation 1945; 24:532-536.
 688. Rosenblatt P, Beaver PC, Orihel TC. A filarial infection apparently acquired in New York City. American Journal of Tropical Medicine & Hygiene 1962; 11:641-645.
 689. Routh HB, Bhowmik KR. History of elephantiasis. International Journal of Dermatology 1993; 32:913-916.
 690. Sabesan A, Krishnamoorthy K, Pani SP, Panicker KN. Man days lost due to repeated acute attacks of lymphatic filariasis. Trends in Life Sciences (India) 1992; 7:5-7.
 691. Sabry M. Critical analysis of an epidemetrans model for the assessment of Bancroftian filariasis endemicity in some areas in Egypt. Journal of Tropical Medicine 1992; 95:260-265.
 692. Sahai R, Narasimham MVVL, Datta A, Chandrasekharan A, Rao CK. Solar eclipse and microfilaraemia. Journal of Communicable Diseases 1980; 12:161-163.
 693. Selkirk ME, Maizels RM, Yazdanbakhsh M. Immunity and the prospects for vaccination against filariasis. Immunobiology 1992; 184:263-281.
 694. Sullivan TJ, Carnahan VA, Cutting RT. Filariasis in American Servicemen. Annual Progress Report of United States Army Medical Research Team and Institute Pasteur Vietnam September July 1968 - June 1969: 69-72.
 695. Sullivan TJ, Carnahan VA, Cutting RT. Filariasis without microfilaraemia: a case report. Military Medicine 1970; 135:797-798.
 696. Santhanam SH, Kumar KVP, Sethumadhavan A, Chandrasekharan DC, Jain A, Malhotra TK, Ghosh GJ, Weil GJ. Detection of *Wuchereria bancrofti* antigen in serum and finger prick blood samples by enzyme immunoassay: field evaluation. Tropical Medicine and Parasitology 1989; 40:440-444.
 697. Santiago-Stevenson D, Oliver-Gonzalez J, Hewitt RI. Treatment of *Filariasis bancrofti* with 1-diethylcarbamyl-4-methylpiperazine hydrochloride (Hetrazan). Journal of the American Medical Association 1947; 135:708-712.
 698. Saphir W. Filariasis. Journal of the American Medical Association 1945; 128:1142-1145.
 699. Sapico FL, Baltazar RF. Filariasis presenting as a case of gross haematuria. Acta Medica Philippina 1967; 3:273-275.
 700. Sartono E (a), Kruize YCM, Partono F, Kurniawan A, Maizels RM, Yazdanbakhsh M. Specific T cell

- unresponsiveness in human filariasis: diversity in underlying mechanisms. *Parasite Immunology* 1995; 17:587-594.
701. Sartono E (b), Kruize YCM, Partono F, Kurniawan A, Maizels RM, Yazdanbakhsh M, Kurniawan A, van der Meide PH, Partono F, Maizels RM, Yazdanbakhsh M. Elevated cellular immune responses and interferon release after long-term diethylcarbamazine treatment of patients with human lymphatic filariasis. *Journal of Infectious Diseases* 1995; 171:1683-1687.
 702. Sartono E, Kruize YCM, Kurniawan A, Maizels RM, Yazdanbakhsh M. Depression of antigen-specific interleukin-5 and interferon responses in human filariasis as a function of clinical status and age. *Journal of Infectious Diseases* 1997; 175:1276-1280.
 703. Sasa S. Human filariasis: a global survey of epidemiology and control. University Park Press, Baltimore 1976. Sawada T, Takei K, Goll M.M. Immunodiagnosis studies on filariasis 111. Isolation and purification of antigen for intradermal skin test.
 704. *Japanese Journal of Experimental Medicine* 1965; 35:125-132.
 705. Sabena KC, Sabena RG, Tandon A, Bhutto B, Srivastava VK. Some observations on immunological status and immuno diagnosis in Bancroftian filariasis. *Biochemical Society Transactions* 1981;9:531-533.
 706. Scott MG, Briles DE, Hackelford PA, Smith DS, Nahm MH. Human antibodies to phosphocholine IgG anti-PC antibodies express restricted numbers of V and C regions. *Journal of Immunology* 1987; 138:3325-3337.
 707. Schacher JF, Sahyuni PF. A chronological study of the histopathology of filarial disease in cats and dogs by *Brugia pahangi*. *Transactions of the Royal Society of Tropical Medicine & Hygiene* 1967; 61:234-245.
 708. Schmidt LH, Esslinger JH. The course of infections with *Plasmodium falciparum* in owl monkeys displaying microfilaremia. *American Journal of Tropical Medicine and Hygiene* 1981; 30:5-11.
 709. Schofield FD, Rowley RE. The effect of prednisone on persistent microfilaremia during treatment with diethylcarbamazine. *American Journal of Tropical Medicine and Hygiene* 1961; 10:849-853.
 710. Schrater AF, Spielman A, Piessens WF. Predisposition to *Brugia malayi* microfilaremia in progeny of infected gerbils. *American Journal of Tropical Medicine & Hygiene* 1983;32:1306-1308.
 711. Schultz GW. A study of Bancroftian filariasis on the islands of Bataan and Rapu, Philippines. *Southeast Asian Journal of Tropical Medicine & Public Health* 1988; 19:207-214.
 712. Schuurkamp GJT, Matango M, Kereu R, Napil J. Malaria, splenomegaly and filariasis in the Ok Tedi area of the Star mountains Papua New Guinea: three years after residual DDT spray. *Papua New Guinea Medical Journal* 1987;30:39-44.
 713. Schuurkamp GJT, Kereu RK, Bulungol PK. Diethylcarbamazine in the control of Bancroftian filariasis in the highly endemic Ok Tedi area of Papua New Guinea: phase 1. *Papua New Guinea Medical Journal* 1990;33:89-98.
 714. Schuurkamp GJT. The epidemiology of malaria and filariasis in the Ok Tedi region of Western province, Papua New Guinea. PhD thesis: University of Papua New Guinea, 1992.
 715. Schuurkamp GJT, Kereu RK, Bulungol PK, Kawereng A, Popon W, Crane GG, Greenidge J, Spicer PE. Diethylcarbamazine in the control of splenomegaly associated with Bancroftian filariasis in the Ok Tedi area of Papua New Guinea. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1992; 86:531-536.
 716. Seurat LG. Orthogenese des filaires. *Bulletin de la Societe d'Histoire Naturelle de l'Afrique du Nord* 1921; 12:28-37.
 717. Shaoqing Z, Feng C, Webber R. A successful control programme for lymphatic filariasis in Hubei, China. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1994; 88:510-512.
 718. Shenoy RK, Kumaraswami V, Rajan K, Thankom S, Jalajakumari. A comparative study of the efficacy and tolerability of single and split doses of ivermectin and diethylcarbamazine in periodic Brugian filariasis. *Annals of Tropical Medicine and Parasitology* 1993; 87:459-467.
 719. Shenoy RK, Sandhya K, Suma TK, Kumaraswami V. A preliminary study of filariasis-related acute adenolymphangitis with special reference to precipitating factors and treatment modalities. *Southeast Asian Journal of Tropical Medicine and Public Health* 1995; 26:301-305.
 720. Shenoy RK(a), Suma TK, Rajan K, Kumaraswami V. Prevention of acute adenolymphangitis in brugian filariasis: comparison of the efficacy of ivermectin and diethylcarbamazine each combined with local treatment of the affected limb. *Annals of Tropical Medicine and Parasitology* 1998; 92:587-594.
 721. Shenoy RK (b) Varghese J, Kuttikkal VV. The efficacy, tolerability and safety of diethylcarbamazine-fortified salt in the treatment of the microfilaremiias of brugian filariasis: an open, hospital based study. *Annals of Tropical Medicine and Parasitology* 1998; 92:285-293.
 722. Shenoy RK, Dalia S, John A, Suma TK, Kumaraswami V. Treatment of the microfilaremia of asymptomatic brugian filariasis with single doses of ivermectin, diethylcarbamazine or albendazole, in various combinations. *Annals of Tropical Medicine and Parasitology* 1999; 93:643-651.
 723. Shenoy RK, Kumaraswami V, Suma TK, Rajan K, Radhakuttyamma G. A double blind, placebo-controlled study of the efficacy of oral penicillin, diethylcarbamazine or local treatment of the affected limb in preventing acute adenolymphangitis in lymphodema caused by brugian filariasis. *Annals of Tropical Medicine and Parasitology* 1999; 93:367-377.
 724. Shin Z, Sun D, Shi H. Cost benefit analysis Malayan filariasis control in Miaoxi township, Huzhou city, Zhejiang province during 1964-1987. *Chung Kuo Chi Sheng Chung Hsueh Yu Chi Sheng Chung Ping Tsa Chih* 1995; 13:193-196.
 725. Sibthorpe BN. On the adult male *Filaria sanguinis hominis*. *British Medical Journal* 1889; 1:1334-1335.
 726. Sidkey HA, Hassan ZA, Hassan RR, Gaafar SA, Awadallah FEM. Disseminated *Hymenolepis nana* in blood of a filarial patient. *Journal of the Egyptian Society of Parasitology* 1987; 17:155-159.
 727. Simon F, Prieto G, Muro A, Cancrini G, Cordero M, Genchi C. Human humoral response to *Dirofilaria* species. *Parassitologia* 1997; 39:397-400.
 728. Simonsen PE, Lemnge MM, Msangeni HA, Jakobsen, Bygbjerg IBC. Bancroftian filariasis: the patterns of filarial-specific immunoglobulin G1 (IgG1), IgG4 and circulating antigens in an endemic community of northeastern Tanzania. *American Journal of Tropical Medicine & Hygiene* 1996; 55:69-75.
 729. Simonsen PE (a), Niemann L, Meyrowitsch DW. *Wuchereria bancrofti* in Tanzania: microfilarial periodicity and the effect of blood sampling time on microfilarial densities. *Tropical Medicine and International Health* 1997; 2:153-158.

730. Simonsen PE (b), Pedersen EM, Lemnge MM, Michael E. Lymphatic Filariasis research and control in Africa. *Parasitology Today* 1997; 13:413-415.
731. Simonsen PE, Dunyo SK. Comparative evaluation of three new tools for diagnosis of bancroftian filariasis based on detection of specific antigens. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1999; 93:278-282.
732. Simpson CF, Jackson RF, Bradley RE. Alterations of glomerular capillaries in canine heart worm disease. In: Bradley RE & Pacheco G. *Canine Heart worm Disease: the Current Knowledge*. University of Florida Press, Gainesville 1972.
733. Singh SN, Bhatnagar S, Fatma N, Chauhan PMS, Chatterjee RK. Antifilarial activity of a synthetic marine alkaloid, aplysinopsin (CDRI compound 92/138). *Tropical Medicine and International Health* 1997; 2:535-543.
734. Sivam NS, Jayaanthi S, Ananthakrishnan N, Elango A, Yuvaraj J, Hoti SL, Pani SP. Tropical vaginal hydroceles: are they all filarial in origin? *Southeast Asian Journal of Tropical Medicine and Hygiene* 1995; 26:739-742.
735. Sivan VM, Kaleysa Raj R. Quinone analogue irreversibly paralyses the filarial parasites in vitro. *Biochemical and Biophysical Research Communications* 1999; 256:81-83.
736. Sharma S, Misra S, Rathaur S. Secretory acetylcholinesterase of *Setaria cervi* microfilariae and its antigenic cross-reactivity with *Wuchereria bancrofti*. *Tropical Medicine and International Health* 1998; 3:46-51.
737. Smith D.H, Wilson T, Berezancev J.A, Lykov V, Pang M, Chari M.V, Davis A. Evaluation of the *Dirofilaria immitis* filarial skin test antigen in the diagnosis of filariasis. *Bulletin of the World Health Organisation* 1971; 44:771-782.
738. Smith FR. Filariasis: a study of 737 cases so diagnosed. *Naval Medical Bulletin* 1945; 44:719-726.
739. Southgate BA. Studies on filariasis in the Pacific: a field trial of a counting chamber technique for the determination of microfilarial rates and densities. *Transactions of the Royal Society of Tropical Medicine & Hygiene* 1973; 65:521-524.
740. Southgate BA, Hamilton PJS. Problems of clinical and biological measurements in the epidemiology and control of filarial infections. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1974; 68:177-185.
741. Southgate BA. Recent advances in the epidemiology and control of filarial infection including entomological aspects of transmission. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1984; 78 (supplement):19-28.
742. Southgate BA. Intensity and efficiency of transmission and the development of microfilaraemia and disease: their relationship in lymphatic filariasis. *Journal of Tropical Medicine and Hygiene* 1992; 95:1-12.
743. Southgate BA, Bryan JH. Factors affecting transmission of *Wuchereria bancrofti* by Anopheline mosquitoes 4. Facilitation and its epidemiological significance. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1992; 86:120-127.
744. Sri Kumari Srisaillapathy CR, Ramesh A, Ganesan J. Association of ABO and Rh(D) blood groups with filariasis. *Human Hereditary* 1990; 40:381-385.
745. Sriwastava RG, Gupta PC, Sharma RS, Vajpayee GN. Haematuria in filariasis. *Journal of the Indian Medical Association* 1979; 62:19.
746. Srivastava SK, Sabena RG, Bhutto B, Tandon A, Srivastava BC, Sabena KC. Antibody response to TAB vaccine human filariasis - a preliminary communication. *Indian Journal of Medical Research* 1981; 74:524-526.
747. Srivastava SK, Agarwal A, Chauhan PM, Agarwal SK, Bhaduri AP, Singh SN, Fatima N, Chatterjee RK. Potent 1,3-substituted-9H-pyridol [3,4-b] indoles as new lead compounds in antifilarial chemotherapy. *Bioorganic Medicinal Chemistry* 1999; 7:1223-1236.
748. Srividya A, Pani SP. Filariasis and blood groups. *National Medical Journal of India* 1993; 6:207-209.
749. Srividya A, Das PK, Subramanian S, Ramaiah KD, Grenfell BT, Michael E, Bundy DAP. Past exposure and the dynamics of infection in young children. *Epidemiology and Infection* 1996; 195-201.
750. Steel C, Guinea A, McCarthy J, Ottesen EA. Long term effects of prenatal exposure to maternal microfilaraemia on immune responsiveness to filarial antigens. *Lancet* 1994; 343:890-893.
751. Steel C, Guinea A, Ottesen EA. Evidence for protective immunity to Bancroftian filariasis in the Cook Islands. *Journal of Infective Diseases* 1996; 174:598-605.
752. Stingl P, Pierce PF, Connor DH, Gibson DW, Straessle T, Ross MA, Ribas JL. Does dexamethasone suppress the Mazzotti reaction in patients with onchocerciasis? *Acta Tropica* 1988; 45:77-85.
753. Story DM, Mukhtar AS. Vaccination of Jirds, *Meriones unguiculatus*, against *Litomosoides carinii* and *Brugia malayi* using irradiated larvae or *L. carinii*. *Tropical Medicine and Parasitology* 1982;33:23-24.
754. Subbu VSV, Biswas AR. Ecobolic action of diethylcarbamazine. *Indian Journal of Medical Research* 1971; 59:646-647.
755. Sucharit S, Vutikes S. Comparison of counting chamber, Sasa and standard smear methods of counting *Brugia pahangi* microfilariae from cat blood. *Southeast Asian Journal of Tropical Medicine & Public Health* 1975; 6:437.
756. Sucharit S, Harinasota C, Choochote W. Experimental transmission of sub-periodic *Wuchereria bancrofti* to the leaf monkey *Presbytis melalophos* and its periodicity. *American Journal of Tropical Medicine & Hygiene* 1982; 31:423-429.
757. Sucharit S, Rongsriyam Y. Transplacental transmission of microfilaria of *Brugia pahangi* in white rat. *Southeast Asian Journal of Tropical Medicine & Public Health* 1980; 11:416.
758. Suma TK, Shenoy RK, Varghese J, Kuttikal VV, Kumaraswami V. Estimation of ASO titre as an indicator of streptococcal infection precipitating acute adenolymphangitis in Brugian lymphatic filariasis. *Southeast Asian Journal of Tropical Medicine and Public Health* 1997; 28:827-829.
759. Sundaray NK, Das AK, Sanchette PC, Anand AC, Singh W. Acute filarial myositis. *Journal of the Association of Physicians of India* 1992; 40:693-694.
760. Sundberg RJ, Biswas S, Kumar Murthi K, Rowe D, McCall JW, Dzimiński MT. Bis-cationic hetero aromatics as macrofilaricides: synthesis of bis-amidine and bis-guanyldihydrazone derivatives of substituted imidazo[1,2-a]pyridines. *Journal of Medicinal Chemistry* 1998; 41:4317-4328.
761. Sunny PJ, Jain DC, Sorrian TP, Ghosh TK. Factors influencing acceptance and non-acceptance of DEC in a Bancroftian endemic area. *Journal of Communicable Diseases* 1986; 18:317-319.
762. Suresh NP, Thilagavathy HP, Kaleysa Raj R. A

- preliminary report on the relationship between ABO blood group and DC levels in filariasis. Proceedings of All India Symposium on Vectors and Vector-borne Diseases, Trivandrum, India 1982.
763. Suresh S, Kumaraswami V, Rajesh K, Suguna G, Vijayasekaran V, Ruckmani A, Rajamanickam MG. Ultrasonographic diagnosis of subclinical filariasis. *Journal of Ultrasound in Medicine* 1997; 16:45-49.
 764. Suswillo RR, Doenhoff MJ, Denham DA. Successful development of *Brugia pahangi* in T-cell deprived CBA mice. *Acta Tropica* 1981; 38:305-308.
 765. Swartzwelder JC. Filariasis bancrofti. In: Preventative Medicine in World War 2. Volume 7 - Communicable Diseases: Arthropod-borne diseases other than malaria. Edited by Coates JB, Hoff EC, Hoff PM. Published by Medical Department, United States Army, 1963>
 766. Tada I, Otsuji Y, Harada R, Mimori T, Fukumoto H. Skin test study of Bancroftian filariasis in Kuroshima island, Okinawa: a 13 year longitudinal study during a control campaign. *American Journal of Tropical Medicine & Hygiene* 1982; 31:262-267.
 767. Tajima K, Fujita K, Tsukidate S, Oda T, Tominga S, Suchi T, Hinuma Y. Sero epidemiological studies on the effects of filarial parasites on infestation of adult T-cell leukaemia virus in the Goto Islands, Japan. *Gann Japanese Journal of Cancer Research* 1983; 74:188-191.
 768. Tandon A, Saxena RP, Bhatia B, Saxena KC. Rodent and bovine filarial antigens in detection of antibodies in human Bancroftian filariasis by ELISA. *Indian Journal of Medical Research* 1981 73(suppl):93-96.
 769. Taylor AER, Denham DA. Immunology and diagnosis of filariasis. *Tropical Diseases Bulletin* 1985; 83:R1-R20
 770. Taylor MJ, Cross HF, Mohammed AA, Trees AJ, Bianco AE. Susceptibility of *Brugia malayi* and *Oncocerca lienalis* microfilariae to nitric oxide and hydrogen peroxide in cell free culture and from IFN gamma-activated macrophages. *Parasitology* 1996; 112:315-322.
 771. Taylor MJ, Turner PF. Control of lymphatic filariasis. *Parasitology Today* 1997; 13:85-86.
 772. Taylor MJ, Hoerauf A. *Wolbachia* bacteria of filarial nematodes. *Parasitology Today*; 1999; 15:437-442.
 773. Taylor MJ, Cross HF, Bilo K. Inflammatory responses induced by the filarial nematode *Brugia malayi* are mediated by lipopolysaccharide-like activity from Endosymbiotic *Wolbachia* bacteria. *Journal of Experimental Medicine* 2000; 191:1429-1435.
 774. Terhell AJ, Haarbrink M, Abadi K, Bronneberg DCH, Tieleman MC, Asri M, Yazdanbakhsh M. A filter paper technique for the detection of anti-filarial IgG4 in lymphatic filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1996; 90:196-198.
 775. Tesh RB, Guzman H. Mortality and infertility in adult mosquitoes after the ingestion of blood containing ivermectin. *American Journal of Tropical Medicine and Hygiene* 1990; 43:229-233.
 776. Tha Mya. Cerebral symptoms associated with filariasis. *Indian Medical Gazette* 1928; 63:636-637.
 777. Thompson KJ, Rifkin H, Zarrow M. Early filariasis in young soldiers: clinical and pathological analysis. *Journal of the American Medical Association* 1945; 129:1074:1079.
 778. Thorpe VG. *Filaria sanguinis hominis* in the South Sea Islands, with photomicrographs from Tonga & the friendly Islands. *British Medical Journal* 1896; ii:922-924.
 779. Tobie JE, Beye HK. Human infection with a subcutaneous filarial worm. *American Journal of Tropical Medicine and Hygiene* 1962; 11:64-67.
 780. Townson S, Tagboto SK, Castro J, Lujan A, Awadzi K, Titanji VPK. Comparison of the sensitivity of different geographical races of *Oncocerca volvulus* microfilariae to ivermectin: studies *in-vitro*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1994; 88:101-106.
 781. Trent SC (a). Re-evaluation of World War 2 veterans with filariasis acquired in the South Pacific. *American Journal of Tropical Medicine and Hygiene* 1963; 12:877-887.
 782. Trent SC (b). Treatment of filariasis without microfilaraemia with Hetrazan and Triancinolone. *Military Medicine* 1963; 128:1104-1107.
 783. Triteraprapab S, Jongwutiwes S, Eisinger SW, Scott AL. Bancroftian filariasis: an emerging disease of Thailand. *American Journal of Tropical Medicine and Hygiene* 1997; 57 (supplement):288 (abstract).
 784. Turner LH, Edeson JF. Studies on filariasis in Malaya; the periodicity of the microfilaria of *Wuchereria malayi*. *Annals of Tropical Medicine & Parasitology* 1957; 51:271-277.
 785. Turner PF. Filariasis in the Western Province of Papua New Guinea. PhD thesis: Department of Public Health and Tropical Medicine, James Cook University of North Queensland, Australia, 1993.
 786. Turner PF, Copeman D, Gerisi D, Speare R. A comparison of the Og4C3 antigen capture ELISA, the Knott test, and IgG4 test and clinical signs in the diagnosis of Bancroftian filariasis. *Tropical Medicine and Parasitology* 1993; 44:45-48.
 787. Turner PF, Usurup JP. A possible case of lymphatic filariasis in a white miner in Papua New Guinea (letter). *Medical Journal of Australia* 1997; 166:223.
 788. Vanamail P, Ramaiah KD, Pani SP, Das PK, Grenfell BT, Bundy DA. Estimation of the fecund life span of *Wuchereria bancrofti* in an endemic area. *Transaction of the Royal Society of Tropical Medicine and Hygiene* 1996; 90:119-121.
 789. Van Bogaert L, Dubois A, Janssens PG. Encephalitis in *Loa loa* filariasis. *Journal of Neurology, Neurosurgery and Psychiatry* 1955; 18:103-119.
 790. Van der Sar A, Hartz PH. The syndrome, tropical eosinophilia and microfilaria. *American Journal of Tropical Medicine* 1945; 25:83-87.
 791. Van der Winkel JGJ, Capel PJA. Human IgG Fc receptor heterogeneity: molecular aspects and clinical implications. *Immunology Today* 1993; 14:215-221.
 792. Van Hoegaerden M, Ivanoff B. A rapid, simple method for isolation of viable microfilariae. *American Journal of Tropical Medicine & Hygiene* 1986; 35:148-151.
 793. Varma Rajendra S, Sarin S, Chatterjee RK. Synthesis of beta-amino substituted 4-amino-3-nitropropionophenones as diethylcarbamazine analogs and their antifilarial activity. *Indian Journal of Chemistry Section B - Organic Chemistry Including Medicinal Chemistry* 1992; 31:711-713
 794. Ventura ER. Factors affecting participation in blood examination and treatment of filariasis in Sorsogon, Philippines. *Southeast Asian Journal of Tropical Medicine and Public Health* 1986; 17:464-469.
 795. Vickery AC, Albertine KH, Nayar JK, Kwa BH. Histopathology of *Brugia malayi* infected nude mice after immune-reconstitution. *Acta Tropica* 1991; 49:45-55.
 796. Vincent AL, Vickery AC, Lotz MJ, Desai U. The lymphatic pathology of *Brugia pahangi* in nude (athymic)

- and thymic mice C3H/HeN. *Journal of Pathology* 1984; 70:48-56.
797. Vincent AL, Rojas CAU, Ayoub EM, Ottesen EA, Harden EG. Filariasis and erisipela in Santo Domingo. *Journal of Parasitology* 1998; 84:557-561.
 798. Voller A, de Savigny D. Diagnostic serology of tropical parasitic diseases. *Journal of Immunological Methods* 1981; 46:1-29.
 799. Von Lichtenberg F. Inflammatory responses to filarial connective tissue parasites. *Parasitology* 1987; 94 (supplement) 5:101-122.
 800. Wada Y, Kimura E, Takagi M, Tsuda Y. Facilitation in *Anopheles* and spontaneous disappearance of filariasis: has the concept been verified with sufficient evidence? *Tropical Medicine and Parasitology* 1995; 46:27-30.
 801. Wadee AA, Vickery AC, Piessens WF. Characterization of immuno suppressive proteins of *Brugia malayi* microfilaria. *Acta Tropica* 1987; 44:343-352.
 802. Walker AS. Clinical Problems of War, chapter 11. Published by the Canberra War Memorial, Australia, 1952.
 803. Wamae CN, Roberts JM, Eberhard ML, Lammie PJ. Kinetics of circulating IgG4 after diethylcarbamazine and ivermectin treatment of Bancroftian filariasis. *Journal of Infectious Diseases* 1992; 165: 1158-1160.
 804. Wamae CN. Advances in the laboratory diagnosis of human lymphatic filariasis: a review. *East African Medical Journal* 1994; 71:171-181.
 805. Wamae CN, Lammie PJ, Muttunga JN. Bancroftian filariasis: profile of serum antifilarial antibody and circulating parasite antigen. *East African Medical Journal* 1995; 72:492-494.
 806. Wang LC. Evaluation of quantitative buffy coat analysis in the detection of canine *Dirofilaria immitis* infection: a model to determine its effectiveness in the diagnosis of human filariasis. *Parasitology Research* 1998; 84:246-268.
 807. Wang SH, Zheng HJ, Dissanayake S, Cheng WF, Tao ZH, Lin SZ, Piessens WF. Evaluation of recombinant chitinase and SXP1 antigens as anti microfilarial vaccines. *American Journal of Tropical Medicine and Hygiene* 1997; 56:474-481.
 808. Wartman WB. Lesions of the lymphatic system in early filariasis. *American Journal of Tropical Medicine* 1944; 24:299-313.
 809. Wartman WB, King BG. Early filariasis in American Soldiers. *Bulletin of the United States Army Medical Department* 1944; 75:45-51.
 810. Wartman WB. Filariasis in American armed forces in World War 2. *Medicine* 1947; 26:394.
 811. Waugh DA, Alexander JH, Ibels LS. Filarial chyluria associated glomerulonephritis and therapeutic considerations in the chyluric patient. *Australian & New Zealand Journal of Medicine* 1980; 10:559-562.
 812. Webber RH. The natural decline of *Wuchereria bancrofti* infection in a vector control situation in the Solomon Islands. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1977; 71:396-400.
 813. Webber RH. The maximum density of anopheline mosquitos that can be permitted in the absence of continuing transmission of filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1981; 75:499-505.
 814. Webber RH. Can anopheline-transmitted filariasis be eradicated? *Journal of Tropical Medicine and Hygiene* 1991; 94:241-244.
 815. Webster EA. Filariasis among white immigrants in Samoa. *United States Naval Medical Bulletin* 1946; 16:186-192.
 816. Wegesa P, McMahon JE, Abaru DE, Hamilton PJS, Marshall TF de C, Vauhan JP. Tanzania filariasis project: survey methodology and clinical manifestations of Bancroftian filariasis. *Acta Tropica* 1979; 36:369-377.
 817. Weil GJ, Powers KG, Parbuoni EL, Line BR, Furrow RD, Ottesen EA. *Dirofilaria immitis*. VI: Anti microfilarial immunity in experimental filariasis. *American Journal of Tropical Medicine & Hygiene* 1982; 31:477-485.
 818. Weil GJ, Hussain R, Kumaraswami V, Tripathy Sp, Phillips KS, Ottesen EA. Prenatal allergic sensitisation to helminth antigens in offspring of parasite-infected mothers. *Journal of Clinical Investigation* 1983; 75:1124-1129.
 819. Weil GJ, Lifits F. Identification and partial characterisation of a parasite antigen in sera from humans infected with *Wuchereria bancrofti*. *Journal of Immunology* 1987; 138:3035-3041.
 820. Weil GJ, Jain A, Santhanam SH, Malhotra TK, Kumar KVP, Sethumadhavan A, Lifits F, Ghosh GJ. A monoclonal antibody-based enzyme immuno assay for detecting parasite antigenaemia in Bancroftian filariasis. *Journal of Infectious Diseases* 1987; 156:350-355.
 821. Weil GJ, Sethumadhavan KVP, Santhanam S, Jain DC, Ghosh TK. Persistence of parasite antigenaemia following diethylcarbamazine therapy of Bancroftian filariasis. *American Journal of Tropical Medicine and Hygiene* 1988; 38:589-595.
 822. Weil GJ, Lammie PJ, Richards FO, Eberhard ML. Changes in circulating parasite antigen levels after treatment of Bancroftian filariasis with diethylcarbamazine. *Journal of Infectious Diseases* 1991; 164:814-816.
 823. Weil GJ, Ramzy RM, Chandrashekar R, Gad AM, Lourie RC jnr, Faris R. Parasite antigenaemia without microfilaraemia in Bancroftian filariasis. *American Journal of Tropical Medicine and Hygiene* 1996; 55:333-337.
 824. Weil GJ, Lammie PJ, Weiss N. The ICT filariasis test: a rapid-format antigen test for diagnosis of bancroftian filariasis. *Parasitology Today* 1997; 13:40-404.
 825. Welch JS, Dobson C. Antibodies to *Dirofilaria immitis* in Caucasian and Aboriginal Australians diagnosed by immunofluorescence and passive arthus hypersensitivity. *American Journal of Tropical Medicine and Hygiene* 1974; 23:1037-1045.
 826. Weller PF. Cell-mediated immunity in experimental filariasis: lymphocyte reactivity to filarial stage-specific antigens and to B and T cell mitogens during acute and chronic infection. *Cellular Immunology* 1978; 37:369-382.
 827. Weller PF, Ottesen EA, Heck L, Tere T, Neva FA. Endemic filariasis on a Pacific island. 1. Clinical, epidemiologic, and parasitologic aspects. *American Journal of Tropical Medicine & Hygiene* 1982; 31:942-952.
 828. Wijers DJB. Bancroftian filariasis in Kenya 1: Prevalence survey among adult males in the Coast Province. *Annals of Tropical Medicine & Parasitology* 1977; 71:313-331.
 829. Wijers DJB. Bancroftian filariasis in Kenya V: mass treatment given by members of the local community. *Annals of Tropical Medicine and Parasitology* 1984; 78:383-394.
 830. Wijetunge HPA (a). Filariasis in a university campus in Ceylon. *Journal of Tropical Medicine and Hygiene* 1967; 70:25-28.

831. Wijetunge HPA (b). Clinical manifestations of early Bancroftian filariasis. A study of 212 cases of microfilaraemia. *Journal of Tropical Medicine and Hygiene* 1967; 70:91-94.
832. Wilson T, Edeson JF, Wharton RH, Reid JA, Turner LH, Laing AB. The occurrence of two forms of *Wuchereria malayi* in man. *Transaction of the Royal Society of Tropical Medicine & Hygiene*.
833. Willenbucher J, Hofle W, Lucius R. The filarial antigens Av/Ov33-3 shown striking similarities to the major pepsin inhibitor from *Ascaris suum*. *Molecular and Biochemical Parasitology* 1993; 57:349-352.
834. Winkler S, El Menyawi, Floris-Linnau K, Graninger W. Short report: total serum levels of the nitric oxide derivatives nitrite/nitrate during microfilarial clearance in human filarial disease. *American Journal of Tropical Medicine and Hygiene* 1998; 59:523-525.
835. Witte MH, Jamal S, Williams WH, Witte CL, Kumaraswami V, McNeil GC, Case TC, Panicker TMR. Lymphatic abnormalities in human filariasis as depicted by lymphoangioscintigraphy. *Archives of Internal Medicine* 1993; 153:737-744.
836. Wong MM, Guest MF. Filarial antibodies and eosinophilia in human subjects in an endemic area. *Transactions of the Royal Society of Tropical Medicine & Hygiene* 1969; 63:797-801.
837. Woolhouse MEJ, Dye C, Etard JF, Smith T, Charlwood JD, Garnett GP, Hagan P, Hii JLK, Ndhlovu PD, Quinnel RJ, Watts CH, Chandiwana SK, Anderson RM. Heterogeneities in the transmission of infectious agents: implications for the design of control programs. *Proceedings of the United States Academy of Science* 1997; 94:338-342.
838. World Health Organisation Scientific Working Group: the immunology of filariasis. *Bulletin of the World Health Organisation* 1981; 59:1-8.
839. World Health Organisation. Lymphatic Filariasis. Fourth report of the WHO expert committee on filariasis. *World Health Organisation Technical Report Series 702*, WHO, Geneva 1984.
840. World Health Organisation. Control of lymphatic Filariasis: a manual for health personnel. *World Health Organisation*, Geneva 1987.
841. World Health Organisation. Lymphatic Filariasis: the disease and its control. Fifth report of the WHO expert committee on filariasis. *World Health Organisation*, Geneva 1992.
842. World Health Organisation expert committee on filariasis. Lymphatic filariasis: diagnosis and pathogenesis. *Bulletin of the World Health Organisation* 1993; 71:135-141.
843. World Health Organisation. Lymphatic filariasis: infection and disease: control strategies. Report of a consultative meeting held at the Universiti Sains Malaysia, Penang. *World health Organisation*, Geneva 1994.
844. World Health Organisation. Gender and tropical diseases. *World health Organisation*, Geneva 1995.
845. World Health Organisation. Report of a WHO informal consultation on epidemiologic approaches to lymphatic filariasis elimination: initial assessment, monitoring and certification. *World health Organisation*, Geneva 1998.
846. World Health Organisation. Building partnerships for lymphatic filariasis - strategic plan. *World health Organisation*, Geneva 1999.
847. Wucherer OE. Noticia preliminar sobre vermes de una especie ainda nao descrita, encontrados na uria de doentes de hematuria intertropical no Brazil. *Gazeta Medica da Bahia* 1868; 3:97-99.
848. Yan Y, Inuo G, Akao N, Tsukidate S, Fujita K. Down-regulation of murine susceptibility to cerebral malaria by inoculation with third-stage larvae of the nematode *Brugia pahangi*. *Parasitology* 1997; 114:333-338.
849. Yap HK, Woo KT, Yeo PPB, Chiang GSC, Singh M, Lim CH. The nephrotic syndrome associated with filariasis. *Annals of the Academy of Medicine of Singapore* 1982; 11:60-63.
850. Yates JA, Higahi GI. *Brugia malayi*: vaccination of jirds with ⁶⁰Cobalt-attenuated infective stage larvae protects against homologous challenge. *American Journal of Tropical Medicine & Hygiene* 1985; 34:1132-1137.
851. Yazdanbakhsh M (a), Paxton WA, Kruize CM, Sartono E, Kurniawan A, van het Wout A, Selkirk ME, Partono F, Maizels RM. T cell responsiveness correlates differentially with antibody isotypes levels in clinical and asymptomatic filariasis. *The Journal of Infectious Diseases* 1993; 167:925-931.
852. Yazdanbakhsh M (b), Sartono E, Kruize YCM, Kurniawan A, van der Pouw-Kraan T, van der Meide PH, Selkirk ME, Partono F, Hintzen RQ, van Lier RAW, Maizels RM. Elevated levels of T cell activation antigen CD27 and increased interleukin-4 production in human lymphatic filariasis. *European Journal of Immunology* 1993; 23:3312-3317.
853. Yazdanbakhsh M (a), Paxton WA, Brandenburg A, Van Ree R, Lens M, Partono F, Maizels RM, Selkirk ME. Differential antibody isotype reactivity to specific antigens in human lymphatic filariasis: gp15/400 preferentially induces immunoglobulin E (IgE), IgG4, and IgG2. *Infection and Immunity* 1995; 63:3772-3779.
854. Yazdanbakhsh M (b), Sartono E, Kruize YCM, Kurniawan A, Partono F, Maizels RM, Schreuder GM, Schipper R, de Vries RR. HLA and elephantiasis in lymphatic filariasis. *Human Immunology* 1995; 44:58-61.
855. Yazdanbakhsh M, Abadi K, de Roo M, Van Wouwe L, Denham D, Medeiros F, Verduijn W, RM, Schreuder GM, Schipper R, Giphart MJ, de Vries RR. HLA and elephantiasis revisited. *European Journal of Immunogenetics* 1997; 24:439-442.
856. Younis TA, el-Sharkawy IM, Youssef FG. Prevalence of intestinal parasites in filariasis-endemic areas in Egypt. *Journal of the Egyptian Society of Parasitology* 1997; 27:291-297.
857. Youssef FG, Hassanein SH, Cummings CE. A modified staining method to detect *Wuchereria bancrofti* microfilariae in thick-smear preparations. *Annals of Tropical Medicine & Parasitology* 1995; 89:93-94.
858. Zang X, Yazdanbakhsh M, Jiang H, Kanost MR, Maizels RM. A novel serpin expressed by blood-borne microfilariae of the parasitic nematode *Brugia malayi* inhibits human neutrophil serine proteases. *Blood* 1999; 94:1418-1428.
859. Zeligs MA. Psychosomatic aspects of filariasis. *Journal of the American Medical Association* 1945; 128:1139-1142.
860. Zheng HJ, Tao Z, Reddy MVR, Harinath BC, Piessens WF. Parasite antigens in sera and urine of patients with Bancroftian and Brugian filariasis detected by sandwich ELISA with monoclonal antibodies. *American Journal of Tropical Medicine & Hygiene* 1987; 36:554-560.
861. Zheng HJ a, Piessens WF, Tao ZH, Cheng WF, Wang SH, Cheng SH, Yee YM, Luo LF, Chen XR, Gan GB. Efficacy of ivermectin for control of microfilaraemia recurring after treatment with diethylcarbamazine. I. Clinical and parasitological observations. *American Journal of Tropical Medicine & Hygiene* 1991; 45:168-174.

862. Zheng HJ b, Tao ZH, Cheng WF, Wang SH, Cheng SH, Yee YM, Luo LF, Chen XR, Gan GB, Piessens WF. Efficacy of ivermectin for control of microfilaraemia recurring after treatment with diethylcarbamazine. 2. Immunologic changes following treatment. American Journal of Tropical Medicine & Hygiene 1991; 45:175-181.
863. Zhong-Xing L, Zhong-Jun S. A field survey on the role of low-density microfilaraemia cases in the transmission of filariasis. Southeast Asian Journal of Tropical Medicine and Public Health 1989; 20:153-156.
864. Zimmer HR. Hindu Medicine. ARNO Press, New York 1979.
865. Zambou NF, Mbiapo TF, Lando G, Tchana KA, Gouado I. Effect of *Onchocerca volvulus* infection on plasma vitamin A concentration in school children in rural region of Cameroon. Sante 1999; 9:151-155.
866. Zucker-Franklin D. Eosinopenia and Eosinophilia In: In Hematology 3rd. Edition. Edited by Williams W.J, Beutler E, Erslev A.J, Lichtman M.A. Marshall-Hill, New York 1983. Chapter 92.

