

Short Report

Absence of ivermectin-associated excess deaths

N. D. E. Alexander¹, M. J. Bockarie², W. A. Kastens^{3,4}, J. W. Kazura⁴ and M. P. Alpers¹
¹Papua New Guinea Institute of Medical Research, P. O. Box 60, Goroka, Papua New Guinea; ²Papua New Guinea Institute of Medical Research, P. O. Box 378, Madang, Papua New Guinea; ³Papua New Guinea Institute of Medical Research, P. O. Box 400, Maprik, Papua New Guinea; ⁴Division of Geographic Medicine, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA

Keywords: filariasis, *Wuchereria bancrofti*, chemotherapy, ivermectin, diethylcarbamazine, mortality, Papua New Guinea

Ivermectin belongs to a group of semisynthetic compounds derived from the soil bacterium *Streptomyces avermitilis*. It was released in 1981, and by the end of the decade had been used in at least 77 countries for control of parasites in agricultural animals (CAMPBELL, 1985, 1988). After its registration for human onchocerciasis in 1987, the number of annual treatments reached 9.2 million in 1993 and continues to grow rapidly (WHO, 1995). Recent clinical trials have shown that ivermectin is also effective against lymphatic filariasis, indicating treatment of millions more people affected by that disease (OTTESEN & CAMPBELL, 1994; WHO, 1997). Ivermectin is also effective against ectoparasites such as head lice (DUNNE *et al.*, 1991) and scabies mites (SULLIVAN *et al.*, 1997), and causes mortality in mosquitoes which have bitten people treated with the drug (WILSON, 1993).

In a recent study, ivermectin was associated with excess deaths when used to treat scabies in a long-term care unit for elderly patients (BARKWELL & SHIELDS, 1997). Although not confirmed by subsequent reports (REINTJES & HOEK, 1997; COYNE & ADDISS, 1997), this finding is a cause for concern because of the very large number of people being treated with the drug.

We present here an analysis of deaths in a part of the East Sepik Province of Papua New Guinea where we maintain demographic surveillance as part of a randomized community-based trial of ivermectin plus diethylcarbamazine (DEC) versus DEC alone for lymphatic filariasis caused by *Wuchereria bancrofti* (see BOCKARIE *et al.*, 1998). Annual treatment started in 1994, and the Table lists the 67 people who died between their first treatment and the beginning of the 1997 annual round. The dose was 400 µg/kg, and those aged less than 5 years were not treated. The death rates were analysed following SMITH & MORROW (1991). The study was randomized by community rather than individual, so it was possible to change regimen as a result of changing residence, although no death occurred among such people.

There is no statistically significant difference in the death rates, with any tendency being for a reduction, not an increase, in those receiving ivermectin. The same is true if the analysis is restricted to those aged at least 60 years at first treatment (Table). The conclusions are similar if the duration of surveillance is ignored and the data analysed as a 2x2 table, as in the original report by BARKWELL & SHIELDS (1997). Had the underlying rate ratio been 3.0 as they estimated, then the statistical power of our study would have been more than 99% for

Table. Death rates following treatment with ivermectin plus diethylcarbamazine or diethylcarbamazine alone

Treatment ^a	No. of deaths/ person-years of surveillance ^b	Rate ratio ^c	P
All treated persons			
Ivermectin plus DEC	28/3259 (0.0086)	0.74 (0.45–1.20)	0.22
DEC alone	39/3344 (0.0117)		
Age ≥60 years			
Ivermectin plus DEC	6/132 (0.045)	0.54 (0.21–1.37)	0.20
DEC alone	17/202 (0.084)		

^aDEC=diethylcarbamazine.

^bDeath rate in parentheses.

^c95% confidence interval in parentheses.

the whole population, and more than 90% for those aged 60 years and over (KIRKWOOD, 1988). In other words, the failure to detect such a difference is unlikely to be due to inadequate sample size.

In conclusion, we found no evidence of an increased death rate associated with ivermectin treatment, even at a much higher dose than the 150–200 µg/kg used by BARKWELL & SHIELDS (1997). This study was randomized by community, not individual, so it was not strictly designed for individually based analyses such as this. Nevertheless, the randomization has some advantages over the retrospective case-control design of BARKWELL & SHIELDS (1997), and the results provide reassurance that the excess mortality associated with ivermectin which they reported may have resulted from pharmacological interactions or other circumstances not generally pertaining in mass control programmes for onchocerciasis or lymphatic filariasis.

References

- Barkwell, R. & Shields, S. (1997). Deaths associated with ivermectin treatment of scabies. *Lancet*, **349**, 1144–1145.
- Bockarie, M. J., Alexander, N. D. E., Hyun, P., Dimber, Z., Bockarie, F., Ibam, E., Alpers, M. P. & Kazura, J. W. (1998). Randomised community-based trial of annual single-dose diethylcarbamazine with or without ivermectin against *Wuchereria bancrofti* infection in human beings and mosquitoes. *Lancet*, **351**, 162–168.
- Campbell, W. C. (1985). Ivermectin: an update. *Parasitology Today*, **1**, 10–16.
- Campbell, W. C. (1988). Ivermectin and faecal degradation. *Parasitology Today*, **4**, 21–22.
- Coyne, P. E. & Addiss, D. G. (1997). Deaths associated with ivermectin treatment of scabies. *Lancet*, **350**, 215–216.
- Dunne, C. L., Malone, C. J. & Whitworth, J. A. G. (1991). A field study of the effects of ivermectin on ectoparasites of man. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **85**, 550–551.
- Kirkwood, B. R. (1988). *Essentials of Medical Statistics*. Oxford: Blackwell Scientific Publications.
- Ottesen, E. A. & Campbell, W. C. (1994). Ivermectin in human medicine. *Journal of Antimicrobial Chemotherapy*, **34**, 195–203.
- Reintjes, R. & Hoek, C. (1997). Deaths associated with ivermectin for scabies. *Lancet*, **350**, 215.
- Smith, P. G. & Morrow, R. H. (1991). *Methods for Field Trials of Interventions Against Tropical Diseases: a 'Toolbox'*. Oxford: Oxford University Press.
- Sullivan, J. R., Walt, G. & Barker, B. (1997). Successful use of ivermectin in the treatment of endemic scabies in a nursing home. *Australasian Journal of Dermatology*, **38**, 137–140.
- WHO (1995). *Onchocerciasis and its Control. Report of a World Health Organization Expert Committee on Onchocerciasis Control*. Geneva: World Health Organization, Technical Report Series, no. 852.
- WHO (1997). *Tropical Disease Research: progress 1995–96. Thirteenth programme report of the UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases*. Geneva: World Health Organization.
- Wilson, M. L. (1993). Avermectins in arthropod vector management—prospects and pitfalls. *Parasitology Today*, **9**, 83–87.

Received 2 February 1998; accepted for publication 19 February 1998