

Are coinfections of malaria and filariasis of any epidemiological significance?

Ephantus J. Muturi · Benjamin G. Jacob ·
Chang-Hyun Kim · Charles M. Mbogo ·
Robert J. Novak

Received: 2 October 2007 / Accepted: 24 October 2007 / Published online: 20 November 2007
© Springer-Verlag 2007

Abstract Africa accounts for about 33 and 90% of the world's burden of lymphatic filariasis (LF) and malaria, respectively. Despite tremendous progress in the approach to their diagnosis, epidemiology, and treatment, and global campaigns for their control and/or elimination, their global burden and economic costs have continued to rise. In most rural areas of the tropics, both diseases co-occur in the same human population and share common mosquito vectors. It is therefore conceived that control of the two diseases can be integrated using tools that have been proven effective recently or in the past. Before implementation of control programs in areas co-endemic for both diseases, it is deemed necessary to understand how the two diseases interact in the vector and human hosts. Here, we summarize available knowledge on coinfections of malaria and LF and provide an insight on how they can be managed.

Introduction

Malaria and lymphatic filariasis (LF) are the world's most important parasitic diseases transmitted by mosquitoes. They are significant causes of morbidity and mortality

wherever they occur, making them priorities for elimination and control programs (Nabarro and Tayler 1998; Ottesen 2000). Both diseases are co-endemic in many areas of the tropics and are transmitted by a number of common vectors (Muirhead-Thomson 1953; Burkot et al. 1990; Chadee et al. 2003). It is therefore common to find coinfections of malaria and LF in a single mosquito vector and in human populations living in these areas. Because interaction among parasites is known to alter disease severity (Buck et al. 1978) and that the nature of interaction is not always predictable, knowledge of interactions among parasites is important in designing effective disease management programs. This paper demonstrates the epidemiological significance of concomitant infections of malaria and LF and highlights the strategies of managing coinfections with the two diseases.

Global burden of malaria and lymphatic filariasis

With half a billion annual infections and a death toll of 1–3 million deaths per year, malaria remains the most important parasitic disease transmitted by mosquitoes (World Health Organization 2005a). The greatest illness and death from malaria results from *Plasmodium falciparum*, the predominant malaria parasite in Africa, and the remaining proportion is accounted for by *Plasmodium vivax*, *Plasmodium ovale*, and *Plasmodium malariae* (Adl et al. 2005). Like many other parasitic diseases, human mortality and morbidity resulting from malaria is mainly concentrated in sub-Saharan Africa, where 90% of global malaria burden occurs. Although numerous multinational research projects have attempted to decrease malaria incidence through improved health care, personal protection, and adult vector control since the 1950s (World Health Organization 1996), the direct and indirect costs of malaria estimated at \$800

E. J. Muturi (✉) · B. G. Jacob · C.-H. Kim · R. J. Novak
Department of Medicine, William C. Gorgas Center
for Geographic Medicine,
University of Alabama at Birmingham,
206C Bevell Biomedical Research Building,
845 19th Street South,
Birmingham, AL 35294, USA
e-mail: emuturi@uab.edu

C. M. Mbogo
Center for Geographic Medicine and Research Coast,
Kenya Medical Research Institute,
Kilifi, Kenya

million (US) per year in 1987, doubled in less than 10 years to an estimated \$1.5–2.0 billion (US) in 1995 (Metcalf 1998).

The escalating burden of malaria has not only slowed economic development in many parts of sub-Saharan Africa but also has posed a global concern because of increased travel and trade. Many areas that were once considered malaria-free have witnessed severe malaria epidemics within the past two decades, and the situation continues to worsen (Roberts et al. 2000). The spread of insecticide and drug resistance, human population growth and movement, socioeconomic conditions, changes in climate, land use, and adaptability of vector mosquitoes to the changing environment has led to serious setbacks in global malaria control (Roberts 1964; Robert et al. 1998; Ghebreyesus et al. 1999; Githeko et al. 2000; Patz et al. 2000; Keating et al. 2004).

To exacerbate the intolerable burden of malaria is the reality that some tropical regions endemic for malaria also have a high prevalence of lymphatic filariasis (LF); another mosquito-borne disease that has continued to degrade human life despite being considered eradicable (Anon 1993). Currently, at least 1 billion people (240 million in Africa) are at risk of infection with LF, more than 120 million people in Africa, Asia, West Pacific, and South America are already infected, 44 million of which are seriously disfigured (Michael and Bundy 1997; World Health Organization 2001). It is estimated that some 4.6 million cases of lymphedema and more than 10 million cases of hydrocele occur in Africa (World Health Organization 2004). The psychosocial consequences associated with LF are enormous, and the disease is ranked the second most important cause of disability after emotional disorders (World Health Organization 1995).

More than 90% of LF cases are caused by *Wuchereria bancrofti*, the only known cause of LF in Africa, with *Brugia malayi* and *Brugia timori* accounting for the remaining 10% (Otessen et al. 1997). LF is responsible for an annual loss of 1 billion US dollars in Africa (Haddix et al. 1999), a figure similar to that incurred in India (Ramaiah et al. 2000). The great morbidity associated with LF provided the impetus for a global elimination program sponsored by the WHO, nongovernment organizations, the drug manufacturers Merck and GlaxoSmithKline, and numerous government health departments (Colatrella 2003). The program is based upon mass community treatment with diethylcarbamazine citrate (DEC) and albendazole in areas where there is no coendemicity with onchocerciasis and ivermectin (Mectizan®) and albendazole in areas where onchocerciasis and LF coexist (World Health Organization 1999; World Health Organization 2001). The goal of the program is to eliminate the disease globally by the year 2020 (Otessen et al. 1997). The program is now active in 20

African countries, nine of which are at mass drug administration (MDA) stage, and 11 have either completed disease mapping and waiting to start MDA, or mapping is still ongoing (WHO 2004). The therapeutic coverage rate is, on average, more than 70% and has shown a continued improvement over the years. However, less than 10% of at risk population is covered by MDA, and the implementation of disease management and vector control components have been delayed due to financial constraints, lack of personnel, technical difficulties with tools for endemicity mapping, and the co-endemicity of loiasis with LF in many endemic countries (Malacela-Lazaro and Twum-Danso 2004; World Health Organization 2004; World Health Organization 2005b; Gyapong and Twum-Danso 2006).

Despite the challenges in the global campaign to eliminate LF, there has been significant progress. For instance, substantial information is available on the vectors of LF and knowledge concerning the site-specific disease transmission and prevalence of human infection (Appawu et al. 2001; Ngwira et al. 2002; Muturi et al. 2006a; Muturi et al. 2006b). Information on the impact of various drug regimens (Kazura 1993; Ramzy et al. 2002; Alleman et al. 2006) and vector control (Maxwell et al. 1990; Maxwell 1999) in disease management is also available.

Anopheles vectors of malaria and filariasis

The major *Anopheles* species that transmit malaria are also important vectors of Bancroftian filariasis. The actual *Anopheles* species involved in transmission of the two parasites varies among geographical regions. In much of sub-Saharan Africa, *Anopheles funestus* and members of the *Anopheles gambiae* complex, mainly, *A. gambiae*, *Anopheles arabiensis*, *Anopheles merus*, and *Anopheles melas* are involved in transmission of malaria and Bancroftian filariasis (McMahon et al. 1981; Trape and Rogier 1996; Muturi et al. 2006b). Other examples of known *Anopheles* vectors of malaria and filariasis include *Anopheles philipinensis* and *Anopheles barbirostris* in India, *Anopheles leucosphyrus*, *Anopheles barbirostris*, and *Anopheles donaldi* in Malaysia (Webber 1991; Chang et al. 1995), *Anopheles darlingi* and *Anopheles aquasalis* in South America (Giglioli 1948; Validum 1998), and *Anopheles farauti* and *Anopheles punctulatus* in Papua New Guinea (Burkot et al. 1990). A comprehensive list of the various *Anopheles* species involved in malaria and filariasis transmission in different regions of the world is shown in Table 1.

Concomitant malaria and filarial infections in mosquitoes

Although the role of *Anopheles* mosquitoes as vectors of both human malaria and LF has long been established, little

Table 1 *Anopheles* vectors involved in malaria and filariasis transmission in various parts of the world

Geographic area	<i>Anopheles</i> species
Africa	<i>A. gambiae</i> , <i>A. arabiensis</i> , <i>A. merus</i> , <i>A. merus</i> , <i>A. funestus</i> ,
India and Bangladesh	<i>A. philippinensis</i> , <i>A. barbirostris</i>
Malaysia	<i>A. leucosphyrus</i> , <i>A. barbirostris</i> , <i>A. donaldi</i> , <i>A. letifer</i> , <i>A. whartoni</i> , <i>A. maculatus</i> , <i>A. campestris</i>
Philippines	<i>A. minimus</i>
Indonesia	<i>A. balabacensis</i>
Papua New Guinea	<i>A. farauti</i> , <i>A. punctulatus</i>
Solomons island	<i>A. koriensis</i>
South America	<i>A. darlingi</i> , <i>A. aquasalis</i>
China and Korea	<i>A. sinensis</i>

Sources: Burkot et al. 1990, Webber 1991, Chang et al. 1995, Validum 1998, Muturi et al. 2006a, and other sources

is known about the occurrence and prevalence of concomitant infections of the two diseases and how they interact within the vector. Earlier reports on occurrence of concomitant infections of the two parasites in the vector were coincidental and were observed during routine dissection of wild caught mosquitoes for either malaria or filarial infections. Muirhead-Thomson (1953) reported high infection rates of malaria and filarial parasites in *A. gambiae* s.l. in Dar-es-Salaam, Tanzania (then Tanganyika) and Accra, Ghana (then Gold Coast), but it was only on rare occasions that a single mosquito could carry malaria sporozoites and mature filaria larvae. Subsequent studies confirmed that concurrent transmission of malaria and filarial parasites by a single mosquito vector was rare in nature (Burkot et al. 1990; Muturi et al. 2006b).

Studies using *Aedes taeniorhynchus* indicated that mosquitoes infected with filarial nematode *B. malayi* were more susceptible to Rift Valley Fever virus because migration of microfilariae in mosquito midgut facilitated virus penetration into the hemocel (Turell 1984). Further studies revealed that mosquito midgut was a significant barrier to malaria infection (Ponnudurai 1988), suggesting that concurrent ingestion of microfilariae and malaria gametocytes could be mutually synergistic. Burkot and colleagues (1990) conducted a field study using wild populations of *A. punctulatus* and observed significantly higher numbers of *W. bancrofti* in *Plasmodium*-infected mosquitoes compared with uninfected mosquitoes. However, there was a greater mortality of mosquitoes carrying both infections when the worms reached the third stage making concurrent transmission of such infections rare. Along the Kenyan coast, *Wuchereria*-infected *A. gambiae* s.l. had significantly higher *P. falciparum* sporozoite rates than uninfected mos-

quitoes suggesting that filarial parasites may enhance malaria infections (Muturi et al. 2006b). Nonetheless, simultaneous occurrence of mature worms and malaria sporozoites was very rare. Alberquerque and Ham (1995) used *Aedes aegypti* to investigate the immune responses invoked by mosquitoes when simultaneously infected with *Plasmodium gallinaceum* and *Brugia pahangi* under laboratory conditions. They found that intrathoracic inoculation of secondary infection of *B. pahangi* reduced the development rate of preexisting infection of *P. gallinaceum* by 9.5–49% in susceptible strains and 50–90% in refractory strain. In both strains, individuals carrying double infections defended themselves against oocytes by causing a reduction in their size and also by melanization in the *B. pahangi* refractory strain. They concluded that activation of phenoloxidase cascade in response to microfilariae in the hemolymph could also be used against oocytes in the midgut. Recent studies in endemic areas of Africa have shown that despite the two diseases sharing common *Anopheles* vectors and environmental factors necessary for replication, one disease tends to dominate the other probably as a result of interspecies competition between microfilaria and *Plasmodium* within the vector or human host (Kelly-Hope et al. 2006; Muturi et al. 2006a). The exact mechanisms of this competition remain a conjecture in the scientific community.

Concomitant infections of malaria and filariasis in humans and other vertebrates

Co-occurrence of malaria and filarial parasites in a single human has been known for a long time (Muirhead-Thomson 1953). However, information on the occurrence, prevalence, and interaction of the two diseases in humans is scant. Three studies conducted in India demonstrated natural occurrence of coinfections of malaria and filariasis in humans but in prevalence lower than 1% (Prasad et al. 1990; Ghosh and Yadav 1995; Ravindran 1998). In Guyana, South America, 0.4 and 3.3% of individuals examined for filariasis and malaria parasites, respectively, were positive for both infections (Chadee et al. 2003). Muturi et al. (2006b) found 4.3% of the 94 persons examined in a coastal Kenyan village to carry both malaria and filariasis parasites. Three of the above studies found *P. falciparum* to co-occur more often with *W. bancrofti* than other *Plasmodium* species (Ghosh and Yadav 1995; Chadee et al. 2003; Muturi et al. 2006b), and one study found *P. vivax* and *W. bancrofti* to be the most common combination (Prasad et al. 1990). From these studies, it was evident that the most prevalent malaria parasite in an area was the one that co-occurred more often with *W. bancrofti*. Moreover, it was shown that coinfections of malaria and filariasis were least likely to occur when the prevalence of either of the two parasites was low (Muturi

et al. 2006b). The lowest threshold prevalence in which coinfections of the two diseases are unlikely to occur has not been established and remains a matter of guess. An epidemiological study along the Kenyan coast did not detect any coinfections of malaria and LF in a study site with 17.4% malaria prevalence and 2.8% LF prevalence (Muturi et al. 2006b). However, the sample size in this study was not large enough and, therefore, might have underestimated the true prevalence of LF and coinfections with the two diseases in the area. This is particularly so because microfilariae are not evenly distributed in the blood and, therefore, blood samples from microfilaremic humans might still yield negative results (Ngwira et al. 2002). Moreover, some studies have shown that mosquitoes are still able to pick filarial infections from ultralow microfilariae carriers (World Health Organization 1983; Lowrie et al. 1989). Because high worm load reduces mosquito survivorship (Klein 1986), mosquitoes that ingest low microfilariae densities are likely to survive longer, a factor that is likely to increase their chances of picking malaria parasites as well. Confirmatory work using animal models is warranted to establish to minimum parasite and prevalence threshold required to sustain the occurrence of coinfections of malaria and filariasis.

Interaction of malaria and filariasis parasites within a single host has been least studied in humans than it is in other vertebrate hosts. Both malaria and filarial infections are known to induce inflammatory cytokines (Richard et al. 1991; Das et al. 1996), and the two diseases have been reported to have antagonistic effect on immunoregulatory T lymphocytes. In *Aotus trivirgatus* monkeys, individuals concurrently infected with microfilariae and *P. falciparum* followed a more benign malaria course than amicrofilaremic individuals (Schmidt and Esslinger 1981). Inoculation of experimental mice with third stage larvae of filarial nematode *B. pahangi* activated Th2 response that conferred protective immunity to *Plasmodium bergeri* cerebral malaria (Yan et al. 1997). A more recent study has shown that mice coinfecting with filarial nematode *Litomosoides sigmodontis* and non-cerebral malaria clone of *Plasmodium chabaudi chabaudi* were more anemic and lost more body mass than mice infected with malaria only even after controlling for parasitemia (Graham et al. 2005). Their study demonstrated that in amicrofilaremic mice, filariasis interferes with delicate immunological balance in malaria infection and aggravates malaria-induced immunopathology. Previous studies had shown that helminth coinfection can impair mechanisms necessary to control malaria parasitemia and/or prevent immunopathological malaria (Mosmann and Sad 1996; Jankovic and Sher 2001; Maizels and Yazdanbakhsh 2003). However, there are also reports that *W. bancrofti* infection has no significant effect on *P. falciparum* parasitemia in humans (Muturi et al. 2006b). Collectively, these findings emphasize the need to understand the interactions

of malaria and filaria parasites in co-endemic areas before implementation of control or elimination programs (Kelly-Hope et al. 2006).

Strategies for reducing the prevalence of coinfections of malaria and filariasis

Control of coinfections of malaria and LF is easier than control of either of the two diseases. This is because transmission of such infections by a single vector is rare, their prevalence in humans is low, and they are more likely to occur when the prevalence of both diseases is high. Because of possible negative effects that might result from differential control of one disease (Kelly-Hope et al. 2006; Muturi et al. 2006a; Muturi et al. 2006b), integrated control of the two diseases is the most efficient way of managing coinfections of malaria and filariasis.

Vector control can help reduce vector densities and the risk of disease transmission. Vector control by indoor spraying of dichlorodiphenyltrichloroethane (DDT) in doses insufficient to interrupt malaria transmission led to successful elimination of filariasis in the Pacific island (Burkot et al. 2002), Solomon Islands (Wada 1974; Webber 1977; Webber 1979), and some parts of Papua New Guinea (Bockarie 1994). The disappearance of *Brugian* filariasis in 18 of the 21 foci previously recognized in Sri Lanka was attributed to the destruction of the aquatic plant *Pistia stratiotes* that is associated with the breeding of *Mansonia* vectors after a malaria vector control program (Abdulcader 1967). Source reduction was one of the methods used to achieve effective eradication of malaria in the United States, Italy, Israel (Kitron and Spielman 1989) and Zambian copper belt (Uttinger et al. 2002). The same method was used to eradicate *A. gambiae* in northeast Brazil, where it was well established and an important vector of malaria (Soper and Wilson 1943).

Currently, vector control in Africa targets adult populations mainly through use of insecticide-treated bed nets (ITNs). ITNs may reduce vector populations by killing them directly through contact with insecticides or by preventing them from feeding, through reduced house entry or early exiting (Mbogo et al. 1996; Bogh et al. 1998; Lengeler 2004; Lindblade et al. 2006). Untreated bed nets have been reported to substantially reduce malaria transmission and significantly reduce LF transmission (Burkot et al. 1990; Bockarie et al. 2002; Mwangi et al. 2003), but the impact is great when they are treated with insecticides (Bogh et al. 1998; Pedersen and Mukoko 2002). The greater impact of bed nets on LF than malaria control is because transmission of LF is very inefficient. There is no parasite multiplication in the vector, and only continuous exposure to infective mosquitoes can produce a patent infection of *W. bancrofti* (Southgate 1992). Nonetheless, it

is now becoming a reality that ITNs alone are insufficient to achieve large scale reduction in vector densities and transmission interruption, and thus, source reduction is being reappraised as an essential component of vector control after five decades of neglect (Fillinger et al. 2003; Fillinger et al. 2004; Killeen et al. 2004).

Chemotherapy is also used widely in the treatment of malaria and LF. In addition to reducing the mortality and morbidity associated with the two diseases, chemotherapy also reduces the number of infected people that can serve as infective agents to the mosquitoes. Interestingly, Ivermectin and DEC; two of the three drugs used in the Global Program for Elimination of Lymphatic Filariasis (World Health Organization 2001) have been reported to possess mosquitocidal activity. Under laboratory conditions, Pampiglione et al (1985) reported mortality of adult mosquitoes of *A. aegypti*, *Culex pipiens*, and *Anopheles stephensi* after ingestion of ivermectin in sugar solutions or after feeding on mice previously inoculated with the chemical. Tesh and Guzman (1990) reported death and reduced fecundity in *A. aegypti*, *Aedes albopictus*, and *Culex quinquefasciatus* mosquitoes fed blood–ivermectin mixture through a chick skin membrane. Cartel et al. (1991) demonstrated for the first time a significant reduction in mosquito survival rates after feeding on people treated with ivermectin and DEC up to 3 months previously. However, the lethal effect of ivermectin was too short-lived to impact significantly on wild-caught mosquitoes fed naturally on ivermectin-treated people following their normal life styles (Bockarie et al. 1999). Collectively, these findings suggest that timely DEC and ivermectin treatment of people at risk and their livestock on whom the vector populations depends on for blood meal may block epidemics of mosquito-borne diseases and may represent an additional tool in control of malaria, filariasis and coinfections of the two diseases. Surprisingly, despite the proven mosquitocidal effects of DEC and ivermectin, little is known about their specific effects on the major vectors of malaria and filariasis in the world.

Conclusion

The evidence provided here demonstrates the epidemiological significance of coinfections of malaria and LF. The literature shows that the two parasites might interact within the vector and human populations in a way that may alter the dynamics of transmission and the course of disease development in humans. To achieve successful results in the ongoing global campaigns for malaria and LF control, it is important to understand the interactions between malaria and LF parasites in areas where they co-occur. More importantly, it is evident that timely provision of treatment to

infected humans along with vector control in endemic communities can drastically reduce the prevalence of malaria, LF, and their coinfections.

Acknowledgements This study was funded by NIH/NIAID grant UO1054889 (to R.N.).

References

- Abdulcader MHM (1967) Present status of *Brugia Malayi* infection in Ceylon. *J Trop Med Hyg* 70:199–202
- Adl SM, Simpson A, Farmer M, Andersen R, Anderson O, Barta J, Bowser S, Brugerolle G, Fensome R, Fredericq S, James T, Karpov S, Kugrens P, Krug J, Lane C, Lewis L, Lodge J, Lynn D, Mann D, McCourt R, Mendoza L, Moestrup O, Mozley-Standridge S, Nerad T, Shearer C, Smirnov A, Spiegel F, Taylor M (2005) The new higher level classification of eukaryotes with emphasis on the taxonomy of protists. *J Eukaryot Microbiol* 52:399–351
- Albuquerque CM, Ham PJ (1995) Concomitant malaria (*Plasmodium gallinaceum*) and filaria (*Brugia pahangi*) infections in *Aedes aegypti*: Effect on parasite development. *Parasitol* 110:1–6
- Alleman MM, Twum-Danso N, Thylefors B (2006) The Mectizan donation program - highlights from 2005. *Filaria J* 5:11
- Anon (1993) Recommendations of the international task force for disease eradication. *Morbidity and Mortality Weekly Rep* 42:1–38
- Appawu AM, Dadzie SK, Baffoe-Wilmot A, Wilson DM (2001) Lymphatic filariasis in Ghana: entomological investigation of transmission dynamics and intensity in communities served by irrigation systems in the Upper East Region of Ghana. *Trop Med Int Health* 6:511–516
- Bockarie M (1994) Can lymphatic filariasis be eradicated in Papua New Guinea? *Papua New Guinea Med J* 37:61–64
- Bockarie M, Tavul L, Kastens W, Michael E, Kazura JW (2002) Impact of untreated bednets on prevalence of *Wuchereria bancrofti* transmitted by *Anopheles farauti* in Papua New Guinea. *Med Vet Entomol* 16:116–119
- Bockarie MJ, Hii JL, Alexander ND, Bockarie F, Dagoro H, Kazura JW, Alper MP (1999) Mass treatment with ivermectin for filariasis control in Papua New Guinea: Impact on mosquito survival. *Med Vet Entomol* 13:120–123
- Bogh C, Pedersen E, Mukoko D, Ouma J (1998) Permethrin-impregnated bednet effects on resting and feeding behavior of lymphatic filariasis vector mosquitoes in Kenya. *Med Vet Entomol* 12:52–59
- Buck A, Anderson R, MacRae A (1978) Epidemiology of poly-parasitism IV.. Combined effects on the state of health. *TropenMed Parasitol* 29:253–268
- Burkot T, Garner P, Paru R, Dagoro H, Barnes A, McDougall S, Wirtz R, Campbell G, Spark R (1990) Effect of untreated bednets on the transmission of *Plasmodium falciparum*, *Plasmodium vivax*, and *Wuchereria bancrofti* in Papua New Guinea. *Trans R Soc Trop Med Hyg* 84:773–779
- Burkot T, Molineaux L, Graves R, Paru R, Battistutta D, Dagoro H, Barnes A, Wirtz R, Garner P (1990) The prevalence of naturally acquired multiple infections of *Wuchereria bancrofti* and human malaria in anophelines. *Parasitol* 100:369–375
- Burkot T, Taleo G, Toeaso V, Ichimori K (2002) Initial progress towards and challenges to filariasis elimination in Pacific island communities. *Ann Trop Med Parasitol* 96:S61–69

- Cartel JL, Sechan Y, Spiegel A, Nguyen L, Barbazan P, Martin PMV, Roux JF (1991) Cumulative mortality rates in *Aedes polynesiensis* after feeding on Polynesian *Wuchereria bancrofti* carriers treated with single dose ivermectin, diethylcarbamazine and placebo. *Ann Trop Med Parasitol* 42:343–345
- Chadee DD, Rawlins C, Tiwari T (2003) Short communication: concomitant malaria and filariasis infections in Georgetown, Guyana. *Trop Med Int Health* 8:140–143
- Chang M, Dorasingham P, Hardin S, Nagum N (1995) Malaria and filariasis transmission in a village/forest setting in Baram District, Sarawak, Malaysia. *J Trop Med Hyg* 98:192–198
- Colatrella B (2003) The Mectizan® Donation Programme—successful collaboration between the public and private sectors. *Essential Drugs Monitor* 33:27–28
- Das BK, Sahoo PK, Ravindran B (1996) A role of tumor necrosis factor-alpha in acute lymphatic filariasis. *Parasite Immunol* 18:421–423
- Fillinger U, George S, Gerry FK, Bart GJK, Norbert B (2004) The practical importance of permanent and semipermanent habitats for controlling aquatic stages of *Anopheles gambiae sensu lato* mosquitoes: operational observations from a rural town in western Kenya. *Trop Med Int Health* 9:1274–1289
- Fillinger U, Knols BG, Becker N (2003) Efficacy and efficiency of new *Bacillus thuringiensis* var *israelensis* and *Bacillus sphaericus* formulations against Afrotropical anophelines in Western Kenya. *Trop Med Int Health* 8:37–47
- Ghebreyesus T, Haile M, Witten K, Getachew A, Yohannes A, Yohannes M, Teklehaimanot H, Lindsay S, Byass P (1999) Incidence of malaria among children living near dams in northern Ethiopia: community based incidence survey. *BMJ* 319:663–666
- Ghosh S, Yadav R (1995) Naturally acquired concomitant infections of bancroftian filariasis and human Plasmodia in Orissa. *Indian J Malariol* 32:32–36
- Giglioli G (1948) The transmission of *Wuchereria bancrofti* by *Anopheles darlingi* in the America tropics. *Am J Trop Med* 28:71–85
- Githeko A, Lindsay S, Confalonieri U, Patz J (2000) Climate change and vector-borne diseases: a regional analysis. *Bull World Health Organ* 78:1136–1147
- Graham A, Lamb TJ, Read A, Allen J (2005) Malaria–filaria coinfection in mice makes malarial disease more severe unless filaria infection achieves patency. *J Infect Dis* 191:410–421
- Gyapong J, Twum-Danso N (2006) Global elimination of lymphatic filariasis: fact or fantasy? *Trop Med Inter Health* 11:125–128
- Haddix A, Carter S, Michael E, Benton B (1999) Cost effectiveness of integrating lymphatic filariasis elimination into APOC onchocerciasis control programs. Report prepared for the Joint Action Forum/Joint Program Committee
- Jankovic D, Sher A (2001) Th1/Th2 effector choice in the immune system: a developmental program influenced by cytokine signals.. In: Segel LA, Cohen IR (eds) *Design principles for the immune system and other distributed autonomous systems*. Oxford University Press, New York
- Kazura J, Greenberg J, Perry R, Weil G, Day K, Alpers M (1993) Comparison of single dose diethylcarbamazine and ivermectin for treatment of bancroftian filariasis in Papua New Guinea. *Am J Trop Med Hyg* 49:804–811
- Keating J, Macintyre K, Mbogo C, Githure J, Beier J (2004) Characterization of potential larval habitats for *Anopheles* mosquitoes in relation to urban land-use in Malindi, Kenya. *Int J Health Geogr* 3:9
- Kelly-Hope L, Diggle P, Rowlingson B, Gyapong J, Kyelem D, Coleman M, Thomson M, Obsomer V, Lindsay S, Hemingway J, Molyneux D (2006) Short communication: Negative spatial association between lymphatic filariasis and malaria in West Africa. *Trop Med Int Health* 11:129–135
- Killeen GF, Seyoum ABK (2004) Rationalizing historical success of malaria control in Africa in terms of mosquito resource availability management. *Amer J Trop Med Hyg* 71:87–93
- Kitron U, Spielman A (1989) Suppression of transmission of malaria through source reduction: Anti-anopheline measures applied in Israel, the United States, and Italy. *Rev Infect Dis* 11:391–406
- Klein T, Harrison BA, Grove JS, Dixon SV, Andre RG (1986) Correlation of survival rates of *Anopheles dirus* A (Diptera: Culicidae) with different infection densities of *Plasmodium cynomolg*. *Bull World Health Organ* 64:901–907
- Lengeler C (2004) Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database Syst Rev* 2:CD000363
- Lindblade KA, Gimnig JE, Kamau L, Hawley WA, Odhiambo F, Olang G, ter Kuile FO, Vulule JM, Slutsker L (2006) Impact of sustained use of insecticide-treated bednets on malaria vector species distribution and culicine mosquitoes. *J Med Entomol* 43:428–432
- Lowrie RJ, Eberhard M, Lammie P, Raccurt C, Katz S, Duverseau Y (1989) Uptake and development of *Wuchereria bancrofti* in *Culex quinquefasciatus* that fed on Haitian carriers with different microfilaria densities. *Am J Trop Med Hyg* 41:429–435
- Maizels R, Yazdanbakhsh M (2003) Immune regulation by helminth parasites: cellular and molecular mechanisms. *Nat Rev Immunol* 3:733–744
- Malacela-Lazaro M, Twum-Danso N (2004) Lymphatic filariasis research forum, 2.4 program implementation. *Amer J Trop Med Hyg* 71:16–19
- Maxwell C, Curtis C, Haji H, Kisumku S, Thalib AI, Yahya SA (1990) Control of Bancroftian filariasis by integrating therapy with vector control using polystyrene beads in wet pit latrines. *Trans Roy Soc Trop Med Hyg* 84:709–714
- Maxwell CA, Mohammed K, Kisumku U, Curtis CF (1999) Can vector control play a useful supplementary role against Bancroftian filariasis? *Bull World Health Organ* 77:138–143
- Mbogo C, Baya N, Ofulla A, Githure J, Snow R (1996) The impact of permethrin-impregnated bednets on malaria vectors of the Kenyan coast. *Med Vet Entomol* 10:251–259
- McMahon J, Magayuka S, Kolstrup N, Mosha F, Bushford F, Abaru D (1981) Studies on the transmission and prevalence of Bancroftian filariasis in four coastal villages of Tanzania. *Ann Trop Med Parasitol* 75:415–431
- Metcalf RL (1998) Pesticides. In: Metcalf RL, Ziegler EN (eds) *In Encyclopedia of Environmental Science and Engineering*. Gordon and Breach Science Publishers
- Michael E, Bundy D (1997) Global mapping of lymphatic filariasis. *Parasitol Today* 13:472–477
- Mosmann T, Sad S (1996) The expanding universe of T-cell subsets: Th1, Th2 and more. *Immunol Today* 17:138–146
- Muirhead-Thomson R (1953) Inter-relations between filarial and malarial infections in *Anopheles gambiae*. *Nature* 172:352–353
- Muturi E, Mbogo C, Ng'ang'a Z, Kabiru E, Mwandawiro C, Novak R, Beier J (2006a) Relationship between malaria and filariasis transmission indices in an endemic area along the Kenyan Coast. *J Vector Borne Dis* 43:77–83
- Muturi JE, Mbogo C, Ng'ang'a Z, Kabiru E, Mwandawiro C, Beier J (2006b) Concomitant infections of *Plasmodium falciparum* and *Wuchereria bancrofti* on the Kenyan coast. *Filaria J* 5:8
- Mwangi T, Ross A, Marsh K, Snow RW (2003) The effects of untreated bednets on malaria infection and morbidity on the Kenyan coast. *Trans R Soc Trop Med Hyg* 97:369–372
- Nabarro DN, Tayler EM (1998) The 'Roll back malaria' campaign. *Science* 280:2067–2068
- Ngwira M, Jabu C, Kanyongoroka H, Mponda M, Crampin A, Branson K, Alexander D, Fine P (2002) Lymphatic filariasis in the Karonga district of northern Malawi: a prevalence survey. *Ann Trop Med Parasitol* 96:137–144

- Otessen E, Duke B, Karam MKB (1997) Strategies and tools for control/elimination of lymphatic filariasis. *Bull World Health Organ* 75:491–503
- Otessen EA (2000) The global programme to eliminate lymphatic filariasis. *Trop Med Int Health* 5:591–594
- Pampiglione S, Majori G, Petrangeli G, Romi R (1985) Avermectins, MK-933 and MK-936, for mosquito control. *Trans R Soc Trop Med Hyg* 79:797–799
- Patz J, Graczyk T, Geller N, Vittor AY (2000) Effects of environmental change on emerging parasitic diseases. *Int J Parasitol* 30:1395–1405
- Pedersen E, Mukoko D (2002) Impact of insecticide-treated materials on filaria transmission by the various species of vector mosquito in Africa. *Ann Trop Med Parasitol* 96:S91–S95
- Ponnudurai T, Billingsley PF, Rudin W (1988) Differential infectivity of *Plasmodium* for mosquitoes. *Parasitol Today* 4:319–321
- Prasad R, Virk K, Prasad H, Sharma V (1990) Concomitant occurrence of malaria and filariasis in man in India. *Mosquito-Borne Dis Bull* 7:51–53
- Ramaiah K, Das P, Michael E, Guyatt H (2000) The economic burden of lymphatic filariasis in India. *Parasit Today* 16:251–253
- Ramzy R, Setouhy M, Helmy H, Kandil A, Ahmed E, Farid H, Faris R, Weil G (2002) The impact of single-dose diethylcarbamazine treatment of bancroftian filariasis in a low-endemicity setting in Egypt. *Amer J Trop Med Hyg* 67:196–200
- Ravindran B, Sahoo PK, Dash AP (1998) Lymphatic filariasis and malaria: Concomitant parasitism in Orissa, India. *Trans Roy Soc Trop Med Hyg* 91:21–23
- Richard G, Barbara L, Antony S (1991) The involvement of TNF- α , IL-1, and IL-6 in immune response to protozoan parasites. *Parasitol Today* 12:13–16
- Robert V, Awono-Ambene H, Thioulouse J (1998) Ecology of larval mosquitoes with special reference to *Anopheles arabiensis* (Diptera: Culicidae) in market-garden wells in urban Dakar, Senegal. *J Med Entomol* 35:948–955
- Roberts DR, Manguin S, Mouchet J (2000) DDT house spraying and re-emerging malaria. *Lancet* 356:330–332
- Roberts M (1964) The control of epidemic malaria in the highlands of Western Kenya. Part II the campaign. *J Trop Med Hyg* 67:191–199
- Schmidt L, Esslinger J (1981) Courses of infections with *Plasmodium falciparum* in owl monkeys displaying a microfilaremia. *Am J Trop Med Hyg* 30:5–11
- Soper F, Wilson D (1943) *Anopheles gambiae* in Brazil. The Rockefeller Foundation, New York
- Southgate BA (1992) Intensity and efficiency of transmission and the development of microfilaremia and disease: Their relationship in lymphatic filariasis. *J Trop Med and Hyg* 95:1–12
- Tesh R, Guzman H (1990) Mortality and infertility in adult mosquitoes after the ingestion of blood containing ivermectin. *Am J Trop Med Hyg* 43:229–233
- Trape JF, Rogier C (1996) Combating malaria morbidity and mortality by reducing transmission. *Parasitol Today* 12:236–240
- Turell M, Rossignol PA, Spielman A, Rossi CA, Bailey CL (1984) Enhanced arboviral transmission by mosquitoes that concurrently ingested microfilariae. *Science* 225:1039–1041
- Utzinger J, Doumani F, Singer BH (2002) The economic payoffs of integrated malaria control in the Zambian copper-belt between 1930 and 1950. *Trop Med Int Health* 7:657–677
- Validum L (1998) Malaria returns: *Anopheles darlingi*, the scourge of generations is back. *Guyana Review* 5:8–16
- Wada Y (1974) In: Pal R, Wharton RH (eds) *Culex tritaeniorhynchus*. Control of arthropods of medical and veterinary importance. Plenum, New York
- Webber R (1977) The natural decline of *Wuchereria bancrofti* infection in a vector control situation in the Solomon Islands. *Trans R Soc Trop Med Hyg* 71:396–400
- Webber R (1979) Eradication of *Wuchereria bancrofti* infection through vector control. *Trans R Soc Trop Med Hyg* 73:722–724
- Webber R (1991) Can anopheline-transmitted filariasis be eradicated? *J Trop Med Hyg* 94:241–244
- World Health Organization (1983) Report of the steering committee of the scientific working group on filariasis. World Health Organization, Geneva
- World Health Organization (1995) World Health Report 1995: Bridging the gaps. WHO, Geneva
- World Health Organization (1996) Operational manual on the application of insecticides for control of the mosquito vectors of malaria and other diseases. WHO/CTD/VBC96
- World Health Organization (1999) Building partnerships for lymphatic filariasis-strategic plan. World Health Organization, Geneva
- World Health Organization (2001) The programme to eliminate lymphatic filariasis (PELF), Geneva.
- World Health Organization (2004) Lymphatic filariasis elimination in the African region: progress report. WHO, Congo Brazaville
- World Health Organization (2005a) World Malaria Report 2005. World Health Organization, WHO/UNICEF 2005., Geneva
- World Health Organization (2005b) Sixth meeting of the Technical Advisory Group on the Global Elimination of Lymphatic Filariasis, Geneva, Switzerland, 20–23 September 2005. *Weekly Epidemiol Rec* 80:401–408
- Yan Y, Inuo G, Akao N, Tsukidate S, Fujita K (1997) Down-regulation of murine susceptibility to cerebral malaria by inoculation with third-stage larvae of the filarial nematode *Brugia pahangi*. *Parasitol* 114:333–338