

Re-assessing the global prevalence and distribution of lymphatic filariasis

E. MICHAEL¹, D. A. P. BUNDY² and B. T. GRENFELL¹

¹ Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK

² Centre for the Epidemiology of Infectious Disease, University of Oxford, South Parks Road, Oxford OX1 3PS, UK

(Received 11 September 1995; revised 7 November 1995; accepted 7 November 1995)

SUMMARY

This paper estimates the global burden of lymphatic filariasis based on a review of the published literature on infection and disease surveys. A method for aggregating and projecting prevalence data from individual studies to national, regional and global levels, which also facilitates the estimation of gender and age-specific burdens, is presented. The method weights in favour of the larger, and hence presumably more reliable, studies and relies on estimated empirical relationships between gender, age, infection and disease in order to correct studies with incomplete data. The results presented here suggest that although the overall prevalence of filariasis cases is 2.0% globally (approximately totalling 119 million cases), the disease continues to be of considerable local importance, particularly in India and Sub-Saharan Africa. Estimates by age and gender clearly show that, unlike other helminth infections, filariasis is mainly a disease of the adult and older age-classes and appears to be more prevalent in males. This work suggests that the derivation of more accurate estimates of the burden of filariasis will require a better understanding of both the epidemiology and the spatial aspects of infection and disease. It also suggests that filariasis is preventable based on a geographically targeted strategy for control.

Key words: lymphatic filariasis, global burden, epidemiology.

INTRODUCTION

Reliable information about the distribution and prevalence of a parasitic disease is fundamental to the estimation of its public health significance (World Bank, 1993). For lymphatic filariasis, the distribution is thought to be confined to the tropics and sub-tropics while the most recent estimate of infection suggests the possibility of up to some 78.6 million individuals infected globally with the 3 major lymphatic filarial species – *Wuchereria bancrofti*, *Brugia malayi* and *B. timori* (WHO, 1992). These estimates are based largely on reports by Member States to the 1992 WHO Expert Committee on Filariasis, and have drawn attention to the fact that filariasis continues to be a sizeable problem in many parts of the world.

However, because these estimates do not differentiate between asymptomatic and clinical infection (WHO, 1984, 1992), they provide little information on the magnitude of morbidity due to the infection. Reliable estimates of disability have become increasingly important to informing the debate on the public health importance of helminth infections (Warren *et al.* 1993; Evans, Gelband & Vlassoff, 1993; World Bank, 1993; Chan *et al.* 1994). This is despite the fact that the chronic disease manifestations of filariasis, such as hydroceles in men and lymphoedema, in contrast to most other helmin-

thiasis (Guyatt & Bundy, 1991; Chan *et al.* 1994), are directly observable and is fairly well-documented in the literature.

These global estimates also do not provide data on the gender and age-distribution of infection (microfilaria-positive) and disease cases. Information on age and gender patterns are essential for attempts to quantify disease burden, especially those based on metrics, such as the Disability Adjusted Life Year (DALY), which measure the burden in terms of future stream of disability-free life lost, and used recently by the World Bank (1993) as a standardized approach to comparing the public health impact of different diseases and conditions.

Furthermore, the previous approaches to estimating infection (WHO, 1984, 1992) do not provide information on the technical bases of the reported estimates, which are likely to have varied between endemic countries. On the other hand, published literature on infection or disease surveys has been used successfully to estimate the burden of geo-helminth infections (Crompton, 1988, 1989; Bundy & Cooper, 1989; Chan *et al.* 1994). Development of appropriate methods for calculating national, regional and global infection and disease prevalences, however, is integral to exploiting this data resource for estimating the disease burden of lymphatic filariasis.

The major aim of this study is therefore to re-assess the current global and regional prevalence of lymphatic filariasis using published data of the

* Corresponding author.

Table 1. Sources of the estimates of prevalence (%) of lymphatic filariasis (infection plus disease) by country

Region ¹ and country	Prevalence ¹¹ (%)	Sources
(A) Bancroftian filariasis		
Sub-Saharan Africa		
Angola	8.89	Azevedo, J. F. de, 1964 ^d (423,234) ⁵
Benin	11.90	Brengues, Subra & Bouchite, 1969 ^{d,age,sex-specific; f,g} (2360)
Burkina Faso	12.88	Brengues <i>et al.</i> 1975 ^d (72,206) ¹²
Cameroon	11.11	Ripert <i>et al.</i> 1982 ^{d,sex-specific; f,g,males,adults} (626), Marceau <i>et al.</i> 1986 ^d (1000)
Cape Verde	0.99	Azevedo, J. F. de <i>et al.</i> 1969 ^d (607)
Central African Republic	1.29	Hawking, 1977 ^d (100)
Chad	2.84	Hawking, 1977 ^d (1681) ⁵
Comoros	26.73	Blanchy & Galtier 1982 ^{d,age,sex-specific} (353), Charafoudine & Pesson, 1986 ^{d,age,sex-specific} (419) ⁷ , Joire, Julivez & Galtier, 1988 ^{c,sex-specific,adults} (943)
Republic of Congo	0.70	Fain, Wery & Tilkin, 1969 ^{d,sex-specific} (720) ⁸
The Gambia	17.57	Knight, 1980 ^{d,f,g, < and ≥ 15 yrs.} (2,309) ¹⁵
Guinea	6.21	Hawking, 1977 ^d
Guinea Bissau	36.50	Azevedo, J. F. de 1964 ^{d,f} (1988) ^{3(sample size not given for one study)}
Ivory Coast	8.62	Brengues <i>et al.</i> 1975 ^a (607) ^d , (3555) ³
Kenya	7.03	Wijers & Kinyanjui, 1977 ^a (1794) ² , Wijers & Kaleli, 1984 ^d
Liberia	11.11	Brinkmann 1976 ^{d,age,sex-specific; f,g} (925), Kuhlow & Zielke, 1976 ^d (6,695) ⁶⁹ , Zielke & Chlebowsky, 1979 ^{d,age,sex-specific; g,age-specific} (1968) ¹⁷
Madagascar	12.50	Brygoo, 1958 ^d (468)
Malawi	17.35	Oram, 1958 ^{f,g,sex-specific} (1642) ³ , Oram, 1960 ^b
Mali	11.50	Brengues <i>et al.</i> 1975 ^d (4472) ⁶
Mauritius	13.95	Huehns, 1953 ^{d,age,sex-specific} (1930)
Mozambique	10.09	Pinhao, 1961 ^{d, < and ≥ 20 yrs} (1420) ⁴
Niger	4.90	Hawking, 1977 ^d
Nigeria	22.11	Wijayarathne <i>et al.</i> 1982 ^{d,age-specific} (1561), Udonsi, 1986 ^{d,age-specific} (1351) ⁴ , Udonsi, 1988 ^{b,d,age,sex-specific} (1418) ⁷ , Akogun, 1991 ^{d,age-specific; f,g} (2552) ⁹
Sao Tome and Principe	18.90	Azevedo J. F. de <i>et al.</i> 1969 ^d
Senegal	12.37	Jumier, Diallo & Diagne, 1971 ^{d,age-specific} (1841) ⁹
Tanzania	11.74	McMahon <i>et al.</i> 1981 ^a (2003) ⁴ , Matola, 1985 ^a (1947) ^{inf} , 2168 ^{dis}
Togo	5.42	Brengues <i>et al.</i> 1969 ^{d,age,sex-specific; f,g} (2360) ³
Zambia	0.76	Hawking, 1977 ^d (459)
Zimbabwe	4.63	Hawking, 1977 ^d (109)
Latin America and Caribbean		
Brazil	0.03	Franco & Lima, 1966 ^d (811,361) ²⁴ , Sherlock & Serafim, 1967 ^{d,age,sex-specific} (22,174) ³¹
Costa Rica	0.11	Weinstock <i>et al.</i> 1977 ^{d,age-specific; f,g} (1196), Madrigal, <i>et al.</i> 1979 ^d (119), Paniagua <i>et al.</i> 1983 ^{d,age,sex-specific} (2879) ⁴
Dominica Republic	1.55	Vincent, 1987 ^{d,sex-specific; e}
Guyana	7.33	Nathan & Stroom, 1990 ^{d,age,sex-specific} (2818)
Haiti	2.85	Raccurt, Mojon & Hodges, 1984 ^{d,age,sex-specific} (1428), Raccurt, 1986 ^d (6779), Raccurt <i>et al.</i> 1988 ^{d,age,sex-specific; f,g} (421)
Surinam	0.42	Oostbrug, 1985 ^{d,age,sex-specific; f,age,sex-specific} (107,415)
Trinidad	0.61	Nathan <i>et al.</i> 1982 ^{d,f,age,sex-specific; g,age-specific} (562) ^{inf} , 326 ^{dis}
Middle Eastern Crescent		
Egypt	0.65	Fahmy <i>et al.</i> 1977 ^{d,age,sex-specific; f} (1998), Southgate, 1979 ^d , El-Ridi <i>et al.</i> 1984 ^{d,f,g} (3215), Feinsod <i>et al.</i> 1987 ^{d,f,sex-specific; g,age-specific} (191), Wahdan, 1991 ^d (434,826) ³⁰
India	5.36	Rahman & Bhattacharya, 1971 ^{d,age,sex-specific} (690) ² , Marwah, Rao & Gaur, 1972 ^{d,e,age,sex-specific; e,f,sex-specific; g} (7194) ¹⁵ , Chandra <i>et al.</i> 1973 ^d (531) ³ , Bhattacharya & Gubler, 1973 ^{d,age,sex-specific} (1338) ² , Russel <i>et al.</i> 1975 ^a (6781) ⁵ , Dondero <i>et al.</i> 1976 ^a (660), Rao <i>et al.</i> 1980 ^a (11,300), Rajagopalan <i>et al.</i> 1981 ^{d,age,sex-specific} (10,522), Rath <i>et al.</i> 1984 ^{d,e,age,sex-specific; f,sex-specific; g} (9323), Sharma <i>et al.</i> 1984 ^{d,e} , Sarma <i>et al.</i> 1987 ^{d,age,sex-specific; e,age-specific; f,sex-specific; g} (2921), Biswas <i>et al.</i> 1989 ^{d,e} , Jain <i>et al.</i> 1989 ^{d,age,sex-specific} (11,105), Rajagopalan <i>et al.</i> 1989 ^{d,age,sex-specific} (24,946), Hyma, Ramesh & Gunasekharan, 1989 ^{d,age,sex-specific; e,f,sex-specific; g} (93,402), Ramaiah <i>et al.</i> 1989 ^{d,age,sex-specific; e,f,sex-specific; g} (7976), Kumar & Chand, 1990 ^{d,age,sex-specific} (13,918), Pani <i>et al.</i> 1991 ^a (34,615) ^{inf} , 6493 ^{dis}
China	0.48	Self, 1991 ^d
Other Asia and Islands		
Bangladesh	2.19	Barry <i>et al.</i> 1971 ^{a,f,sex-specific; g} (9624), Wolfe & Aslamkhan, 1971 ^{d,sex-specific; institutions} (4190) ³⁹ , Wolfe & Aslamkhan, 1972 ^{a,f,sex-specific; g} (1443) ²
Burma	0.10	Hawking, 1976 ^d (880,258) ¹⁴
Other Polynesia	22.77	Ciferri <i>et al.</i> 1969 ^{d,f,g} (826) ^{4,inf} , 1008 ^{4,dis} , Hawking & Denham, 1971 ^d (Nauru, Tokelau(326), Wallis Islands), Weller <i>et al.</i> 1982 ^{d,e,age-specific} (459)

Table 1. (cont.)

Region ⁱ and country	Prevalence ⁱⁱ (%)	Sources
Fiji	10.69	Hawking & Denham, 1971 ^a , Mataika <i>et al.</i> 1971 ^{b,d,age,sex-specific} (4483), Mataika <i>et al.</i> 1971 ^{a,f,age,sex-specific}
French Polynesia	12.62	Laigret, 1984 ^{d,f,g} , Perolat <i>et al.</i> 1986 ^d
Indonesia	1.77	van Djik, 1961 ^a (1206), Self <i>et al.</i> 1978 ^a (536), Joesef & Cross, 1978 ^d (14,219) ¹⁸ , Harbut, 1983 ^{d,age,sex-specific} (358) ²
Laos	0.07	Tawil, 1979 ^c (2339)
Malaysia	0.40	Hii <i>et al.</i> 1985 ^{d,age,sex-specific; f,sex-specific; g} (1680)
Nepal	0.84	Jung, 1973 ^a (5302)
New Caledonia	18.72	Hawking & Denham, 1971 ^{d,f}
Other Micronesia	3.07	Hawking & Denham, 1971 ^d
Micronesia	11.12	Hawking & Denham, 1971 ^d
Papua New guinea	72.02	Knight <i>et al.</i> 1979 ^a (233), Kazura <i>et al.</i> 1984 ^a (99)
Philippines	4.12	Rozeboom & Cabrera, 1964 ^{d,age,sex-specific} (280), Cabrera & Jueco, 1972 ^{d,age,sex-specific} (248), Wenclesloa <i>et al.</i> 1972 ^{d,age,sex-specific} (7,222), Grove <i>et al.</i> 1978 ^a (555), Enarson & Enarson, 1982 ^a (590), Ishii <i>et al.</i> 1983 ^{d,age,sex-specific} (267), Schultz, 1988 ^{d,age,sex-specific} (1546) ⁸
Western Samoa	7.73	Kimura, Penaia & Spears, 1985 ^a (8385) ²⁸
Seychelles	10.02	Lambrecht, 1971 ^{d,age-specific} (2474) ⁶
Sri Lanka	1.62	Hawking, 1976 ^d , Dissayanake, 1989 ^{d,age,sex-specific; f,sex-specific} (4004) ²
Thailand	2.82	Harinasuta <i>et al.</i> 1970 ^{d,age,sex-specific; f,sex-specific; g,age-specific} (1549)
Republic of Tongo	48.04	Desowitz & Hitchcock 1974 ^{d,f,age,sex-specific} (296), Desowitz, Berman & Puloka, 1976 ^{d,f,g,age-specific} (606) ²
Vanuatu	11.99	Bouree, Sauvagnac & Montaville, 1987 ^{d,f} (7137) ¹³
Vietnam	1.35	Hawking, 1976 ^d
(B) Brugian filariasis		
India	0.30	Russel <i>et al.</i> 1980 ^c (2542), Chandrasekharan, Balaraman & Rao, 1981 ^{d,f,age-specific} (533), Rath, Mohapatra & Das, 1989 ^d (529), Raina <i>et al.</i> 1990 ^{d,age,sex-specific} (2186), Pani <i>et al.</i> 1990 ^{f,age,sex-specific} (7766), Srividya <i>et al.</i> 1991 ^{a,d,age,sex-specific} (22,579)
China	0.37	Anti-epidemic station, Zhejiang Province 1981 ^d , Ling, 1985 ^d (2100), Liu, Liu & Chen, 1989 ^d (494,000)
Other Asia and Pacific Islands		
Indonesia	2.58	Partono <i>et al.</i> 1972 ^a (635) ³ , Partono <i>et al.</i> 1977 ^{a,d,age,sex-specific; f} (3658) ⁷ , Partono <i>et al.</i> 1977 ^{b,d,age,sex-specific; f,sex-specific} (2764) ⁷ , Joesef & Cross, 1978 ^d (125,312) ¹¹⁰ , Sudomo <i>et al.</i> 1982 ^{d,age,sex-specific} (197) ²
Malaysia	1.88	Mak <i>et al.</i> 1977 ^{d,age,sex-specific; f,sex-specific} (858), Hii 1978 ^{d,sex-specific} (1305) ¹⁷ , Rubis <i>et al.</i> 1981 ^{d,age,sex-specific} (1613) ⁷
Thailand	0.39	Guptavanji <i>et al.</i> 1977 ^a (5021) ^{2,c} (36,457) ⁸ , Harinasuta & Sucharit, 1977 ^{d,age,sex-specific; f,sex-specific} (4291) ¹¹
Philippines	0.82	Wencleslao <i>et al.</i> 1972 ^{d,age,sex-specific} (14,543) ³⁹ , Cabrera & Jueco, 1972 ^{d,age,sex-specific} (248) ²
Republic of Korea	0.03	Kim, No & Lee, 1987 ^d (1863) ²
Vietnam	0.77	Hawking, 1976 ^d

ⁱ As defined for the Global Burden of Disease Study (World Bank, 1993).

ⁱⁱ Estimates of disease prevalence are for both sexes combined following the methods described in the text.

Superscripts in literature citations

^a Study provides age-sex stratified data for both infection and disease.

^b Study provides age-stratified data on infection and disease.

^c Study provides overall prevalence of infection and disease.

^d Study contains prevalence data on infection.

^e Study contains prevalence data on overall disease. This pertains to the case in bancroftian filariasis when disease prevalence figures refer to hydrocele and lymphoedema combined.

^f Study contains prevalence data on lymphoedema.

^g Study contains prevalence data on hydrocele.

When the superscripted letters are followed by suffixes, age and/or sex-specific, they refer to the availability of the corresponding age/sex stratified data. Thus, ^{d,age,sex-specific} specifies the availability of infection prevalence data stratified by age and sex.

Numbers with brackets denote the number of subjects surveyed and examined. Note that some studies do not provide sample sizes. Superscripted numbers (> 1) following bracketed figures represent the number of separate surveys reported in the citation. The sample sizes listed are simply the sum of the separate samples examined. Note that in some studies, sample sizes of infection and clinical or disease surveys differed. Thus (562^{inf}, 326^{dis}) means that 562 individuals were examined for infection status, while 326 individuals were examined for the presence of disease conditions.

Countries in Africa potentially at-risk of infection for which there are either no data or recent information (Hawking, 1977) include: Ascension, Botswana, Burundi, Djibouti, Equatorial Guinea, Gabon, Ghana (but see Gyapong *et al.* 1993), Lesotho, Reunion, Rwanda, St Helena, Somalia, Tristan da Cunha, Uganda and Zaire.




Fig. 1. The relationship between overall microfilarial prevalence (both sexes combined) and male infection proportion. The line represents the weighted best least squares fit ($t = 100.39$, D.F. = 95, $P < 0.0001$) of the linear regression model through the origin (equation (2) in the text) with coefficients: $E(Y_i) = 1.07 X_i$. The data are from 96 separate studies from Sub-Saharan Africa, (■), Other Asia and Islands (+) and India regions (◇). Analysis of co-variance indicated no significant regional differences in the observed relationship ($F = 1.503$, D.F. = 2, 93, $P > 0.5$), as can also be seen clearly from the graph (see however Hedges & Olkin (1985) for discussion on the validity of the F-test in meta-regression). Note that the estimated relationship suggests significantly higher infection in males (slope significantly greater than 1, $t = 7.00$, D.F. = 95, $P < 0.001$). Data sources: SSA (30 data points): Blanchy & Galtier, 1982; Akogun, 1991; Udonsi 1988a, b; Matola, 1985; Wijeyaratne *et al.* 1982; Ripert *et al.* 1982; Kuhlow & Zielke, 1976; Zielke & Chlebowsky, 1979; Brunhes, 1975; Gyapong *et al.* 1993; Brengues, 1975; McMahon *et al.* 1981; Wijers & Kinyanjui, 1977; OAI (33 data points): Ciferri *et al.* 1969; Mataika *et al.* 1971b; Rosen, 1955; Beye *et al.* 1953; van Dijk, 1961; Cartel *et al.* 1992; Self *et al.* 1978; Barry *et al.* 1971; Schultz, 1988; Wolfe & Aslamkhan, 1972; Rozebroom & Cabrera, 1964, 1965; Enarson & Enarson, 1982; Harinasuta *et al.* 1970; Jachowski & Otto, 1955; David & Edeson, 1965; Kazura *et al.* 1984; Tanaka *et al.* 1980; Mahfuddin *et al.* 1977; Ishii *et al.* 1983; McCarthy & Fitzgerald, 1955; Knight *et al.* 1979; Lie, Hudajo & Amaliah, 1960; Symes, 1960; Iyengar *et al.* 1959; IND (33 data points): Bhattacharya & Gubler, 1973; Dondero *et al.* 1976, Jain *et al.* 1989; Nair, 1960; Basu *et al.* 1965; Jain *et al.* 1987; Rajagopalan, Shetty & Arunachalam, 1981; Rajagopalan *et al.* 1989, 1971; Joseph & Peethambaran, 1963; Kant, Sen & Puri, 1956; Krishnaswami, 1955; Kumar & Chand, 1990; Nair, 1962, Rahman & Bhattacharya, 1971; Rath *et al.* 1984; Singh, Raghavan & Krishnaswami, 1956; Singh *et al.* 1963, 1964; Varma *et al.* 1960; Varma, Dass & Sinha, 1961a, b, c; Ramaiah *et al.* 1989; Chand *et al.* 1961, 1962; Russel, Das & Rao, 1975; Sarma *et al.* 1987.

occurrence of infection (defined as positive for microfilaria) and chronic disease (restricted to hydrocele and lymphoedema). The secondary aim is to

address the methodological issues of rate projections and aggregations arising from using the published data on community infection and disease prevalences in endemic countries. This work forms part of the Global Burden of Disease study, a World Bank initiative to develop indices of disability for the purposes of comparison between different diseases and illnesses (World Bank, 1993). It also forms part of a WHO initiative to update the existing estimates of the global prevalence of infection (WHO, 1984, 1992).

MATERIALS AND METHODS

Data sources

The estimates derived here are for 6 out of the 8 regions defined for the Global Burden of Disease study (see World Bank, 1993). The Established Market Economies and Formerly Socialist Economies of Europe are excluded as infection is not considered endemic in these regions (WHO, 1984, 1992). The data used for the estimates are prevalence values of microfilaraemia and chronic disease derived from published community-based studies in each country. They are based on extensive computer literature searches complemented with reference tracing and scrutiny of reports held by the WHO. As far as possible, the most recent data published for each country were used, older data (largely restricted to the 1960s) were used only when no recent data were available for a particular country.

The above searches located some 118 publications (published between 1953 and 1991) with relevant data for disease estimation. A complete bibliography, including the data content of each study, is given in Table 1, and highlights the considerable variation in the information content of published studies. Only 20 studies provide complete information on the age-sex distribution for both microfilaraemia and the 2 forms of chronic disease in the community, the majority of studies tending to give only partial (with a greater bias for infection than disease) or no age-data (see Table 1). The studies also varied considerably in reporting infection and disease prevalences by gender, with most studies presenting only combined results for the sexes. Most studies provide results for 'combined' disease, or give results for either lymphoedema or hydrocele but not for both. Most publications were concerned with one survey only; however, a few dealt with several surveys either within the same country or within a group of countries. The number of studies available for each country also varied (Table 1).

Methodological approach

Overview. The methodology developed is primarily founded on the need to extrapolate individual study prevalences to larger populations at the national and




Fig. 2. The relationship between overall infection prevalence (combined sexes) and overall disease proportion (combined sexes) for (A) Sub-Saharan Africa (17 studies), (B) Other Asia and Islands (15 studies) and (C) India (19 studies) regions. Lines show the best fit linear regression models (equation (1)) for each region yielding the following equations: SSA = $E(Y_i) = -0.20 + 0.25X_i$ ($t = 25.00$, D.F. = 15, $P < 0.0001$), OAI = $E(Y_i) = 3.4 + 0.16X_i$ ($t = 4.85$, D.F. = 13, $P < 0.001$), IND = $E(Y_i) = -0.21 + 0.60X_i$ ($t = 7.00$, D.F. = 17, $P < 0.001$). Note the high degree of between-study variability observed for the relationship in each region. The asterisks mark those studies that contribute more than 5% each to the overall heterogeneity about the regression line, and hence have the least weights in the regression. The fitted lines clearly show that the influence of these studies is heavily discounted during the weighted least squares fit of the models. The estimated relationships indicate slight regional differences ($F = 10.806$, D.F. = 2, 43, $P < 0.01$, steeper rise in disease with infection in the India and Sub-Saharan Africa regions but higher disease prevalence at lower infection in the Other Asia and Islands region), facilitating the incorporation of some geographical variation in the estimation of disease prevalences. *Data sources:* Akogun, 1991; Matola, 1985; Brunhes, 1975; Gyapong *et al.* 1993; Brengues, 1975; McMahon *et al.*

regional levels, and the need to deal with the incomplete age-gender-disease information of published studies (see Table 1). These considerations required the development of the following 3-phased approach to burden estimation. First, to incorporate gender and disease-specific (hydrocele and lymphoedema) distributions into those studies lacking this information. Second, to derive age-specific prevalences for each condition in each study and, third, to sum these individual study prevalences within age groups to provide age-specific prevalences at the country, regional and global levels, or across age-groups to provide a summary age-adjusted figure. The approach is described in detail in the following 3 sections.

Incorporation of gender and disease-specific distributions. This primarily required the estimation of (a) the relationship between overall point prevalences of infection and chronic disease, (b) the sex ratios of infection and disease prevalences and (c) in the case of bancroftian filariasis, the ratio of hydrocele to lymphoedema for a given overall male disease prevalence. We derive simple linear regression models to estimate these relationships taking appropriate account, however, of the considerable between-study variation in sample size (Table 1, see Hedges & Olkin, 1985). Specifically, given the non-constant error variance, we used a weighted least-squares regression model of the form :

$$E(Y_i) = \beta_0 + \beta_1 X_i + e_i \quad i = 1, \dots, N, \quad (1)$$

where $E(\cdot)$ denotes expected value; Y_i is the prevalence of overall infection or disease, the prevalence of male infection or disease, or the prevalence of hydrocele as a proportion of overall male disease for study i ; X_i is the overall microfilarial or infection prevalence, the prevalence of overall disease (undifferentiated by gender), or the prevalence of male disease (both hydrocele and lymphoedema) observed in each study; N is the number of studies; and $\text{var}(e_i) = \sigma^2/w_i$, where σ^2 represents the error variance or residual mean square of the model (Weisberg, 1985; Hedges & Olkin, 1985). Thus,

1981; Wijers & Kinyanjui, 1976; van Dijk, 1961; Wolfe & Aslamkhan, 1972; Mahfuddin *et al.* 1977; McCarthy & Fitzgerald, 1955; Weller *et al.* 1982; de Rook, 1959; Knight *et al.* 1979; Kazura *et al.* 1984; Enarson & Enarson, 1982; Barry *et al.* 1971; Self *et al.* 1978; Ciferri *et al.* 1969; Iyengar *et al.* 1959; Galliard, Mille & Robinson, 1949; Dondero *et al.* 1976; Nair, 1960, 1962; Joseph & Peethambaran, 1963; Krishnaswami, 1955; Rath *et al.* 1984; Varma *et al.* 1960; Ramaiah *et al.* 1989; Chand *et al.* 1961, 1962; Iyengar, 1952; Srivastava & Prasad, 1962; Singh *et al.* 1963, 1964; Sarma *et al.* 1987; Rahman, Singh & Srivastava, 1959.




Fig. 3. The relationship between overall disease prevalence (combined sexes) and male disease proportion. Data from studies in Sub-Saharan Africa (■), data from studies in Other Asia and Islands (+), data from India (◇). Lines indicate the best fit linear model (equation (2)) in each region: SSA = $E(Y_i) = 1.8X_i$ ($t = 36.38$, D.F. = 15, $P < 0.0001$), OAI = $E(Y_i) = 1.7X_i$ ($t = 39.22$, D.F. = 9, $P < 0.0001$), IND = $E(Y_i) = 1.2X_i$ (D.F. = 16, $P < 0.0001$). As in Fig. 3, the asterisks mark studies contributing to $> 5\%$ of the total variance about the regression line. Male disease in the community appears to be significantly lower in India for a given overall disease prevalence ($F = 13.827$, D.F. = 2, 40, $P < 0.0001$). Data sources: SSA: Ufomadu *et al.* 1991; Matola, 1985; Brunhes, 1975; Gyapong *et al.* 1993; Brengues, 1975; McMahon *et al.* 1981; Wijers & Kinyanjui, 1977; Estambale *et al.* 1994; OAI: Ciferri *et al.* 1969; van Djik, 1961; Self *et al.* 1978; Barry *et al.* 1971; Enarson & Enarson, 1982; Kazura *et al.* 1984; Mahfuddin *et al.* 1977; McCarthy & Fitzgerald, 1955; Knight *et al.* 1979; Iyengar *et al.* 1959; IND: Pani *et al.* 1991; Dondero *et al.* 1976; Nair, 1960; Joseph & Peethambaran, 1963; Krishnaswami, 1955; Nair, 1962; Rath *et al.* 1984; Varma *et al.* 1960; Ramaiah *et al.* 1989; Chand *et al.* 1961, 1962; Iyengar, 1952; Singh *et al.* 1963, 1964; Russel *et al.* 1975; Sarma *et al.* 1987.

given the binomial response variable Y_i , approximate weights w_i are readily given by

$$w_i = \frac{1}{p_i(1-p_i)/n_i} \quad (2)$$

where $p_i(1-p_i)/n_i$ gives the binomial variance of p_i , and $p_i = Y_i$ is the studied prevalence or proportion, and n_i is the study sample size. The intuitive rationale for these weights is that reliable estimates (small variance) have relatively large weight and unreliable estimates (large variance) have relatively small weights in the respective regressions (Hedges & Olkin, 1985).

Analysis of geographical variation in the relationships was limited by restriction of available data to Sub-Saharan Africa, Other Asia and Pacific Islands and India. This was a particular problem for brugian filariasis, as we were able to locate only 11 studies

with concomitant information on overall infection and disease prevalences, and only 10 studies giving gender data on lymphoedema prevalences. This precludes the analysis of geographical variation for this disease, and we apply the estimated relationships globally to all relevant *Brugia* studies regardless of study region.

Figures 1–3 and Table 2 summarize the results of the linear regression analyses used to derive the relationships, together with the details of the studies used in the estimation. The results for bancroftian filariasis (Figs 2 and 3) indicate some variation in the infection–disease–gender relationships by region. Disease estimates for Latin America and the Caribbean, the Middle Eastern Crescent and China, for which no or incomplete data were available were extrapolated simply by using the mean overall global relationships in each case (data not shown).

Hydrocele cases as a proportion of male disease appeared independent of the overall prevalence of male disease in each region (none of the regressions were significant). Hence, the weighted mean proportion of hydrocele (as a proportion of male disease) was calculated from the studies in each region using the formula:

$$\hat{p}_{hj} = \frac{\sum_i w_i p_{hi}}{\sum_i w_i} \quad (3)$$

where \hat{p}_{hj} is the weighted mean proportion, p_{hi} is the proportion in each study i and w_i represent the study weights following equation (2). The results indicated that 94% (S.E. = 0.017, $n = 14$, data sources: Brengues, 1975; Brunhes, 1975; Wijers & Kinyanjui, 1977; McMahon *et al.* 1981; Matola, 1985; Ufomadu *et al.* 1991; Gyapong *et al.* 1993) and 93% (S.E. = 0.03, $n = 10$, data sources: Beye *et al.* 1952; McCarthy & Fitzgerald, 1955; Iyengar, de Rook & van Djik, 1959; van Djik, 1961; Ciferri *et al.* 1969; Barry, Ahmed & Khan, 1971; Wolfe & Aslamkhan, 1972; Mahfuddin *et al.* 1977; Self *et al.* 1978; Knight *et al.* 1979; Enarson & Enarson, 1982) of observed male disease prevalence may be attributed to hydrocele in Sub-Saharan Africa and Other Asia and Islands regions. In India, the corresponding proportion was significantly lower (Analysis of co-variance: $F = 32.60$, D.F. = 2, 35 $P < 0.0001$) at 80% (S.E. = 0.026, $n = 14$, data sources: Iyengar, 1952; Nair, 1960; Varma, Sinha & Dass, 1960; Chand *et al.* 1961; Chand, Singh & Vyas, 1959; Singh *et al.* 1963, 1964; Dondero *et al.* 1976; Rath *et al.* 1984; Sarma *et al.* 1987; Ramaiah *et al.* 1989).

Specific prevalence estimates were extrapolated from the empirical data using the above relationships in the following manner. Overall community infection or disease prevalences s_i , and male infection

Table 2. Results of weighted linear regressions used for estimating infection and disease prevalences from individual studies in brugian filariasis providing only overall infection and/or partial disease data^a(i) Relationship between overall community infection prevalence X_i and male infection prevalence Y_i^b

$$Y_i = 1.2X_i, t = 37.33 \text{ D.F.} = 14 \text{ } P < 0.0001$$

(ii) Relationship between community infection X_i and disease prevalence Y_i^c

$$Y_i = 0.5 + 0.11X_i, t = 2.220 \text{ D.F.} = 16 \text{ } P = 0.05$$

(iii) Relationship between overall lymphoedema X_i and male lymphoedema Y_i^d

$$Y_i = 1.13X_i, t = 13.07 \text{ D.F.} = 9 \text{ } P < 0.0001$$

^a Data are from studies from all *Brugia* endemic regions (OAI, CHN and IND).^b Regression (through the origin) of male infection prevalence on overall (combined sexes) infection prevalence; used to derive sex-specific estimates of infection from overall community infection prevalence. Note that the slope indicates significant higher infection in males. *Data sources*: Ramachandran *et al.* 1971; Barclay, 1969; Partono *et al.* 1972, 1977a, b, 1978; Kapojos *et al.* 1976; Guptavanij *et al.* 1971, 1977; Harinasuta *et al.* 1970; Raina *et al.* 1990; Srividya *et al.* 1991a; Wenceslao *et al.* 1972; Mak *et al.* 1977; Rubis *et al.* 1981.^c Regression of disease prevalence on infection prevalence; used to derive disease estimates from studies giving only overall infection prevalence (see text). *Data sources*: Barclay, 1969; Partono *et al.* 1972, 1977b; Guptavanij *et al.* 1971, 1977; Harinasuta *et al.* 1970; Dondero & Menon, 1972; Mak *et al.* 1977.^d Regression of male lymphoedema prevalence on overall lymphoedema prevalence. Slope value indicates significant higher disease in males. *Data sources*: Barclay, 1969; Partono *et al.* 1972, 1978; Guptavanij *et al.* 1971, 1977; Harinasuta *et al.* 1970; Dennis *et al.* 1976; Pani *et al.* 1990; Estambale *et al.* 1994.

or disease prevalences s_{im} , are obtained from the corresponding regression equations ($Y_i = s_i, s_{im}$) for the particular region (*i.e.* equation coefficients in Figs 1–3 and the corresponding mean global regression relationships). Similarly, hydrocele prevalence for a given male disease prevalence is calculated by direct multiplication with the mean proportions estimated for each region (equation (3)), and in the case of regions with incomplete data (Latin America and the Caribbean, China and Middle Eastern Crescent) by the global mean of 0.89 (S.E. = 0.025, $n = 38$). Overall female infection or disease prevalences, s_{if} , for study i are derived from estimated male disease prevalences, s_{im} , as follows

$$s_{if} = \hat{p}_i + (\hat{p}_i - s_{im}) \quad (4)$$

where $\hat{p}_i \doteq X_i$ is the observed overall disease prevalence for the study. Equation (3) is similarly adapted to calculate male lymphoedema prevalences from estimated hydrocele prevalences.

Generation of age-specific prevalences for each study. These were directly recorded to the nearest age-class of interest (0–4, 5–14, 15–44, 45–59 and 60+ yrs) for studies providing such data. Age-prevalences for studies that provided only crude overall infection and/or disease (point) prevalences were derived as follows. First, a search for studies with age-specific details of infection and disease was made for each region. Sufficient numbers of studies for bancroftian filariasis (see Table 3 for details) were found only for Sub-Saharan Africa, Other Asia and Pacific Islands and India. The age-specific prevalences from these studies, s_{ij} , where j denotes the age groups 0–4, 5–14, 15–44, 45–59 and 60+ for each study i , were then combined (see method below) to produce reference age-prevalence curves for each region. Age-weights, a_j , calculated from these curves were then applied to the crude overall community prevalences, s_i , of

studies in the corresponding region to derive infection and disease age-specific prevalences. Given s_i , the community prevalence, the age-specific prevalence for each condition is then given by

$$s_{ij} = s_i \cdot a_j \quad (5)$$

where the age-weights a_j are as shown in Table 3. Note that global weights were calculated (not shown) and applied to studies in Middle Eastern Crescent, China and Latin American and the Caribbean.

For brugian filariasis, there were insufficient studies to derive separate reference curves for Other Asia and Pacific Islands, India and China, the 3 endemic regions for this infection (WHO, 1992). In total, we were able to locate 5 studies with complete disease age-prevalence data for females and only 4 with similar disease (lymphoedema) age-details for males, all from Other Asia and Pacific Islands (Table 3). Thus, a common reference curve was derived and used for all 3 endemic regions.

Data aggregation to produce nation and regional age-prevalence estimates. National and regional age-prevalence estimates were obtained by combining the observed and derived age-specific data of individual studies. First, age-prevalences from each study in a particular country are averaged to obtain the corresponding national gender-disease specific prevalences by age-class, c_{ij} . For each age group, and for each sex and condition, this is given by

$$c_{ij} = \frac{\sum_i w_{ij} s_{ij}}{\sum_i w_{ij}} \quad (6)$$

where w_{ij} is the weight associated with age group j for study i and is given by equation (2); s_{ij} is the prevalence for age group j and study i ; and k denotes

Table 3. Age weights to generate age-specific disease prevalences from overall community prevalences

Species and region	Age class	Age-weight				
		Male			Female	
		Inf ¹	Hyd ²	Lymph ³	Inf ¹	Lymph ³
(A) <i>Wuchereria bancrofti</i>						
Sub-Saharan Africa	0-4	0.30	0.01	0.01	0.50	0.05
	5-14	0.50	0.10	0.02	0.60	0.10
	15-44	1.1	0.80	0.50	1.2	1.1
	45-59	1.4	1.9	1.9	1.6	3.7
	60+	1.8	2.4	3.4	1.6	3.9
Other Asia and Pacific Islands	0-4	0.30	0.01	0.01	0.50	0.10
	5-14	0.50	0.15	0.25	0.70	0.16
	15-44	1.2	1.2	1.1	1.2	1.1
	45-59	1.7	2.7	3.0	1.5	3.5
	60+	1.5	3.5	3.7	1.6	3.1
India	0-4	0.50	0.05	0.10	0.65	0.15
	5-14	0.75	0.20	0.30	0.90	0.35
	15-44	1.2	1.1	1.0	1.0	1.1
	45-59	1.3	2.1	2.3	1.3	2.2
	60+	1.2	2.5	2.9	1.7	2.7
(B) <i>Brugia malayi</i>						
(all regions)	0-4	0.20		0.0	0.35	0.0
	5-14	0.65		0.01	0.70	0.01
	15-44	1.2		0.90	1.3	0.70
	45-59	1.5		3.2	1.2	4.1
	60+	1.7		5.3	2.4	5.5

¹ Microfilarae + ves.² Hydrocele cases.³ Lymphoedema cases.

Data sources: SSA: McMahon *et al.* 1981; Brunhes, 1975; Gyapong *et al.* 1993; Udonso, 1986; Zielke & Chlebowosky, 1979; Brengues *et al.* 1969; Brengues, 1975; Oram, 1960; Akogun, 1991; Wijers, 1977; Wijers & Kinyanjui, 1977; OAI: Ciferri *et al.* 1969; Mataika *et al.* 1971a, b; Rosen, 1955 Beye *et al.* 1952, 1953; van Dijk, 1961; Cartel *et al.* 1992; Self *et al.* 1978; Barry *et al.* 1971; Schultz, 1988; Wolfe & Aslamkhan, 1972; Rozeboom & Cabrera, 1964, 1965; Enarson & Enarson, 1982; Harinasuta *et al.* 1970; Jachowski & Otto, 1955; David & Edeson, 1965; Kazura *et al.* 1984; Tanaka *et al.* 1980; Mahfuddin *et al.* 1977; Ishii *et al.* 1983; McCarthy & Fitzgerald, 1955, 1956; Abdulkader & Padley, 1960; Knight *et al.* 1979; Lie *et al.* 1960; Symes, 1960; IND: Bhattacharya & Gubler, 1972; Pani *et al.* 1991; Dondero *et al.* 1976; Jain *et al.* 1987, 1989; Nair, 1960, 1962; Basu *et al.* 1967; Rajagopalan, Kazmi & Mani, 1977; Rajagopalan *et al.* 1981, 1989; Joseph & Peethambaran, 1963; Kant *et al.* 1956; Krishnaswami, 1955; Kumar & Chand, 1990; Rahman & Bhattacharya, 1971; Rath *et al.* 1984; Singh *et al.* 1956; Varma *et al.* 1961a, b, c.

the number of studies. As before, this weighting scheme is used to discount the impact of small (and hence less reliable) studies (Hedges & Olkin, 1985).

The next step involved the estimation of the number of affected individuals in a particular country by gender and age-class from the estimated age-prevalences and demographic data. For each sex and disease, the numbers affected in each age-class is calculated from

$$T_{+ij} = c_{ij} \cdot T_{ij} \quad (7)$$

where T_{+ij} is the number of estimated infected or diseased individuals in age group j in country i , and T_{ij} is the age group population size. Where available, information on the size of the population at-risk of infection were used in these calculations (Table 4). In calculating the population at risk in China, it has been customary to ignore the populations of provinces having infection prevalences below 1%

(Self, 1991). Here, we have considered the population of any province with endemic infection to be potentially at risk.

Regional prevalences were calculated by summing the number of cases across the endemic countries of the region and dividing by the total regional population. Thus for each sex and disease condition, the age-specific regional prevalences R_{ij} are given by

$$R_{ij} = \frac{\sum_{i=1}^{N_+} T_{+ij}}{P_{ij}} \cdot 100 \quad (8)$$

where T_{+ij} is the number of affected individuals in each age group j for each endemic country i ; N_+ is the total number of endemic countries in the region; and P_{ij} is the total regional population in age-class j . This method provides a crude weighting for national population size differences.

Table 4. Estimates of at-risk population size for lymphatic filariasis by country

(Modified from information held by the Filariasis Expert Committee, WHO.)

Country	Population in millions ¹	At-risk population size (millions)
Bangladesh	106·656	68·316
Brazil	150·367	3·00
China	1133·693	805·111 (55·562)
Costa Rica	2·807	0·041
Dominican Republic	7·074	1·500
Egypt	52·063	3·700
Fiji	0·745	0·413
Guyana	0·799	0·650
Haiti	6·471	1·000
India	849·514	394·067 (152·110)
Indonesia	178·228	22·000 (22·000)
Kenya	24·158	9·856 ²
Malaysia	17·860	1·994 (1·994)
Myanmar	41·607	3·120
Nepal	18·916	1·018
Philippines	61·481	15·803 (15·803)
Sri Lanka	17·003	4·400
Tanzania	24·516	10·000
Thailand	55·854	9·000 (6·950)
Trinidad	1·237	0·004
Vietnam	66·311	10·007

¹ Population statistics from the World Bank (1993).² Proportion of population at-risk assumed to be similar to that of Tanzania. Numbers in brackets denote the estimated at-risk population size for brugian filariasis. The total national population is assumed to be the at-risk population for all other countries.

The estimation of overall filariasis cases. The procedures described above allow the extrapolation of separate estimates for each of the disease conditions observed in filariasis. To obtain an estimate of the overall number of filariasis cases (*i.e.* microfilarial and diseased cases combined), however, required the development of methods to control for the number of individuals with both types of condition (infection with disease symptoms) in order to avoid double counting. We address this by analysing available data for hydrocele (h_{mf+}) and lymphoedema (l_{mf+}) cases who are also microfilaraemic in order to estimate the weighted proportions of such cases (see equation (3)), and use these to predict the proportion of individuals with simultaneous disease and infection, h_{mf+i} , l_{mf+i} . (data sources: *hydrocele*: de Rook, 1957, 1959; Nair, 1960; van Dijk, 1961; Barry *et al.* 1971; Wolfe & Aslamkhan, 1972; Brengues, 1975; Dondero *et al.* 1976; Wijers & Kinyanjui, 1977; Abaru *et al.* 1980; McMahon *et al.* 1981; *lymphoedema*: Causey *et al.* 1945; de Rook, 1959; Nair, 1960; van Dijk, 1961; Barry *et al.* 1971; Brengues, 1975; Dondero *et al.* 1976; Wijers &

Kinyanjui, 1977; McMahon *et al.* 1981; Weller *et al.* 1982). The analysis indicated that on average 22 % of hydrocele patients are microfilaraemic, while up to 8·9 % of lymphoedema patients may be microfilaraemic. Overall filariasis cases in each study are then given for males, say, by :

$$T_{ijm} = s_{ijmf} + s_{ijh} + s_{ijl} - s_{ijmf} (s_{ijhmf+} + s_{ijlmf+}) \quad (9)$$

where T_{ijm} is total male filariasis cases in study i and age-group j ; s_{ijmf} is the observed age-specific male microfilarial prevalence; s_{ijh} is the hydrocele prevalence; s_{ijl} is the male lymphoedema prevalence; and s_{ijhmf+} and s_{ijlmf+} are the corresponding expected proportions of male microfilaraemic hydrocele and lymphoedema cases. Note that the proportion of infected individuals for both disease conditions is assumed to be independent of age-group, gender and the community prevalence of infection, as a first approximation (see Michael, Grenfell & Bundy, 1994). The overall filariasis cases at the country, regional and global levels are then calculated employing the same methodology as outlined for the separate infection and disease conditions.

Table 5. Global estimate of the number of cases and prevalence of filariasis (infection and chronic disease) due to *Wuchereria bancrofti* by region

(Estimates of the number of cases are given in millions while prevalences are in percentages (%). Figures are for both sexes combined.)

Region ^a	Population	Infections ^b	Lymphoedema	Hydrocele	Total no. of cases ^c	Percentage of total global cases
SSA	512	27.87	4.64	10.20	40.02 7.81	37.69
OAI	793	11.33	1.79	1.90	14.46 1.83	13.62
IND	849	29.46	6.58	12.88	45.53 5.36	42.88
CHN	1134	4.05	0.11	1.68	5.46 0.48	5.14
LAC	441	0.32	0.031	0.057	0.395 0.09	0.37
MEC	391	0.24	0.056	0.060	0.34 0.09	0.32
World	4119	73.27	13.21	26.79	106.19 2.58^d	

^a Regions as defined in the Global Burden of Disease study (World Bank, 1993). SSA = Sub-Saharan Africa, OAI = Other Asia and Islands, IND = India, CHN = China, LAC = Latin America and Caribbean, MEC = Middle Eastern Crescent.^b Microfilaraemia cases.^c Denotes infection and chronic disease cases, including hydrocele in males, combined as described in the text. Lower numbers denote the prevalence in each region.^d Represents the global prevalence for endemic populations only.Table 6. Global estimate of the number of cases and prevalence of filariasis (infection and chronic disease) due to *Wuchereria bancrofti* by age group

(Estimates of the number of cases are given in millions, prevalences are in percentages (%). Figures are for both sexes combined.)

Age class	Population ^a	Infections ^b	Lymphoedema	Hydrocele	Total no. of cases ^c	Percentage of total global cases
0-4	552	2.71	0.13	0.06	2.88 0.52	2.72
5-14	918	11.44	0.94	1.82	13.63 1.48	12.84
15-44	1932	42.76	6.45	15.62	60.72 3.14	57.18
45-59	432	10.70	2.57	5.65	18.13 4.20	17.07
60+	285	5.74	2.44	3.64	10.83 3.80	10.20
Total	4119	73.27	13.21	26.79	106.19 2.58^d	

^a Population in endemic regions only (see Table 5).^b Microfilaraemia cases.^c Denotes infection and disease cases, including hydrocele in males, combined as described in the text. Lower numbers represent the prevalence in each age group.^d Represents the global prevalence for endemic populations only.

RESULTS

The estimates of number of cases and prevalence of infection, lymphoedema and hydrocele in males are given in Tables 5-10. The results are in terms of the endemic populations only, i.e. those living in the regions where significant risk of infection exists. For

each parasite species, these estimates are given for different age classes, by gender and for different endemic regions.

The estimated total number of *W. bancrofti* infections is 73 million, while the numbers of lymphoedema and hydrocele cases are estimated to be 13 and 27 million respectively (Tables 5 and 6).

Table 7. Global estimate of the number of cases and prevalence of filariasis (infection and chronic disease) due to *Brugia malayi* by endemic region

(Estimates of the number of cases are given in millions while prevalences are in percentages (%). Figures are for both sexes combined.)

Region ^a	Population	Infections ^b	Lymphoedema	Total no. of cases ^c	Percentage of total global cases
OAI	793	5.07	1.21	6.18 0.78	47.84
IND	849	1.80	0.86	2.58 0.30	20.01
CHN	1134	3.49	0.73	4.15 0.37	32.16
World	2776	10.36	2.81	12.91 0.47 ^d	

^a Regions as defined in Table 5.^b Microfilaraemia cases.^c Denotes infection and chronic disease cases, including hydrocele in males, combined as described in the text. Lower numbers denote the prevalence in each region.^d Represents the global prevalence for endemic populations only.Table 8. Global estimate of the number of cases and prevalence of filariasis (infection and lymphoedema) due to *Brugia malayi* by age group

(Estimates of the number of cases are given in millions, prevalences are in percentages (%). Figures are for both sexes combined.)

Age class	Population ^a	Infections ^b	Lymphoedema	Total no. of cases ^c	Percentage of total global cases
0-4	340	0.40	0.02	0.42 0.12	0.71
5-14	578	1.45	0.07	1.52 0.26	2.63
15-44	1342	6.32	1.11	7.33 0.55	39.50
45-59	309	1.34	0.81	2.08 0.67	28.17
60+	207	0.83	0.80	1.56 0.75	28.51
Total	2776	10.36	2.81	12.91 0.46 ^d	

^a Population in endemic regions only (see Table 7).^b Microfilaraemia cases.^c Denotes infection and chronic disease cases, including hydrocele in males, combined as described in the text. Lower numbers denote the prevalence in each age group.^d Represents the global prevalence for endemic populations only.

The total number of cases (infection and chronic disease) arising from *W. bancrofti* infection, however, is estimated at 106 million. The estimates for *B. malayi* (Tables 7 and 8) indicate a global total of some 10 million cases of infection with a further 3 million cases suffering from lymphoedema. The total number of *B. malayi* filariasis cases (infection and chronic disease) is estimated to be around 13 million, yielding a total filariasis burden from both species combined of 119 million cases. The overall prevalence of filariasis cases is thus low: 2.6% in the exposed population or 2.0% globally for bancroftian filariasis, and 0.5% and 0.25% respectively for brugian filariasis.

The regional distribution of cases is shown in Tables 5 and 7 for bancroftian and brugian filariasis respectively. For bancroftian filariasis, India, with 45.5 million cases, and Sub-Saharan Africa, with 40 million cases, have similar burdens (Table 5). These regions account for 43% and 38% of the world burden. The highest prevalence of cases occurs in Sub-Saharan Africa (8% compared to 5% in India). The region with the third highest number of cases and prevalence for bancroftian filariasis is 'Other Asia and Islands' (Asia, excluding China, India and Pakistan and including Pacific islands) (Table 5). China, with a total burden of 5.5 million cases, comes next in order of importance, while lower burdens of

Table 9. Estimates of the number of cases, in millions, of filariasis infection and lymphoedema due to *Wuchereria bancrofti* by gender and region

Region ^a	Population		Infections ^b		Lymphoedema		Total no. of cases ^c	
	Males	Females	Males	Females	Males	Females	Males	Females
SSA	253	259	14.74	13.13	1.78	2.86	24.28	15.74
OAI	401	392	6.54	4.79	0.92	0.87	8.87	5.59
IND	440	410	17.00	12.46	2.60	3.98	29.43	16.10
CHN	585	549	2.25	1.80	0.06	0.05	3.62	1.84
LAC	220	221	0.19	0.13	0.014	0.017	0.25	0.15
MEC	198	193	0.13	0.11	0.027	0.029	0.21	0.13
World ^d	2096	2023	40.86	32.41	5.36	7.81	66.65	39.54
			1.95^e	1.60	0.26	0.39	3.18	1.95

^a Regions as defined in Table 5.^b Microfilaraemia cases.^c Infection and chronic disease cases, including hydrocele in males, combined as described in the text.^d Population in endemic regions only.^e This and other figures in bold represent the gender prevalences for endemic populations only.Table 10. Estimates of the number of cases, in millions, of filariasis infection and lymphoedema due to *Brugia malayi* by gender and region

Region ^a	Population		Infections ^b		Lymphoedema		Total no. of cases ^c	
	Males	Females	Males	Females	Males	Females	Males	Females
OAI	401	392	3.17	1.90	0.76	0.45	3.87	2.30
IND	440	410	1.11	0.69	0.58	0.28	1.63	0.95
CHN	585	549	2.24	1.25	0.46	0.27	2.65	1.50
World ^d	1426	1351	6.52	3.84	1.81	1.00	8.16	4.75
			0.45^e	0.28	0.13	0.07	0.57	0.35

^a Regions as defined in Table 5.^b Microfilaraemia cases.^c Infection and chronic disease cases combined as described in the text.^d Population in endemic regions only.^e This and other figures in bold represent the gender prevalences for endemic populations only.

infection and disease are observed for 'Latin America and Caribbean' and 'Middle Eastern Crescent' regions. Note that the entire burden (some 340 thousand cases) attributed to the 'Middle Eastern Crescent' region occurs in Egypt (Table 1).

The regional distribution of the estimates for brugian filariasis (Table 7) indicates that almost half of the global burden (48%) is confined to the 'Other Asia and Pacific islands' region, mainly the South-East Asian countries of Indonesia, Thailand, Malaysia and Philippines (Table 1). China and India account for the remaining 32% and 20% of the global burden respectively.

The national prevalence figures given in Table 1 need to be interpreted cautiously as they are likely to be considerably more imprecise than regional figures. In part, this reflects the smaller number of studies of varying sample sizes used in their estimation. However, since they also reflect extrapolations from specific study areas, a fuller treatment of precision would require more detailed information than currently available on the spatial distribution of

the disease within each country. Nonetheless, the present figures show that although there is wide variation, Sub-Saharan Africa proportionately possess the largest number of countries with moderate to high prevalence of bancroftian filariasis cases, with 16/28 countries indicating prevalences over 10%. Countries belonging to 'Other Asia and Pacific islands' seem to exhibit even greater variation, with some of the highest prevalences in the world apparently occurring in Papua New Guinea and Republic of Togo (Table 1). By contrast, the endemic countries of 'Latin America and Caribbean' and 'Middle Eastern Crescent' (Egypt) all tend towards low prevalence values. In general, national prevalences of brugian filariasis are low, irrespective of region (Table 1).

The age-specific estimates clearly show the marked age-dependency of infection and disease (Tables 6 and 8). For both species, infection prevalence tends to rise asymptotically with age, while chronic disease rises linearly. Numerically, the largest number of cases occur in the 15 to 44 year

age-groups, but this partially reflects their much larger sampling sizes since prevalences, especially of disease, tend to increase with age (Tables 6 and 8). About 57% of the total burden for bancroftian filariasis and 40% of the total brugian filariasis burden occur in the 15 to 44 year age-group. Similar age-patterns were also observed at the regional level (data not shown), with only slight differences in the degree of age dependency occurring between regions. There was also no gender-specific difference in this pattern.

The prevalence and number of total cases are significantly higher among males for both parasite species (Tables 9 and 10). On average, there are about 25% more cases of bancroftian infection among males and nearly twice as many males as females with *B. malayi* infections. Chronic disease due to bancroftian filariasis is also more prevalent among males. However, this is largely due to the incidence of hydroceles, since when the numbers of cases of lymphoedema are compared the bias appears to be in the opposite direction, with a significantly greater (1.5 times more) amount of disease among females (Table 9). This female bias for lymphoedema is largely the result of the estimates for Sub-Saharan Africa and India, the 2 regions with the highest infection and disease prevalences. All the other regions indicate either a male or only a slight (Middle Eastern Crescent) female bias. By contrast, both infection and lymphoedema cases are more prevalent among males in all endemic regions in the case of brugian filariasis (Table 10).

DISCUSSION

The present estimates suggest that a total of some 119 million individuals may be currently afflicted with lymphatic filariasis. This figure is higher than the most recent estimate by WHO suggesting a global total of 78.6 million cases (WHO, 1992). Two possibilities may account for this discrepancy. First, the increase in number of cases may be an inevitable result of the continuing increase in world population even when case prevalences remain unchanged. Second, the present estimates may be an over-estimation of the 'true' prevalence despite being based on a broader (and hence more representative) database.

Ascribing some level of statistical certainty to the present estimates, however, has proved difficult. The approach adopted here essentially involves data aggregation at different population levels and the use of empirical relationships between key parameters to estimate case prevalences when no local information exists. Both these procedures were adjusted for only one type of bias, namely sample size differences between studies. Although weighting positively in favour of larger and presumably more reliable studies, this adjustment does not control for other

biases in individual studies, such as the reliability of the sampling or diagnostic method used (Southgate, 1974; Sasa, 1976; Grenfell *et al.* 1990), demographic differences (Brabin, 1990; Pani *et al.* 1991), socio-economic status (Wijers & Kinyanjui, 1977; Grove, Valeza & Cabrera, 1978; Jain *et al.* 1989; Evans *et al.* 1993), and spatio-environmental factors (Grove, 1983; Southgate, 1992). Statistical adjustment for these factors, however, is hampered by the lack of relevant information on such biases in many of the published studies, and is also difficult to justify because of inadequate understanding of the epidemiology of infection and disease (Grove, 1983; Srividya *et al.* 1991b; Bundy, Grenfell & Rajagopalan, 1991; Grenfell & Michael, 1992; Michael *et al.* 1994).

It is clear that at least 2 areas warrant particular attention for the development of more reliable methods for extrapolating from published data. The first concerns the need for more accurate estimations of the epidemiological relationships between key variables, used to derive indirect estimates when local data are unavailable, and the second relates to the need to address the effects of spatial heterogeneity in infection and disease. Spatial heterogeneity in lymphatic filariasis and other helminth infections is a relatively neglected area of study (Booth & Bundy, 1992) but its resolution is not only important for more accurate estimations of global means but also for understanding the population risk factors underlying disease clustering. The resolution of both these questions, however, requires the collection of detailed, standardized data, and current attempts by the WHO to develop Rapid Assessment methods for determining community infection and disease prevalences can only aid this process.

Nonetheless, the present estimates have proved valuable in extending and in identifying areas of continuing weakness in our knowledge of the distribution, epidemiology and public health significance of lymphatic filariasis (WHO, 1994; World Bank, 1993). The estimates suggest that the prevalence of infection and disease is relatively low when considered at a global or regional scale, although it is clear that great variation may occur between endemic countries. Although this spatial heterogeneity may reflect the effect of sampling variation, it could also, on the other hand, suggest that the disease may be of considerable local importance, particularly in India, Africa and among the Pacific islands, where country-specific prevalences range from near 0% up to the very high rates of > 20% estimated for some African and Pacific island nations. Support for new national disease surveys, perhaps through the development and deployment of Rapid Assessments methods, will be crucial in addressing this problem.

A finding of potential importance regarding the distribution of bancroftian filariasis at the regional scale is that Sub-Saharan Africa has the highest over-

all prevalence, has proportionately more countries with high prevalences, and with 40 million cases, almost as many cases of infection and disease as India (45.5 million). This contrasts with previous estimates (WHO, 1984, 1992), which suggested that a significant majority of filarial infection and disease cases occurred in India. Given the potential public health significance of this result for Africa, it is clear that there is a particular need for more precise information on infection incidence in this region, including information to quantify current at-risk population sizes or, at the very least, to determine remaining endemic countries.

This work also provides, for the first time, some information on the age and gender distribution of infection and disease cases at the global, regional and national scales. This has proved vital for assessing the public health impact of the disease using metrics, such as the DALY (World Bank, 1993), which measures burden due to disease in terms of future stream of disability-free life lost (more details on disability derivation given in Michael *et al.* (1995)). The age-distributions show that, unlike most other nematode infections (Chan *et al.* 1994), lymphatic filariasis is an affliction of the adult and older age classes, particularly in the case of chronic disease. Furthermore, this age-pattern was consistent for both species, across regions, and for both sexes. This implies a relatively low disease impact for lymphatic filariasis.

The current estimates of disease necessarily, through lack of data, focus on the more gross manifestations of lymphoedema and hydrocele. How justified this limited focus may be is debatable given that other forms of disease, particularly 'acute' disease, such as adenolymphangitis (ADL) (WHO, 1994; Pani *et al.* 1991, 1995) and 'subclinical' pathology in microfilaraemics (Dreyer *et al.* 1992; Witte *et al.* 1993; Freedman *et al.* 1994), are thought to be significant additional sources of disability. The implications of these disease manifestations for calculating filariasis burden is discussed in detail by Michael *et al.* (1995), but it is clear that their inclusion will have a significant effect on estimates of disease burden, particularly as they are likely to appear earlier in life than chronic manifestations, are likely to be at least as prevalent (Pani *et al.* 1991), and appear to occur similarly in both genders (Pani *et al.* 1991, 1995).

A final implication of the estimates is that filarial disease, because of the general low prevalence, may be preventable using simple, low cost approaches to control (WHO, 1994). The local importance of disease, however, argues for a geographically targeted strategy for control.

We thank the Wellcome trust for financial support, and Drs C. P. Ramachandran and E. Ottesen of the WHO for helpful discussions and access to unpublished data on filariasis distribution held at the WHO.

REFERENCES

- ABARU, D. E., MCMAHON, J. E., DE C. MARSHALL, T. F., HAMILTON, P. J. S., VAUGHAN, J. P. & WEGESA, P. (1980). Tanzania Filariasis Project. Studies on microfilaraemia and selected clinical manifestations of Bancroftian filariasis. *Acta Tropica* **37**, 63–71.
- ABDULCADER, M. H. M. & PADLEY, R. (1960). Filariasis records in Ceylon. *Indian Journal of Malariology* **14**, 521–55.
- AKOGUN, O. B. (1991). Filariasis in Gongola State Nigeria I: Clinical and parasitological studies in Mutum-Biyu district. *Journal of Hygiene, Epidemiology, Microbiology and Immunology* **35**, 383–93.
- ANTI-EPIDEMIC STATION, ZHEJIANG PROVINCE (1981). Follow-up survey on the long-term effects of diethylcarbamazine (Hetrazan) - medicated salt in the control of malayan filariasis. *Chinese Journal of Preventive Medicine* **15**, 234.
- AZEVEDO, J. F. D. (1964). Distribution and incidence of filariae of the genera 'Wuchereria' and 'Brugia' in the Portuguese overseas territories. *Anais do Instituto Medicina Tropical* **21**, 313–19.
- AZEVEDO, J. F. D., PINHAO, R., MEIRA, M. & GARDETTE, M. (1969). Bancroftian and Malayan Filariasis in Overseas Portuguese Territories. *Anais da Escola Nacional de Saude Publica Medicina Tropical* **3**, 3–9.
- BARCLAY, R. (1969). Filariasis in Sabah, East Malaysia. *Annals of Tropical Medicine and Parasitology* **63**, 473–88.
- BARRY, C., AHMED, A. & KHAN, A. Q. (1971). Endemic filariasis in Thakurgaon, East Pakistan. *American Journal of Tropical Medicine and Hygiene* **20**, 592–7.
- BASU, P. C., RAS, V. N. & PATTANAYAK, S. (1967). Filariasis in Greater Bombay – results of a rapid survey conducted in June, 1965. *Bulletin of the Indian Society of Malaria and Communicable Disease* **4**, 296–317.
- BEYE, H. K., EDGAR, S. A., MILLE, R., KESSEL, J. F. & BAMBRIDGE, B. (1952). Preliminary observations on the prevalence, clinical manifestations and control of filariasis in the Society islands. *American Journal of Tropical Medicine and Hygiene* **1**, 637–61.
- BEYE, H. K., KESSEL, J. F., HEULS, J., THOORIS, G. & BAMBRIDGE, B. (1953). Nouvelles recherches sur l'importance, les manifestations cliniques, et la lutte contre la filariose a Tahiti, Oceanie francaise. *Bulletin de la Société de Pathologie Exotique* **1**, 144–63.
- BHATTACHARYA, N. C. & GUBLER, D. J. (1973). A survey for bancroftian filariasis in the Calcutta area. *Indian Journal of Medical Research* **61**, 8–11.
- BISWAS, H., SHARMA, S. P., DAS, M., RAO, V. G., YADAVA, R. L. & NARASIMHAM, M. V. (1989). Filariasis control in rural areas through detection and treatment with diethylcarbamazine. *Journal of Communicable Disease* **21**, 272–81.
- BLANCHY, S. & GALTIER, J. (1982). La filariose de Bancroft a Mayotte. Etat actuel de l'endemie filarienne et programme de lutte. *Cahiers ORSTOM ser Entomologie Medicale et Parasitologie* **20**, 153–9.
- BOOTH, M. & BUNDY, D. A. P. (1992). Comparative prevalences of *Ascaris lumbricoides*, *Trichuris trichiura* and hookworm infections and the prospects for combined control. *Parasitology* **105**, 151–7.
- BOUREE, P., SAUVAGNAC, B. & MONTAVILLE, B. (1987).

- Lymphatic filariasis in Vanuatu. *Bulletin de la Société de Pathologie Exotique et de ses Filiales* **80**, 634–45.
- BRABIN, L. (1990). Sex differentials in susceptibility to lymphatic filariasis and implications for maternal child immunity. *Epidemiology and Infection* **105**, 335–53.
- BRENGUES, J. (1975). La filariose de bancroft en afrique de L'ouest. *Memoires ORSTOM* **79**, 299pp.
- BRENGUES, J., SUBRA, R. & BOUCHITE, B. (1969). Etude parasitologique, clinique et entomologique sur la filariose de Bancroft dans le sud du Dahomey et du Togo. *Cahiers ORSTOM ser Entomologie Medicale et Parasitologie* **8**, 279–305.
- BRINKMANN, U. K. (1976). Epidemiological investigations of bancroftian filariasis in the coastal zone of Liberia. *Tropenmedizin und Parasitologie* **28**, 71–6.
- BRUNHES, J. (1975). La filariose de Bancroft dans la sous-region Malgache. *Memoires ORSTOM* **81**, 212pp.
- BRYGOO, E. R. (1958). La filariose humaine a Madagascar. *Archives de L'institut Pasteur de Madagascar* **26**, 23–39.
- BUNDY, D. A. P. & COOPER, E. S. (1989). *Trichuris* and trichuriasis in humans. *Advances in Parasitology* **28**, 107–73.
- BUNDY, D. A. P., GRENFELL, B. T. & RAJAGOPALAN, P. K. (1991). Immunoepidemiology of lymphatic filariasis: the relationship between infection and disease. In *Immunoparasitology Today*, (ed. Ash, C. & Gallagher, R. B.) pp. A71–A75. Cambridge: Elsevier Trends Journals.
- CABRERA, B. D. & JUECO, N. L. (1972). Filariasis survey among indigenous tribes of Palawan, Republic of the Philippines. *Southeast Asian Journal of Tropical Medicine and Public Health* **3**, 31–9.
- CARTEL, J. L., NGUYEN, N. L., SPIEGEL, A., MOULIA-PELAT, J. P., PLICHART, R., MARTIN, P. M. V., MANUELLAN, A. B. & LARDEUX, F. (1992). *Wuchereria bancrofti* infection in human and mosquito populations of a Polynesian village ten years after the interruption of mass chemoprophylaxis with diethylcarbamazine. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **86**, 414–16.
- CAUSEY, O. R., DEANE, M. P., DA COSTA, O. & DEANE, L. M. (1945). Studies on the incidence and transmission of filaria, *Wuchereria bancrofti*, in Belem, Brazil. *American Journal of Hygiene* **41**, 143–9.
- CHAN, M. S., MEDLEY, G. F., JAMISON, D. & BUNDY, D. A. P. (1994). The evaluation of potential global morbidity attributable to intestinal nematode infections. *Parasitology* **109**, 373–87.
- CHAND, D., SINGH, M. V., GUPTA, B. B. & SRIVASTAVA, R. N. (1961). A note on filariasis in Gonda Town (Uttar Pradesh). *Indian Journal of Malariology* **15**, 39–47.
- CHAND, D., SINGH, M. V. & VYAS, L. C. (1962). Filariasis in Gorakhpur district, Uttar Pradesh. *Indian Journal of Malariology* **16**, 269–76.
- CHANDRA, S., CHANDRA, R., KATIYAR, J. C., GOVIL, P., GEORGE, P. A. & SEN, A. B. (1973). Observations on filariasis in some villages around Lucknow, Uttar Pradesh. *Indian Journal of Medical Research* **61**, 1127–33.
- CHANDRASEKHARAN, A., BALARAMAN, K. & RAO, C. K. (1981). Decline in brugian filariasis in hill tribe settlements of Kerala. *Indian Journal of Medical Research* **73**, S82–S86.
- CHARAFOUDINE, H. & PESSON, B. (1986). Bancroftian filariasis in Anjouan (Comores). *Bulletin de la Société de Pathologie Exotique et de ses Filiales* **79**, 229–36.
- CIFERRI, F., SILIGA, N., LONG, G. & KESSEL, J. F. (1969). A filariasis-control programme in American Samoa. *American Journal of Tropical Medicine and Hygiene* **18**, 369–78.
- CROMPTON, D. W. T. (1988). The prevalence of ascariasis. *Parasitology Today* **4**, 162–9.
- CROMPTON, D. W. T. (1989). Prevalence of ascariasis. In *Ascariasis and its Prevention and Control* (ed. Crompton, D. W. T., Nesheim, M. C. & Pawlowski, Z. S.), pp. 45–59. London: Taylor & Francis.
- DAVID, H. L. & EDESON, J. F. B. (1965). Filariasis in Portuguese Timor, with observations on a new microfilaria found in man. *Annals of Tropical Medicine and Parasitology* **59**, 193–207.
- DENNIS, D. T., PARTONO, F., PURNOMO, ATMOSOEDJONO, S. & SAROSO, J. S. (1976). Timor filariasis: epidemiologic and clinical features in a defined community. *American Journal of Tropical Medicine and Hygiene* **25**, 797–802.
- DE ROOK, H. (1957). Filariasis on the island of Pam. *Tropical and Geographical Medicine* **9**, 197–212.
- DE ROOK, H. (1959). Filariasis in the village of Inanwatan (South coast of the Vogelkop, Netherlands New Guinea). *Tropical and Geographical Medicine* **11**, 313–31.
- DESOWITZ, R. S., BERMAN, S. J. & PULOKA, T. (1976). Hyperendemic subperiodic Bancroftian filariasis: a search for clinical and immunological correlates of microfilaremia. *Bulletin of the World Health Organization* **54**, 565–71.
- DESOWITZ, R. S. & HITCHCOCK, J. C. (1974). Hyperendemic bancroftian filariasis in the Kingdom of Tonga: the application of the membrane filter concentration technique to an age-stratified blood survey. *American Journal of Tropical Medicine and Hygiene* **23**, 877–9.
- DISSANAYAKE, S. (1989). Microfilaraemia, serum antibody and development of clinical disease in microfilaraemic subjects infected with *Wuchereria bancrofti* and treated with diethylcarbamazine citrate. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **83**, 384–8.
- DONDERO, T. J. & MENON, V. V. V. (1972). Clinical epidemiology of filariasis due to *Brugia malayi* on a rubber estate in West Malaysia. *Southeast Asian Journal of Tropical Medicine and Public Health* **3**, 355–65.
- DONDERO, T. J., JR., BHATTACHARYA, N. C., BLACK, H. R., CHOWDHURY, A. B., GUBLER, D. J., INUI, T. S. & MUKERJEE, M. (1976). Clinical manifestations of bancroftian filariasis in a suburb of Calcutta, India. *American Journal of Tropical Medicine and Hygiene* **25**, 64–73.
- DREYER, G., OTTESEN, E. A., GALDINO, E., ANDRADE, L., ROCHA, A., MEDEIROS, Z., MOURA, I., CASIMIRO, I., BELIZ, R. & COUTINHO, A. (1992). Renal abnormalities in microfilaraemic patients with bancroftian filariasis. *American Journal of Tropical Medicine and Hygiene* **46**, 745–51.
- EL-RIDI, A. M. S., ARAFA, M. S., EL-REFAIE, S. A. & ABOULMAGED, L. A. (1984). Bancroftian filariasis in young adult males from Nile delta. *Journal of the Egyptian Society for Parasitology* **14**, 471–6.
- ENARSON, D. A. & ENARSON, P. M. (1982). Filariasis in an

- upland population in the Philippines. *Tropical and Geographical Medicine* **34**, 353–8.
- ESTAMBALE, B. B. A., SIMONSEN, P. E., KNIGHT, R. & BWAYO, J. J. (1994). Bancroftian filariasis in Kwale district of Kenya. I. Clinical and parasitological survey in an endemic community. *Annals of Tropical Medicine and Parasitology* **88**, 145–51.
- EVANS, D. B., GELBAND, H. & VLASSOFF, C. (1993). Social and economic factors and the control of lymphatic filariasis: a review. *Acta Tropica* **53**, 1–26.
- FAHMY, M. A. M., ARAFA, M. S., MANDOUR, A. M. & MAKLOUF, L. M. (1977). Bancroftian filariasis in Assiut area, Upper Egypt. *Journal of the Egyptian Society for Parasitology* **7**, 3–9.
- FAIN, A., WERY, M. & TILKIN, J. (1969). Recherches sur les filarioses humaines dans la region de la Cuvette Centrale (Republique democratique du Congo). *Annales de la Société Belge de Médecine Tropicale* **49**, 629–48.
- FEINSOD, F. M., FARIS, R., GAD, A., EL SAID, S., SOLIMAN, B. A., ABD-EL AZEM, I. S. & SAAH, A. J. (1987). Clinical manifestations of *Wuchereria bancrofti* filariasis in an endemic village in the Nile delta. *Annales de la Société Belge de Médecine Tropicale* **67**, 259–65.
- FRANCO, O. & LIMA, M. D. (1966). Alguns aspectos das atividades contra a filariose bancroftiana no Brasil. *Revista Brasileira de Malariologia e Doencas Tropicas* **18**, 73–89.
- FREEDMAN, D. O., FILHO, P. J., BESH, S., SILVA, M., BRAGA, C. & MACIEL, A. (1994). Lymphoscintigraphic analysis of lymphatic abnormalities in symptomatic and asymptomatic human filariasis. *Journal of Infectious Diseases* **170**, 927–33.
- GALLIARD, H., MILLE, R. & ROBINSON, W. H. (1949). Filariasis due to *W. bancrofti* var. *pacifica* at Tahiti and in the Society Archipelago. *Tropical Diseases Bulletin* **47**, 60–1.
- GRENFELL, B. T., DAS, P. K., RAJAGOPALAN, P. K. & BUNDY, D. A. P. (1990). Frequency distribution of lymphatic filariasis microfilariae in human populations: population processes and statistical estimation. *Parasitology* **101**, 417–27.
- GRENFELL, B. T. & MICHAEL, E. (1992). Infection and disease in lymphatic filariasis – an epidemiologic approach. *Parasitology* **104**, S81–S90.
- GROVE, D. I. (1983). Selective primary health care: strategies for the control of disease in the developing world. VII. Filariasis. *Reviews of Infectious Diseases* **5**, 933–44.
- GROVE, D. I., VALEZA, F. S. & CABRERA, B. D. (1978). Bancroftian filariasis in a Philippine village: clinical, parasitological, immunological, and social aspects. *Bulletin of the World Health Organization* **56**, 975–84.
- GUYATT, H. L. & BUNDY, D. A. P. (1991). Estimating prevalence of community morbidity due to intestinal helminths: prevalence of infection as an indicator of the prevalence of disease. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **85**, 778–82.
- GUPTAVANI, P., HARINASUTA, C., SURATHIN, K. & VUTIKES, S. (1971). Studies on sub-periodic *Brugia malayi* in Southern Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health* **2**, 44–50.
- GUPTAVANI, P., HARINASUTA, C., SURATHIN, K., VUTIKES, S. & DEESIN, T. (1977). Studies on the prevalence of Malayan filariasis in South Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health* **8**, 42–52.
- GYAPONG, J. O., BADU, J. K., ADJEI, S. & BINKA, F. (1993). Bancroftian filariasis in the Kassena Nankana District of the upper east region of Ghana: a preliminary study. *Journal of Tropical Medicine and Hygiene* **96**, 317–22.
- HARBUT, C. L. (1983). Filariasis in West Irian, Indonesia: a survey of six villages. *American Journal of Tropical Medicine and Hygiene* **32**, 764–6.
- HARINASUTA, C., SUCHARIT, S., DEESIN, T., SURATHIN, K. & SAMRAN, V. (1970). Bancroftian filariasis in Thailand, a new endemic area. *Southeast Asian Journal of Tropical Medicine and Public Health* **1**, 233–45.
- HARINASUTA, C. & SUCHARIT, S. (1977). Observation on the reduction of the prevalence of Malayan filariasis in Chumphon province of South Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health* **8**, 179–84.
- HAWKING, F. (1976). The distribution of human filariasis throughout the world. Part II. Asia. *Tropical Diseases Bulletin* **73**, 967–1013.
- HAWKING, F. (1977). The distribution of human filariasis throughout the world. Part III. Africa. *Tropical Diseases Bulletin* **74**, 649–79.
- HAWKING, F. & DENHAM, D. A. (1971). The distribution of human filariasis throughout the world. Part I. The Pacific region, including New Guinea. *WHO/FIL/71.94* 31pp. (mimeogr).
- HEDGES, L. V. & OLKIN, I. (1985). *Statistical Methods for Meta-Analysis*, San Diego: Academic Press.
- HII, J., KAN, S., PEREIRA, M., PARMAR, S. S., CAMPOS, R. L. & CHAN, M. K. C. (1985). Bancroftian filariasis and malaria in island and hinterland populations in Sabah, Malaysia. *Tropical and Geographical Medicine* **37**, 93–101.
- HII, J. L. K. (1978). A re-survey of the prevalence of Malayan filariasis in south-west Sabah, Malaysia. *Medical Journal of Malaysia* **33**, 26–9.
- HUEHNS, E. R. (1953). Filariasis in Mauritius. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **47**, 549–55.
- HYMA, B., RAMESH, A. & GUNASEKARAN, K. (1989). Lymphatic filariasis in Madras, India. *Social Science and Medicine* **29**, 983–90.
- ISHII, A., CABRERA, B. D., SUGURI, S., KOBAYASHI, M., GO, T. G. & VALEZA, F. (1983). An epidemiological study of filariasis in Sorsogon province, Republic of the Philippines, with notes on experimental mosquito infection. *Journal of Tropical Medicine and Hygiene* **86**, 59–64.
- IYENGAR, M. O. T. (1952). Filariasis in the Maldives Islands. *Bulletin of the World Health Organization* **7**, 375–403.
- IYENGAR, M. O. T., DE ROOK, H. & VAN DJIK, W. J. O. M. (1959). Interruption of transmission of *Anopheles* borne filariasis by indoor residual spraying in Netherlands New Guinea. *Tropical and Geographical Medicine* **11**, 287–90.
- JACHOWSKI, L. A. & OTTO, G. F. (1955). Filariasis in American Samoa IV. Prevalence of microfilaria in the human population. *American Journal of Hygiene* **61**, 334–48.
- JAIN, D. C., RAVINDRANATH, T. C., GHOSH, T. K., PILLAI, K. B. T. & RAO, C. K. (1987). Prophylactic effect of

- diethylcarbamazine on *W. bancrofti* filariasis. *Journal of Communicable Disease* **19**, 317–21.
- JAIN, D. C., MENON, P. K., SETHUMADHAVAN, K. V., JOHNEY, V. M. & GHOSH, T. K. (1989). Epidemiology of bancroftian filariasis in a semi-urban community of Kerala State. *Journal of Communicable Disease* **21**, 265–71.
- JOESOE, A. & CROSS, J. H. (1978). Distribution and prevalence of cases of microfilaraemia in Indonesia. *Southeast Asian Journal of Tropical Medicine and Public Health* **9**, 480–8.
- JOIRE, E., JULIVEZ, J. & GALTIER, J. (1988). La filariose de bancroft a Mayotte. Evaluation de l'importance des complications cliniques. *Medecine et Maladies Infectieuses* **4**, 251–2.
- JOSEPH, C. & PEETHAMBARAN, P. (1963). A filariasis survey of Trichur (Kerala State). *Indian Journal of Malariology* **17**, 33–6.
- JUMINER, B., DIALLO, S. & DIAGNE, S. (1971). Le foyer de filariose lymphatique du secteur de Sandiara (Senegal). I. Evaluation de l'endemicite. *Archives de L'institute Pasteur de Tunis* **48**, 231–46.
- JUNG, R. K. (1973). A brief study of the epidemiology of filariasis in Nepal. *Journal of the Nepalese Medical Association* **11**, 155–68.
- KANT, C., SEN, S. K. & PURI, B. S. (1956). Filariasis in Patna (Bihar). Part 1. *Indian Journal of Malariology* **10**, 199–217.
- KAPOJOS, D. S., TUMEWU, M., KAPOJOS, F. X. & WAWORUNTU, H. C. (1976). Filariasis in the Minahassa. *Tropical and Geographical Medicine* **28**, 359–61.
- KAZURA, J. W., SPARK, R., FORSYTH, K., BROWN, G., HEYWOOD, P., PETERS, P. & ALPERS, M. (1984). Parasitologic and clinical features of bancroftian filariasis in a community in East Sepik province, Papua New Guinea. *American Journal of Tropical Medicine and Hygiene* **33**, 1119–23.
- KIM, J. S., NO, B. U. & LEE, W. Y. (1987). Brugian filariasis: 10-year follow-up study on the effectiveness of selective chemotherapy with diethylcarbamazine on Che Ju island, Republic of Korea. *Bulletin of the World Health Organization* **65**, 67–75.
- KIMURA, E., PENAI, L. & SPEARS, G. F. S. (1985). Epidemiology of subperiodic bancroftian filariasis in Samoa 8 years after control by mass treatment with diethylcarbamazine. *Bulletin of the World Health Organization* **63**, 869–80.
- KNIGHT, R., MCADAM, K. P. W. J., MATOLA, Y. G. & KIRKHAM, V. (1979). Bancroftian filariasis and other parasitic infections in the Middle Fly River region of Western Papua New Guinea. I. Clinical, parasitological and serological studies. *Annals of Tropical Medicine and Parasitology* **73**, 563–76.
- KNIGHT, R. (1980). Current status of filarial infections in The Gambia. *Annals of Tropical Medicine and Parasitology* **74**, 63–8.
- KRISHNASWAMI, A. K. (1955). Filariasis in Mangalore (South India). *Indian Journal of Malariology* **9**, 1–16.
- KUHLOW, F. & ZIELKE, E. (1976). Distribution and prevalence of *Wuchereria bancrofti* in various parts of Liberia. *Tropenmedizin und Parasitologie* **27**, 93–100.
- KUMAR, A. & CHAND, S. K. (1990). Prevalence of *Wuchereria bancrofti* infection in some coastal villages of Ganjam, Orissa. *Journal of Communicable Disease* **22**, 209–12.
- LAIGRET, J. (1984). The effects of treatment with annual doses of diethylcarbamazide citrate in the control of human lymphatic filariasis: experience in the Pacific islands. *WHO/FIL/84.177* 10pp. (mimeogr).
- LAMBRECHT, F. L. (1971). Preliminary report on the distribution and epidemiology of filariasis in the Seychelles islands, Indian Ocean. *Southeast Asian Journal of Tropical Medicine and Public Health* **2**, 222–32.
- LIE, K. J., HUDAJO, W. & AMALIAH, S. (1960). *Wuchereria bancrofti* infection in Djakarta, Indonesia. A study of some factors influencing its transmission. *Indian Journal of Malariology* **14**, 339–52.
- LING, J. H. (1985). Analysis of residual sources of infection in an area previously endemic for filariasis. *Chinese Journal of Epidemiology* **6**, 222.
- LIU, X. J., LIU, J. Y. & CHEN, Z. (1989). Control of filariasis in the Fujian Province. *Chinese Journal of Parasitic Disease Control* **2**, 65–8.
- MADRIGAL, R. B., OCAMPO, E. M. & MORA, R. A. (1979). El uso de la tecnica de Knott en una zona endemica de filariasis en Puerto Limon, Costa Rica. *Revista Cubana de Medicina Tropical* **31**, 165–8.
- MAHFUDDIN, H., PARTONO, F., PURNOMO & IDRIS, K. N. (1977). Parasitological and clinical survey of Bancroftian filariasis in Kepu district, Central Jakarta, Indonesia. *Southeast Asian Journal of Tropical Medicine and Public Health* **8**, 173–7.
- MAK, J. W., CHEONG, W. H., OMAR, A. H., SIVANANDAM, S. & MAHADEVAN, S. (1977). Filariasis in Perlis, Peninsular Malaysia. *Medical Journal of Malaysia* **31**, 198–203.
- MARCEAU, C., COUPRIE, B., COMBE, A., SAMEEKOBO, A., TRIBOULEY, J., PUEL, V., PIQUEMAL, A. & RIPERT, C. (1986). Epidemiology of bancroftian filariasis and onchocerciasis in the area of Tala Mokolo (North Cameroon). *Bulletin de la Société de Pathologie Exotique et de ses Filiales* **79**, 755–65.
- MARWAH, S. M., RAO, N. S. & GAUR, S. D. (1972). A study of filariasis in an aggregated community of Banaras Hindu University township. *Journal of the Indian Medical Association* **60**, 33–9.
- MATAIKA, J. U., DANDO, B. C., SPEARS, G. F. S. & MACNAMARA, F. N. (1971a). Mosquito-borne infections in Fiji. III. Filariasis in northern Fiji: epidemiological evidence regarding the mechanisms of pathogenesis. *Journal of Hygiene* **69**, 297–306.
- MATAIKA, J. U., DANDO, B. C., SPEARS, G. F. S. & MACNAMARA, F. N. (1971b). Mosquito-borne infections in Fiji I. Filariasis in northern Fiji: epidemiological evidence regarding factors influencing the prevalence of microfilaraemia of *Wuchereria bancrofti* infections. *Journal of Hygiene* **69**, 273–86.
- MATOLA, Y. G. (1985). Prospects of human malaria and bancroftian filariasis infections in the lower Rufiji basin, Tanzania. 2. Bancroftian filariasis. *Tropical and Geographical Medicine* **37**, 108–13.
- MCCARTHY, D. D. & FITZGERALD, N. (1955). Researches in Western Samoa. III. Observations on filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **49**, 82–9.
- MCCARTHY, D. D. & FITZGERALD, N. (1956). Habit, habitat and hyperfilaria in the epidemiology of filariasis in Western Samoa. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **50**, 58–65.
- MCMAHON, J. E., MAGAYUKA, S. A., KOLSTRUP, N., MOSHA,

- F. W., BUSHROD, F. M. & ABARU, D. E. (1981). Studies on the transmission and prevalence of Bancroftian filariasis in four coastal villages of Tanzania. *Annals of Tropical Medicine and Parasitology* **75**, 415–31.
- MICHAEL, E., GRENFELL, B. T. & BUNDY, D. A. P. (1994). The association between microfilaraemia and disease in lymphatic filariasis. *Proceedings of the Royal Society of London, B* **256**, 33–40.
- MICHAEL, E., BUNDY, D. A. P., OTTESEN, E. & RAMACHANDRAN, C. P. (1995). The global burden of lymphatic filariasis. In *The Burden of Disease* (ed. Murray, K. S. & Lopez, A. D.), Geneva: World Health Organization. (In press).
- NAIR, C. P. (1960). Filariasis in centrally administered areas. Part I. Filaria survey of Pondicherry settlement. *Indian Journal of Malariology* **14**, 233–52.
- NAIR, C. P. (1962). Filariasis in Kerala State. Part VII. Survey of Ponani (Palghat District). *Indian Journal of Malariology* **16**, 47–62.
- NATHAN, M. B., BECKLES, G., TIKASINGH, E. S., HAMILTON, P. J. S. & MONTEIL, S. (1982). Parasitological and clinical studies of *Wuchereria bancrofti* and *Mansonella ozzardi* in coastal north Trinidad, West Indies. *West Indies Medical Journal* **31**, 168–76.
- NATHAN, M. B. & STROOM, V. (1990). Prevalence of *Wuchereria bancrofti* in Georgetown, Guyana. *Bulletin of the Pan American Health Organization* **24**, 301–6.
- OOSTBRUG, B. F. J. (1985). Is Wuchereriosis a disappearing disease in Surinam? *Acta Liedensia* **53**, 37–50.
- ORAM, R. H. (1958). Filariasis on the North Nyasa Lake Shore. *The Central African Journal of Medicine* **4**, 99–103.
- ORAM, R. H. (1960). Filariasis on the North Nyasa Lake Shore (II). *The Central African Journal of Medicine* **4**, 144–5.
- PANI, S. P., KRISHNAMOORTHY, K., RAO, A. S. & PRATHIBA, J. (1990). Clinical manifestations in malayan filariasis infection with special reference to lymphoedema grading. *Indian Journal of Medical Research* **91**, 200–7.
- PANI, S. P., BALAKRISHNAN, N., SRIVIDYA, A., BUNDY, D. A. P. & GRENFELL, B. T. (1991). Clinical epidemiology of bancroftian filariasis: effect of age and gender. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **85**, 260–4.
- PANI, S. P., YUVARAJ, J., VANAMAIL, P., DHANDA, V., MICHAEL, E., GRENFELL, B. T. & BUNDY, D. A. P. (1995). Episodic adenolymphangitis and the stages of lymphoedema in bancroftian filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **89**, 72–4.
- PANIAGUA, F., GARCES, J. L., GRANADOS, C., ZUNIGA, A., RAMIREZ, M. & JIMENEZ, L. (1983). Prevalence of bancroftian filariasis in the city of Puerto Limon and the province of Limon, Costa Rica. *American Journal of Tropical Medicine and Hygiene* **32**, 1294–7.
- PARTONO, F., HUDOJO, OEMIJATI, S., NOOR, N., BORAHINA, CROSS, J. H., CLARKE, M. D., IRVING, G. S. & DUNCAN, C. F. (1972). Malayan filariasis in Margolembo, South Sulawesi, Indonesia. *Southeast Asian Journal of Tropical Medicine and Public Health* **3**, 537–47.
- PARTONO, F., OEMIJATI, S., HUDOJO, JOESOEF, A., SAJIDIMAN, H., PUTRALI, J., CLARKE, M. D., CARNEY, W. P. & CROSS, J. H. (1977a). Malayan filariasis in Central Sulawesi (Celebes), Indonesia. *Southeast Asian Journal of Tropical Medicine and Public Health* **8**, 452–8.
- PARTONO, F., OEMIJATI, S., HUDOJO, JOESOEF, A., CLARKE, M. D., DURFEE, P. T., IRVING, G. S., TAYLOR, J. & CROSS, J. H. (1977b). *Brugia malayi* in seven villages in South Kalimantan, Indonesia. *Southeast Asian Journal of Tropical Medicine and Public Health* **8**, 400–7.
- PARTONO, F., PURNOMO, PRIBADI, W. & SOEWARTA, A. (1978). Epidemiological and clinical features of *Brugia timori* in a newly established village, Karakuak, West Flores, Indonesia. *American Journal of Tropical Medicine and Hygiene* **27**, 910–15.
- PEROLAT, P., GUIDI, C., RIVIERE, F. & ROUX, J. (1986). Filariose de bancroft en Polynesie française. situation épidémiologique et perspectives après 35 ans de lutte. *Bulletin de la Société de Pathologie Exotique* **79**, 78–88.
- PINHAO, R. C. (1961). Incidência de filarioses no vale do Zambeze. *Anais do Instituto Medicina Tropical* **18**, 15–18.
- RACCURT, C. P. (1986). Lymphatic filariasis in Haiti – historical sequel or future public health problem in this part of the world. *Bulletin de la Société de Pathologie Exotique et de ses Filiales* **79**, 745–54.
- RACCURT, C. P., LOWRIE, R. C., JR., KATZ, S. P. & DUVERSEAU, Y. T. (1988). Epidemiology of *Wuchereria bancrofti* in Leogane, Haiti. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **82**, 721–5.
- RACCURT, C. P., MOJON, M. & HODGES, W. H. (1984). Parasitological, serological, and clinical studies of *Wuchereria bancrofti* in Limbe, Haiti. *American Journal of Tropical Medicine and Hygiene* **33**, 1124–9.
- RAHMAN, J., SINGH, M. V. & SRIVASTAVA, R. N. (1959). Filariasis in Bahraich Town (Uttar Pradesh). *Bulletin of the Natural Society of India on Malaria and Mosquito Diseases* **7**, 53–117.
- RAHMAN, N. M. I. & BHATTACHARYYA, M. N. (1971). A preliminary survey of filarial infection in a group of labour population at Majan Tea Estate in Lakhimpur District, Assam. *Journal of the Indian Medical Association* **56**, 363–6.
- RAINA, V. K., JOSHI, M. C., SINGH, S., JOSHI, R. D., BHATTACHARJEE, K. K., KUMAR, A. & VERGHESE, T. (1990). Epidemiology of *Brugia malayi* and its co-existence with *Wuchereria bancrofti* in and around Sillaberia PHC, District Midnapur, West Bengal. *Journal of Communicable Disease* **22**, 205–8.
- RAJAGOPALAN, P. K., KAZMI, S. J. & MANI, T. R. (1977). Some aspects of transmission of *Wuchereria bancrofti* and ecology of the vector *Culex pipiens fatigans* in Pondicherry. *Indian Journal of Medical Research* **66**, 200–15.
- RAJAGOPALAN, P. K., DAS, P. K., SUBRAMANIAN, S., VANAMAIL, P. & RAMAIAH, K. D. (1989). Bancroftian filariasis in Pondicherry, South India: 1. Pre-control epidemiological observations. *Epidemiology and Infection* **103**, 685–92.
- RAJAGOPALAN, P. K., SHETTY, P. S. & ARUNACHALAM, N. (1981). A filariasis survey in Pondicherry villages. *Indian Journal of Medical Research* **73**, 73–7.
- RAMACHANDRAN, C. P., DONDERO JR., T. J., MULLIN, S. W., SIVANANDAM, S. & STEVENS, S. (1971). Filariasis in a rubber plantation in Negeri Sembilan, West Malaysia. *The Medical Journal of Malaya* **25**, 273–7.

- RAMAIAH, K. D., PANI, S. P., BALAKRISHNAN, N., SADANANDANE, C., DAS, L. K., MARIAPPAN, T., RAJAVEL, A. R., VANAMAIL, P. & SUBRAMANIAN, S. (1989). Prevalence of bancroftian filariasis and its control by single course of diethyl carbamazine in a rural area in Tamil Nadu. *Indian Journal of Medical Research* **89**, 184–91.
- RAO, C. K., DATTA, K. K., SUNDARAM, R. M., RAMPRASAD, K., RAO, J. S., VENKATANARAYANA, M., NATH, V. V. N., RAO, P. K., RAO, K., DAS, M. & SHARMA, S. P. (1980). Epidemiological studies on bancroftian filariasis in East Godavari district (Andhra Pradesh) : baseline filariometric indices. *Indian Journal of Medical Research* **71**, 712–20.
- RATH, R. N., DAS, R. K., MISHRA, G., MOHAPATRA, B. N. & RAMAKRISHNA, C. (1984). Bancroftian filariasis in two selected rural communities in Puri District: Orissa - A comparative study of filariometric data. *Journal of Communicable Disease* **16**, 104–12.
- RATH, R. N., MOHAPATRA, B. N. & DAS, B. (1989). Detection of a new focus of *Brugia malayi* infection in Orissa. *Journal of Communicable Disease* **21**, 39–40.
- RIPERT, C., EONO, P., EONO, D., TRIBOULEY, J., APPRIOU, M. & ISSOUFA, H. (1982). Etude epidemiologique de la Bancroftose dans la vallee du Logone (Nord Cameroun). *Medecine Tropicale* **42**, 59–66.
- ROSEN, L. (1955). Observations on the epidemiology of human filariasis in French Oceania. *American Journal of Hygiene* **61**, 219–48.
- ROZEBOOM, L. E. & CABRERA, B. D. (1964). Filariasis in mountain province, Luzon, Republic of the Philippines. *Journal of Medical Entomology* **1**, 18–28.
- ROZEBOOM, L. E. & CABRERA, B. D. (1965). Filariasis caused by *Wuchereria bancrofti* in Palawan, Republic of the Philippines. *American Journal of Epidemiology* **81**, 216–21.
- RUBIS, P., SENG, C. M., JUTE, N. A. & JAU, J. L. (1981). Parasitological and entomological studies on filariasis in seven villages, Serian district, Sarawak, East Malaysia. *Southeast Asian Journal of Tropical Medicine and Public Health* **12**, 30–6.
- RUSSEL, S., RAO, A. R., KHAMRE, J. M., PRAHARAJ, B. C. & RAO, C. K. (1980). Disappearance of some *Brugia malayi* foci from Madhya Pradesh, Orissa and Tamil Nadu and co-existence of Malayan and Bancroftian filariasis in some pockets. *Journal of Communicable Disease* **12**, 183–7.
- RUSSEL, S., DAS, M. & RAO, C. K. (1975). Filariasis in Andaman and Nicobar islands. Part I. Survey findings - Nancowry, Terressa, Chowra, Carnicobar and Port Blair. *Journal of Communicable Disease* **1**, 15–30.
- SARMA, R. V. S. N., VALLISHAYEE, R. S., MAYURNATH, S., NARAYANAN, P. R., RADHAMANI, M. P. & TRIPATHY, S. P. (1987). Prevalence survey of filariasis in two villages in Chingleput district of Tamil Nadu. *Indian Journal of Medical Research* **85**, 522–30.
- SASA, M. (1976). *Human Filariasis - a Global Survey of Epidemiology and Control*, Baltimore, Maryland: University Park Press.
- SCHULTZ, G. W. (1988). A study of bancroftian filariasis on the islands of Batan and Rapu Rapu, Philippines. *Southeast Asian Journal of Tropical Medicine and Public Health* **19**, 207–14.
- SELF, L. S. (1991). Report to Expert Committee. *WHO/FIL - F3/81/5*. (mimeogr).
- SELF, L. S., USMAN, S., SAJIDIMAN, H., PARTONO, F., NELSON, M. J., PANT, C. P., SUZUKI, T. & MECHFUDIN, H. (1978). A multidisciplinary study on bancroftian filariasis in Jakarta. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **72**, 581–7.
- SHARMA, S. P., BISWAS, H., DAS, M., KOSARE, G. J. & SHARMA, G. K. (1984). Control of bancroftian filariasis with diethylcarbamazine in rural areas of Valsad district, Gujarat state. *Journal of Communicable Disease* **16**, 282–6.
- SHERLOCK, I. A. & SERAFIM, E. M. (1967). Condições atuais da filariose bancroftiana no estado da Bahia, Brasil. *Revista Brasileira de Malariologia e Doenças Tropicais* **19**, 373–95.
- SINGH, J., RAGHAVAN, N. G. S. & KRISHNASWAMI, A. K. (1956). Filariasis in Travancore-Cochin State. *Indian Journal of Malariology* **10**, 219–38.
- SINGH, M. V., RASTOGI, K. C., SINGH, R. P. & SRIVASTAVA, V. K. (1963). Observations on rural filariasis in Sitapur district (Uttar Pradesh). *Indian Journal of Malariology* **17**, 303–10.
- SINGH, M. V., RASTOGI, K. C., SINGH, R. P. & SRIVASTAVA, V. K. (1964). Observations on urban filariasis in Lucknow. *Bulletin of the Indian Society of Malariology and Communicable Disease* **1**, 141–7.
- SOUTHGATE, B. A. (1974). Problems of clinical and biological measurements in the epidemiology and control of filarial infections. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **68**, 177–86.
- SOUTHGATE, B. A. (1979). Bancroftian filariasis in Egypt. *Tropical Diseases Bulletin* **76**, 1045–68.
- SOUTHGATE, B. A. (1992). Intensity and efficiency of transmission and the development of microfilaremia and disease - their relationship in lymphatic filariasis. *Journal of Tropical Medicine and Hygiene* **95**, 1–12.
- SRIVASTAVA, R. N. & PRASAD, B. G. (1969). An epidemiological study of filariasis in the villages of the rural health training centre, Sarojini Nagar, Lucknow. *Indian Journal of Medical Research* **57**, 528–42.
- SRIVIDYA, A., KRISHNAMOORTHY, K., SABESAN, S., PANICKER, K. N., GRENFELL, B. T. & BUNDY, D. A. P. (1991a). Frequency distribution of *Brugia malayi* microfilariae in human populations. *Parasitology* **102**, 207–12.
- SRIVIDYA, A., PANI, S. P., RAJAGOPALAN, P. K., BUNDY, D. A. P. & GRENFELL, B. T. (1991b). The dynamics of infection and disease in bancroftian filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **85**, 255–9.
- SUDOMO, M., HANIFAH, A., MAK, J. W. & LIM, B. L. (1982). A study of Malayan filariasis in Lubuk Mumpo and Datar Lebar villages in Lais regency, North Bengkulu, Sumatera, Indonesia. *Southeast Asian Journal of Tropical Medicine and Public Health* **13**, 584–9.
- SYMES, C. B. (1960). Observations on the epidemiology of filariasis in Fiji. Part II. Laboratory studies and human infections. *Journal of Tropical Medicine and Hygiene* **63**, 31–44.
- TANAKA, H., NAKAI, H., OMOTO, K., SHIBUYA, T., HIRAI, M., MERCADO, A. S., PAGARAN, I. G. & EVIOTA, W. S. (1980). The high prevalence of *Wuchereria bancrofti* infections

- in indigenous tribes in northern Mindanao, Philippines. *Japanese Journal of Experimental Medicine* **50**, 85–9.
- TAWIL, N. A. (1979). Microfilaraemia survey in Vientiane, Laos. *Southeast Asian Journal of Tropical Medicine and Public Health* **10**, 483–5.
- UDONSI, J. K. (1986). The status of human filariasis in relation to clinical signs in endemic areas of the Niger delta. *Annals of Tropical Medicine and Parasitology* **80**, 425–37.
- UDONSI, J. K. (1988a). Filariasis in the Igwu river basin, Nigeria – an epidemiological and clinical study with a note on the vectors. *Annals of Tropical Medicine and Parasitology* **82**, 75–82.
- UDONSI, J. K. (1988b). Bancroftian filariasis in the Igwu basin, Nigeria: an epidemiological, parasitological, and clinical study in relation to the transmission dynamics. *Folia Parasitologica* **35**, 147–55.
- UFOMADU, G. O., NWOKE, B. E. B., AKOH, J. I., SATO, Y., EKEJINDU, G. O. C., UCHIDA, A., SHIWAKU, K., TUMBAU, M. & UGOMO, K. K. (1991). The occurrence of loiasis, mansonellosis and wuchereriosis in the Jarawa River Valley, Central Nigeria. *Acta Tropica* **48**, 137–47.
- VAN DIJK, W. J. O. M. (1961). Mass treatment of *Wuchereria bancrofti* filariasis with diethylcarbamazine. A trial at Inanwatan, western New Guinea. *Tropical and Geographical Medicine* **13**, 143–59.
- VARMA, B. K., SINHA, V. P. & DASS, N. L. (1960). Filariasis in Sultanganj and its suburbs (Bihar). *Bulletin of the Indian Society of Malariology and Communicable Disease* **8**, 149–52.
- VARMA, B. K., DASS, N. L. & SINHA, V. P. (1961a). Studies on the incidence and transmission of filariasis in Bhagalpur town (Bihar). *Indian Journal of Malariology* **15**, 185–94.
- VARMA, B. K., DASS, N. L. & SINHA, V. P. (1961b). Filariasis in the rural population around Bhagalpur Town. Part I. *Indian Journal of Malariology* **15**, 285–92.
- VARMA, B. K., DASS, N. L. & SINHA, V. P. (1961c). Filariasis in the rural population around Bhagalpur Town. Part II. *Indian Journal of Malariology* **15**, 293–9.
- VINCENT, A. L. (1987). A survey of Bancroftian filariasis in the Dominican Republic. *Journal of Parasitology* **73**, 839–40.
- WAHDAN, M. H. (1991). Updated information on lymphatic filariasis in Egypt. *WHO/FIL – F3/81/5*. (mimeogr).
- WARREN, K. S., BUNDY, D. A. P., ANDERSON, R. M., DAVIS, A. R., HENDERSON, D. A., JAMISON, D. T., PRESCOTT, N. & SENFT, A. (1993). Helminth infections. In *Disease Control Priorities in Developing Countries* (ed. Jamison, D. T., Mosley, W. H., Meashem, A. R. & Bodadilla, J. L.), pp. 131–160. Oxford: Oxford University Press.
- WEINSTOCK, H., PANIAGUA, F., GARCAS, J. L., ZUNIGA, A., GRANADOS, C. & HERNANDEZ, E. (1977). Bancroftian filariasis in Puerto Limon, Costa Rica. *American Journal of Tropical Medicine and Hygiene* **26**, 1148–52.
- WEISBERG, S. (1985). *Applied Linear Regression*, 2nd edn. New York: John Wiley & Sons.
- WELLER, P. F., OTTESEN, E. A., HECK, L., TERE, T. & NEVA, F. A. (1982). Endemic filariasis on a pacific island 1. Clinical, epidemiologic, and parasitologic aspects. *American Journal of Tropical Medicine and Hygiene* **31**, 942–52.
- WENCESLAO, J. M., OBAN, E. & CABRERA, B. D. (1972). Eastern Samar, the fourth endemic focus for Malayan filariasis in the Republic of the Philippines. *Southeast Asian Journal of Tropical Medicine and Public Health* **3**, 552–61.
- WITTE, M. H., JAMAL, S., WILLIAMS, W. H., WITTE, C. L., KUMARASWAMI, V., MCNEIL, G. C., CASE, T. C. & PANICKER, T. M. R. (1993). Lymphatic abnormalities in human filariasis as depicted by lymphoangioscintigraphy. *Archives of Internal Medicine* **153**, 737–44.
- WORLD HEALTH ORGANIZATION (1984). *Lymphatic filariasis. Fourth Report of the WHO Expert Committee on Filariasis*. WHO Technical Report Series 702. WHO, Geneva.
- WORLD HEALTH ORGANIZATION (1992). *Lymphatic Filariasis: The Disease and its Control*. WHO Technical Report Series 821. WHO, Geneva.
- WORLD HEALTH ORGANIZATION (1994). Lymphatic filariasis infection and disease: control strategies. *TDR/CTD/FIL/PENANG/94.1* (mimeogr).
- WORLD BANK (1993). *World Development Report 1993. Investing in Health*. New York: Oxford University Press.
- WIJERS, D. J. B. (1977). Bancroftian filariasis in Kenya 1. Prevalence survey among adult males in the Coast Province. *Annals of Tropical Medicine and Parasitology* **71**, 313–31.
- WIJERS, D. J. B. & KALELI, N. (1984). Bancroftian filariasis in Kenya V. Mass treatment given by members of the local community. *Annals of Tropical Medicine and Parasitology* **78**, 383–94.
- WIJERS, D. J. B. & KINYANJUI, H. (1977). Bancroftian filariasis in Kenya II. Clinical and parasitological investigations in Mambui, a small coastal town, and Jaribuni, a rural area more inland (coast province). *Annals of Tropical Medicine and Parasitology* **71**, 333–45.
- WIJEYARATNE, P. M., VERMA, O. P., SINGHA, P., OSUHOR, P. C., MOTHA, B., SAHA, A. L., SLOTBOOM, A. B., DELEON, A. & BANDIPO, A. B. (1982). Epidemiology of filariasis in Malumfashi district of northern Nigeria. *Indian Journal of Medical Research* **76**, 534–44.
- WOLFE, M. S. & ASLAMKHAN, M. (1971). Filariasis in East Pakistan. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **65**, 63–9.
- WOLFE, M. S. & ASLAMKHAN, M. (1972). Bancroftian filariasis in two villages in Dinajpur district, East Pakistan. I. Infections in Man. *American Journal of Tropical Medicine and Hygiene* **21**, 22–9.
- ZIELKE, E. & CHLEBOWSKY, H. O. (1979). Studies on bancroftian filariasis in Liberia, West Africa I. Distribution and prevalence in the Nord-Western Savanna Area. *Tropenmedizin und Parasitologie* **30**, 91–6.