

Effect of Copper on Expression of Functional Genes and Proteins Associated with *Bradyrhizobium diazoefficiens* Denitrification

Pedro J. Pacheco

Abstract

Nitrous oxide (N₂O) is a powerful greenhouse gas that contributes to climate change. Denitrification is one of the largest sources of N₂O in soils. The soybean endosymbiont *Bradyrhizobium diazoefficiens* is a model for rhizobial denitrification studies since, in addition to fixing N₂, it has the ability to grow anaerobically under free-living conditions by reducing nitrate from the medium through the complete denitrification pathway. This bacterium contains a periplasmic nitrate reductase (Nap), a copper (Cu)-containing nitrite reductase (NirK), a c-type nitric oxide reductase (cNor), and a Cu-dependent nitrous oxide reductase (Nos) encoded by the *napED-ABC*, *nirK*, *norCBQD* and *nosRZDFYLX* genes, respectively. In this work, an integrated study of the role of Cu in *B. diazoefficiens* denitrification has been performed. A notable reduction in *nirK*, *nor* and *nos* gene expression observed under Cu limitation was correlated with a significant decrease in NirK, NorC and NosZ protein levels and activities. Meanwhile, *nap* expression was not affected by Cu, but a remarkable depletion in Nap activity was found, presumably due to an inhibitory effect of nitrite accumulated under Cu-limiting conditions. Interestingly, a post-transcriptional regulation by increasing Nap and NirK activities, as well as NorC and NosZ protein levels, was observed in response to high Cu. Our results demonstrate, for the first time, the role of Cu in transcriptional and post-transcriptional control of *B. diazoefficiens* denitrification. Thus, this study will contribute by proposing useful strategies for reducing N₂O emissions from agricultural soils.

Keywords

Cu-containing nitrite reductase; enzymatic activity; gene expression; nitric oxide reductase; nitrous oxide reductase; periplasmic nitrate reductase

1 Introduction and State of Art

With a 300-fold greater global warming potential than carbon dioxide (CO₂), nitrous oxide (N₂O) is one of the main biogenic greenhouse gases (GHG), and has also been described as the biggest single cause of ozone depletion^[1]. N₂O emissions from human activities, fundamentally Agriculture, Forestry and Other Land Use (AFOLU), have notably increased since the Green Revolution in the early 60s. During the period 2007–2016, these activities represented 81% of the anthropogenic emissions of N₂O, according to the last special report by the Intergovernmental Panel on Climate Change^[2]. In particular, agriculture has become the major source of N₂O emissions, accounting for

approximately 78% of the anthropogenic N₂O sources^[2] because of a global agricultural intensification and a great increase in the non-synchronized use of synthetic nitrogen fertilisers^{[3][4][5]}. Several biological pathways occurring in agricultural soils are involved in N₂O emissions. Among all of them, nitrification and denitrification are the main microbial N₂O sources directly affected by soil nitrogen fertilisation, but only denitrification is known to be the largest source of N₂O^[6].

Apart from other organisms, such as archaea and fungi, some facultative bacteria possess the ability to adapt their metabolism to an oxygen-depleted environment in the presence of nitrate as a respiratory substrate through the activation of denitrification. This pathway consists of the dissimilatory reduction of nitrate or nitrite to dinitrogen (N₂) via the gaseous intermediates nitric oxide (NO) and nitrous oxide (N₂O). In this process, specific metalloenzymes are sequentially involved: periplasmic (Nap) or membrane-bound (Nar) nitrate reductases, copper (Cu)-containing (NirK) or cytochrome cd1-containing (NirS) nitrite reductases, nitric oxide reductases (cNor, qNor or CuANor), and nitrous oxide reductase (Nos). The majority of denitrifiers are found in the phylum Proteobacteria, within the domain Bacteria. The alpha-proteobacterium *Paracoccus denitrificans* and the gamma-proteobacteria *Pseudomonas stutzeri* and *Pseudomonas aeruginosa* are the first model organisms where denitrification were widely studied. Reviews covering the physiology, biochemistry and molecular genetics of denitrification have been published elsewhere^{[7][8][9][10][11][12]}. Over recent years, several reports about denitrification in plant endosymbiotic bacteria emerged^{[13][14][15]}. Thanks to their capacity to establish an N₂-fixing symbiotic relationship with plants, these bacteria can contribute to natural N soil enrichment, while reducing the need for chemical fertilisation. Therefore, symbiotic N₂ fixation is considered a process with economic, ecological and agricultural importance. In this process, a mutualist association between soil bacteria, commonly known as rhizobia, and plants of the Fabaceae family is established. Rhizobia may induce the formation of nodules in the legume roots and on the stems of some aquatic legumes; nodules are specialized structures where N₂ fixation takes place^[16].

Bradyrhizobium diazoefficiens, which establishes nitrogen-fixation symbiosis with soybean (*Glycine max*), is considered a model organism in the study of denitrification in rhizobia because it is the only known rhizobia species able to grow under oxygen-limiting conditions with nitrate as sole electron acceptor and, also, to perform the complete denitrification pathway under both free-living and symbiotic conditions^[13]. Denitrification in *B. diazoefficiens* is carried out by a periplasmic nitrate reductase (Nap), encoded by the *napEDABC* operon^[17], a Cu-containing nitrite reductase (NirK), encoded by the *nirK* gene^[18], a cytochrome c-type nitric oxide reductase (cNor), encoded by the *norCBQD* operon^[19], and a Cu-dependent nitrous oxide reductase (Nos), encoded by the *nosRZDFYLYX* genes^[20]. Nap is a functional heterodimer comprising the catalytic subunit NapA of about 90 kDa that contains a bis molybdopterin guanine dinucleotide (Mo[MGD]₂) cofactor and a [4Fe-4S] centre, and NapB (15 kDa) that contains 2 heme c groups and receives electrons from the membrane-bound NapC (25 kDa) which binds 4 heme c groups. NirK is a homotrimer with a predicted molecular mass of about 35 kDa per monomer that contains type 1 and type 2 Cu centres. The catalytic subunit of cNor, NorB, contains heme b and a binuclear active centre (heme b₃ and FeB). NorC is a membrane-anchored protein (16 kDa) that contains heme c. Finally, the catalytic subunit of Nos, NosZ (120–160 kDa), is a homodimer Cu-containing enzyme with two distinct Cu centres (CuA and CuZ).

Similarly to many other denitrifiers, expression of denitrification genes in *B. diazoefficiens* requires both oxygen limitation and the presence of nitrate or a derived nitrogen oxide (NO_x), this control being mediated by the FixLJ-FixK2-NnrR regulatory cascade^{[21][22][23]}. In fact, the expression of *napEDABC*, *nirK* and *nosRZDFYLYX* genes requires microoxic conditions and directly depends on the transcriptional regulator FixK2^{[23][24]}, while expression of *norCBQD* genes relies on NO, being

NnrR the transcriptional regulator which directly interacts with the *norCBQD* promoter^{[23][25]}. In this context, the molecular discriminatory determinants for selective FixK2 recognition and target activation were recently unveiled^[26].

Besides being a source of N₂O, the ecological and environmental importance of denitrification lies in the fact that Nos is the only known enzyme able to remove N₂O from ecosystems^[4], the expression and activity of this enzyme becoming a natural target to effectively reduce N₂O emissions from agricultural soils. Increasing knowledge of the regulation and biochemistry of N₂O metabolism in rhizobia will raise opportunities for the design of effective mitigation strategies to reduce N₂O emissions from legume crops^[27].

Nowadays, new environmental factors are emerging as candidates for controlling denitrification, such as pH^{[28][29]} or Cu^[30]. In the case of Cu, it is an essential cofactor in critical enzymes, such as multicopper oxidases, as well as the Nos and NirK denitrification enzymes. The role of this metal in denitrification has been studied in a wide range of non-symbiotic microorganisms, such as *Pseudomonas perfectomarinus*^[31], *P. stutzeri*^[30], *P. denitrificans*^{[32][33]} and *Achromobacter xylosoxidans*^[32]. Regarding rhizobia, Serventi et al. (2012)^[34] investigated the role of Cu in cytochrome oxidase biogenesis in *B. diazoefficiens*. Nevertheless, studies covering Cu influence on the denitrification pathway in rhizobia are scarce. This study provides an integral view of the involvement of Cu in *B. diazoefficiens* denitrification, analysing the effect of different Cu regimes on gene expression, as well as on the protein levels and activity of the denitrification enzymes in free-living cultures.

2 Materials and Methods

Bacterial strains used in this study are compiled in Table 1.

Table 1: *B. diazoefficiens* strains used in this study.

Strains	Relevant description	Reference
110spc4	Wild-type	^[35]
BG0602	<i>napE-lacZ</i> fusion-containing strain	^[36]
RJ2498	<i>nirK-lacZ</i> fusion-containing strain	^[21]
RJ2499	<i>norC-lacZ</i> fusion-containing strain	^[21]
BG0301	<i>nosR-lacZ</i> fusion-containing strain	^[24]

B. diazoefficiens cells were cultivated routinely under oxic conditions at 30 Celsius degrees in peptone–salts–yeast extract (PSY) medium supplemented with 0.1% L-arabinose, essentially as described by Mesa et al. (2008)^[22]. Buffered Vincent’s minimal medium, here defined as vitamin-free modified Vincent’s minimal medium (BVM,^{[37][38]}) was used in this study, containing the following ingredients (per litre): KH₂PO₄, 2 g; K₂HPO₄, 2 g; NH₄Cl, 840 mg; MgSO₄, 246.48 mg; CaCl₂, 67.63 mg; FeCl₃, 10 mg; MOPS, 2.09 g. This medium was supplemented with 3 g of 1 M arabinose and 1 mL from a mineral solution^[39] consisting of: H₃BO₃, 145 mg; ZnSO₄, 108 mg; Na₂MoO₄, 125 mg; MnCl₂, 4 mg; FeSO₄, 125 mg; CoSO₄, 70 mg; nitrile triacetate, 7 g; CuSO₄, 5 mg. When needed, the medium was supplemented with 10 mM KNO₃ (referred here as BVMN). Final pH was adjusted around 6.8 with 2 M NH₃.

Final Cu concentration in BVM or BVMN as indicated in the original recipe^[37] was 0.02 μ M, referred to in this manuscript as Cu-standard medium (Cu-S). In this study, 13 μ M Cu was used as high Cu conditions (Cu-H); this concentration was also used as Cu-H in previous studies^{[32][33]}. In the case of the Cu-limiting medium (Cu-L), CuSO₄ was omitted from the mineral solution, and

10 μ M bathocuproine disulfonic acid (BCS) (Cu(I) chelator) and 1 mM L-ascorbate (reducer from Cu(II) to Cu(I)) were added to the medium in order to lower Cu availability [32] [34]. Only for the Cu-L medium, glassware was treated overnight with 0.1 M HCl and rinsed afterwards with double-distilled water [34].

After growing under oxic conditions in the PSY medium, *B. diazoefficiens* cells were collected by centrifugation (8000 g, 8 min, 4 Celsius degrees). Next, cells were washed twice with BVM or BVMN and inoculated at an OD₆₀₀ of 0.05 (or 0.2 when needed). For oxic conditions, 3 mL of medium were added to 17-mL tubes. For anoxic conditions, 17-mL tubes were completely filled with medium. For microoxic conditions, 3, 50 and 100 mL of medium were added to 17-mL, 250 and 500-mL rubber stoppered tubes or Erlenmeyer flasks, respectively. The headspace was then filled with a gas mixture consisting of 2% (v/v) oxygen and 98% (v/v) N₂ and both, tubes and flasks, were incubated at 30 Celsius degrees with agitation at 170 rpm.

When needed, antibiotics were added to *B. diazoefficiens* cultures at the following concentrations (μ g/mL): spectinomycin (Spc), 200 (solid cultures), 100 (liquid cultures); streptomycin (Sm), 200 (solid cultures), 100 (liquid cultures); tetracycline (Tc), 100 (solid cultures), 25 (liquid cultures); kanamycin (Km), 200 (solid cultures), 100 (liquid cultures); chloramphenicol (Cm), 20.

Galactosidase activity was analysed using permeabilised cells from at least three independent cultures (3 mL), assayed in triplicate for each strain and condition, as previously described [40]. Specific activity was calculated in Miller Units [41] applying the following formula:

Galactosidase activity in the presence of NO was analyzed by generating this gas chemically according to Bricio et al. (2014) [42] and adding (50 μ M final concentration) to the tubes 5 h before activity measurements.

3 Results




Figure 1: Growth of *B. diazoefficiens* 110spc4 in Cu limitation (Cu-L) (circle), Cu standard (Cu-S) (square) and high Cu (Cu-H) (triangle) BVMN media under oxic (A), anoxic (B) and microoxic (C) conditions. In (B,C), growth in the Cu-S BVM medium was also included (white square). Error bars represent standard error between triplicates, and where not visible, these were smaller than the symbols




Figure 2: Transcriptional expression of denitrification genes monitored as β galactosidase activity from *napE-lacZ* (A), *nirK-lacZ* (B), *norC-lacZ* (C) and *nosR-lacZ* (D) fusions chromosomally integrated in *B. diazoefficiens* 110spc4 grown aerobically in Cu-S (white bars) and microaerobically in Cu-L (light grey bars), Cu-S (dark grey bars) and Cu-H (black bars) BVMN media for 3 days. A post-hoc Tukey HSD test at p minor to 0.05 was applied in (A–D); same lower-case letters in each figure indicate that values are not statistically different. (E) Expression changes of *napE*, *nirK*, *norC* and *nosR* genes in *B. diazoefficiens* 110spc4 grown microaerobically in Cu-L compared with Cu-S measured by qRT-PCR. Data expressed as Miller Units (MU) and Fold Change (FC) are means with standard deviation from at least three independent cultures assayed in triplicate.

4 Conclusions

The main goal of the present work was to investigate the influence of Cu on denitrification in the soybean endosymbiont *B. diazoefficiens*. Taken together, our results suggest that Cu may act as an essential factor in the regulation of the denitrification gene expression. Therefore, Cu could be involved in the denitrification regulatory network and not only acts as a mere enzymatic cofactor of the Cu-dependent enzymes, but also as an important regulatory signal of this process.

References

- [1] AR Ravishankara, John S Daniel, and Robert W Portmann. Nitrous oxide (N_2O): the dominant ozone-depleting substance emitted in the 21st century. *science*, 326(5949):123–125, 2009.
- [2] Priyadarshi R Shukla, J Skeg, E Calvo Buendia, Valérie Masson-Delmotte, H-O Pörtner, DC Roberts, Panmao Zhai, Raphael Slade, Sarah Connors, S van Diemen, et al. Climate change and land: an ipcc special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems. 2019.
- [3] James N Galloway, John D Aber, Jan Willem Erisman, Sybil P Seitzinger, Robert W Howarth, Ellis B Cowling, and B Jack Cosby. The nitrogen cascade. *Bioscience*, 53(4):341–356, 2003.
- [4] David Richardson, Heather Felgate, Nick Watmough, Andrew Thomson, and Elizabeth Baggs. Mitigating release of the potent greenhouse gas N_2O from the nitrogen cycle—could enzymic regulation hold the key? *Trends in biotechnology*, 27(7):388–397, 2009.
- [5] Philip G Taylor and Alan R Townsend. Stoichiometric control of organic carbon–nitrate relationships from soils to the sea. *Nature*, 464(7292):1178–1181, 2010.
- [6] Andrew J Thomson, Georgios Giannopoulos, Jules Pretty, Elizabeth M Baggs, and David J Richardson. Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1593):1157–1168, 2012.
- [7] Walter G Zumft. Cell biology and molecular basis of denitrification. *Microbiology and molecular biology reviews*, 61(4):533–616, 1997.
- [8] RJM van Spanning. The nitrogen cycle: denitrification and its relationship to N_2 fixation.—277–327. Technical report, En: Nitrogen fixation in agriculture, forestry, ecology, and the environment
- [9] Rob JM van Spanning, David J Richardson, and Stuart J Ferguson. Introduction to the biochemistry and molecular biology of denitrification. In *Biology of the nitrogen cycle*, pages 3–20. Elsevier, 2007.
- [10] Beate Kraft, Marc Strous, and Halina E Tegetmeyer. Microbial nitrate respiration—genes, enzymes and environmental distribution. *Journal of biotechnology*, 155(1):104–117, 2011.
- [11] Emilio Bueno, Socorro Mesa, Eulogio J Bedmar, David J Richardson, and Maria J Delgado. Bacterial adaptation of respiration from oxic to microoxic and anoxic conditions: redox control. *Antioxidants & redox signaling*, 16(8):819–852, 2012.
- [12] MJ Torres, Jorg Simon, Gary Rowley, EJ Bedmar, DJ Richardson, AJ Gates, and MJ Delgado. Nitrous oxide metabolism in nitrate reducing bacteria: physiology and regulatory mechanisms. *Advances in microbial physiology*, 68:353–432, 2016.
- [13] EJ Bedmar, EF Robles, and MJ Delgado. The complete denitrification pathway of the symbiotic, nitrogen-fixing bacterium *bradyrhizobium japonicum*, 2005.

- [14] EJ Bedmar, E Bueno, D Correa, MJ Torres, MJ Delgado, and S Mesa. Ecology of denitrification in soils and plant-associated bacteria. *Beneficial plant-microbial interactions: Ecology and applications*, pages 164–182, 2013.
- [15] Ana Salas, Juan J Cabrera, Andrea Jiménez-Leiva, Socorro Mesa, Eulogio J Bedmar, David J Richardson, Andrew J Gates, and María J Delgado. Bacterial nitric oxide metabolism: Recent insights in rhizobia. *Advances in microbial physiology*, 78:259–315, 2021.
- [16] Philip Poole, Vinoy Ramachandran, and Jason Terpolilli. Rhizobia: from saprophytes to endosymbionts. *Nature Reviews Microbiology*, 16(5):291–303, 2018.
- [17] María J Delgado, Nathalie Bonnard, Alvaro Tresierra-Ayala, Eulogio J Bedmar, and Peter Müller. The bradyrhizobium japonicum napedabc genes encoding the periplasmic nitrate reductase are essential for nitrate respiration. *Microbiology*, 149(12):3395–3403, 2003.
- [18] Leonardo Velasco, Socorro Mesa, Maria J Delgado, and Eulogio J Bedmar. Characterization of the nirk gene encoding the respiratory, cu-containing nitrite reductase of bradyrhizobium japonicum. *Biochimica et Biophysica Acta (BBA)-Gene Structure and Expression*, 1521(1-3):130–134, 2001.
- [19] Socorro Mesa, Leonardo Velasco, Maximino E Manzanera, Maria J Delgado, and Eulogio J Bedmar. Characterization of the norcbqd genes, encoding nitric oxide reductase, in the nitrogen fixing bacterium bradyrhizobium japonicum. The genbank accession number for the b. japonicum norcbqd genes reported in this paper is aj132911. *Microbiology*, 148(11):3553–3560, 2002.
- [20] Leonardo Velasco, Socorro Mesa, Chang-ai Xu, María J Delgado, and Eulogio J Bedmar. Molecular characterization of nosrzdflx genes coding for denitrifying nitrous oxide reductase of bradyrhizobium japonicum. *Antonie Van Leeuwenhoek*, 85(3):229–235, 2004.
- [21] Socorro Mesa, Eulogio J Bedmar, Astrid Chanfon, Hauke Hennecke, and Hans-Martin Fischer. Bradyrhizobium japonicum nnrr, a denitrification regulator, expands the fixlj-fixk2 regulatory cascade. *Journal of bacteriology*, 185(13):3978–3982, 2003.
- [22] Socorro Mesa, Felix Hauser, Markus Friberg, Emmanuelle Malaguti, Hans-Martin Fischer, and Hauke Hennecke. Comprehensive assessment of the regulons controlled by the fixlj-fixk2-fixk1 cascade in bradyrhizobium japonicum. *Journal of bacteriology*, 190(20):6568–6579, 2008.
- [23] Emilio Bueno, Eloy F Robles, María J Torres, Tino Krell, Eulogio J Bedmar, María J Delgado, and Socorro Mesa. Disparate response to microoxia and nitrogen oxides of the bradyrhizobium japonicum napedabc, nirk and norcbqd denitrification genes. *Nitric Oxide*, 68:137–149, 2017.
- [24] María J Torres, Emilio Bueno, Andrea Jiménez-Leiva, Juan J Cabrera, Eulogio J Bedmar, Socorro Mesa, and María J Delgado. Fixk2 is the main transcriptional activator of bradyrhizobium diazoefficiens nosrzdflx genes in response to low oxygen. *Frontiers in microbiology*, 8:1621, 2017.
- [25] Andrea Jiménez-Leiva, Juan J Cabrera, Emilio Bueno, María J Torres, Sergio Salazar, Eulogio J Bedmar, María J Delgado, and Socorro Mesa. Expanding the regulon of the bradyrhizobium diazoefficiens nnrr transcription factor: new insights into the denitrification pathway. *Frontiers in microbiology*, page 1926, 2019.

- [26] Juan J Cabrera, Andrea Jiménez-Leiva, Laura Tomás-Gallardo, Sergio Parejo, Sara Casado, María J Torres, Eulogio J Bedmar, María J Delgado, and Socorro Mesa. Dissection of fixk2 protein–dna interaction unveils new insights into bradyrhizobium diazoefficiens lifestyles control. *Environmental Microbiology*, 23(10):6194–6209, 2021.
- [27] Lars R Bakken and Åsa Frostegård. Sources and sinks for n₂o, can microbiologist help to mitigate n₂o emissions?, 2017.
- [28] Cíntia Carreira, Rute F Nunes, Olga Mestre, Isabel Moura, and Sofia R Pauleta. The effect of ph on marinobacter hydrocarbonoclasticus denitrification pathway and nitrous oxide reductase. *JBIC Journal of Biological Inorganic Chemistry*, 25(7):927–940, 2020.
- [29] Alfonso Olaya-Abril, Jesús Hidalgo-Carrillo, Víctor M Luque-Almagro, Carlos Fuentes-Almagro, Francisco J Urbano, Conrado Moreno-Vivián, David J Richardson, and María Dolores Roldán. Effect of ph on the denitrification proteome of the soil bacterium paracoccus denitrificans pd1222. *Scientific reports*, 11(1):1–12, 2021.
- [30] Amanda Black, Pei-Chun L Hsu, Kelly E Hamonts, Tim J Clough, and Leo M Condrón. Influence of copper on expression of nirs, norb and nosz and the transcription and activity of nir, nor and n₂ or in the denitrifying soil bacteria pseudomonas stutzeri. *Microbial biotechnology*, 9(3):381–388, 2016.
- [31] T Matsubara, K Frunzke, and WG Zumft. Modulation by copper of the products of nitrite respiration in pseudomonas perfectomarinus. *Journal of Bacteriology*, 149(3):816–823, 1982.
- [32] Heather Felgate, Georgios Giannopoulos, Matthew J Sullivan, Andrew J Gates, Thomas A Clarke, Elizabeth Baggs, Gary Rowley, and David J Richardson. The impact of copper, nitrate and carbon status on the emission of nitrous oxide by two species of bacteria with biochemically distinct denitrification pathways. *Environmental microbiology*, 14(7):1788–1800, 2012.
- [33] Matthew J Sullivan, Andrew J Gates, Corinne Appia-Ayme, Gary Rowley, and David J Richardson. Copper control of bacterial nitrous oxide emission and its impact on vitamin b12-dependent metabolism. *Proceedings of the National Academy of Sciences*, 110(49):19926–19931, 2013.
- [34] Fabio Serventi, Zeb Andrew Youard, Valérie Murset, Simona Huwiler, Doris Bühler, Miriam Richter, Ronny Luchsinger, Hans-Martin Fischer, Robert Brogioli, Martina Niederer, et al. Copper starvation-inducible protein for cytochrome oxidase biogenesis in bradyrhizobium japonicum. *Journal of Biological Chemistry*, 287(46):38812–38823, 2012.
- [35] Brigitte Regensburger and Hauke Hennecke. Rna polymerase from rhizobium japonicum. *Archives of microbiology*, 135(2):103–109, 1983.
- [36] EF Robles, C Sanchez, N Bonnard, MJ Delgado, and EJ Bedmar. The bradyrhizobium japonicum napedabc genes are controlled by the fixlj-fixk2-nnrr regulatory cascade, 2006.
- [37] JM Vincent et al. A manual for the root-nodule bacteria. *A manual for the root-nodule bacteria.*, 1970.
- [38] Anke Becker, Hélène Bergès, Elizaveta Krol, Claude Bruand, Silvia Rüberg, Delphine Capela, Emmanuelle Lauber, Eliane Meilhoc, Frédéric Ampe, Frans J de Bruijn, et al. Global changes in gene expression in sinorhizobium meliloti 1021 under microoxic and symbiotic conditions. *Molecular Plant-Microbe Interactions*, 17(3):292–303, 2004.

- [39] Paul E Bishop, Juan G Guevara, Jean A Engelke, and Harold J Evans. Relation between glutamine synthetase and nitrogenase activities in the symbiotic association between rhizobium japonicum and glycine max. *Plant Physiology*, 57(4):542–546, 1976.
- [40] Juan J Cabrera, Ana Salas, María J Torres, Eulogio J Bedmar, David J Richardson, Andrew J Gates, and María J Delgado. An integrated biochemical system for nitrate assimilation and nitric oxide detoxification in bradyrhizobium japonicum. *Biochemical Journal*, 473(3):297–309, 2016.
- [41] JH Miller. Miller jh.. experiments in molecular genetics, 1972.
- [42] Carlos Bricio, Laura Álvarez, Marta San Martín, Lici A Schurig-Briccio, Robert B Gennis, and José Berenguer. A third subunit in ancestral cytochrome c-dependent nitric oxide reductases. *Applied and environmental microbiology*, 80(16):4871–4878, 2014.