

BPC-157

Patient Education Handout

Functional Health & Longevity Program · KORB Health Group · For Patient Reference

This handout is for educational reference only. It is not medical advice. Always follow your prescription label, your pharmacy-specific instructions, and your KORB provider's guidance. If you have questions, contact KORB by phone or email (see contact information below).

What Is BPC-157

BPC-157 is a synthetic peptide derived from a protein found naturally in the stomach lining. Its full name is Body Protection Compound 157. It is being studied for its potential role in soft tissue recovery, musculoskeletal support, and gut health.

What BPC-157 May Support

This medication may be considered as an adjunctive support option for patients with goals related to:

- Soft tissue recovery — tendon, ligament, and connective tissue support
- Musculoskeletal wellness during rehabilitation or recovery
- Exercise recovery quality and return-to-activity support
- Gut support in select clinical contexts (discussed with your provider)

Important: Benefits are not guaranteed.

Response to BPC-157 varies significantly between patients. Preclinical data does not guarantee human response. Some patients notice changes during their active course; others do not. Your provider will assess whether to continue at your renewal visit.

What BPC-157 Is Not

Please do not think of it as any of the following:

- A proven injury treatment — current evidence is preclinical (animal studies only)
- A replacement for physical therapy, imaging, specialist evaluation, or surgery
- A guaranteed pain-relief or tissue-repair program
- A therapy you can extend, restart, or adjust on your own without provider direction
- Appropriate if you have active malignancy or pregnancy

How to Use BPC-157

Your prescription label — or the instructions given to you by your KORB clinical provider — is the authoritative source for your dose and directions. (Some pharmacies label vials “as directed by provider” rather than printing exact instructions on the label.) What follows is a general reference. **Do not adjust anything without speaking to your provider first.**

How to inject	Schedule	Timing	Active weeks
Subcutaneous (SQ) injection — fatty tissue under the skin	Every day (7 days a week)	Any consistent time daily — no food restriction	Foundation: an 8-week course (Weeks 1–8) · Gateway / Peak: a shorter 6-week course starting at Week 3 (Weeks 3–8)

Missed a Dose?

- If you miss a dose: skip it — do not double up the next day
- If you are regularly missing doses: contact KORB by phone or email

Traveling with Your Medication

Traveling doesn't have to interrupt your program. Keep medication in its original labeled vial or packaging, and pack it in an insulated bag with an ice pack if it needs refrigeration — don't let it freeze or sit in a hot car or checked bag. If flying, medication and syringes are permitted in carry-on luggage; TSA recommends keeping the original pharmacy label visible. Traveling across time zones can shift your injection time by a few hours without a problem — if you'll be off your usual schedule for several days, contact KORB for guidance.

Storage and Handling

 Refrigerate 36°F – 46°F (2°C – 8°C)	 Do not freeze Freezing damages medication	 Protect from light Keep away from direct sunlight	 Discard at 28 days Write open date on vial when opened
---	---	---	--

Before first use: refrigerate, protect from light, do not freeze or shake. After opening: write the open date on your vial and discard 28 days after first use — even if medication remains. Do not use if the solution appears cloudy, discolored, or contains particles.

What to Expect — Realistic Timeline

Timeline	Phase	What to Expect
Course Weeks 1–2	Getting started	No immediate noticeable effect is common. The therapy works gradually. Continue your schedule.
Foundation: Weeks 1–8 Gateway/Peak: Weeks 3–8	Active course	Recovery quality may begin to shift. Foundation runs a full 8-week course; Gateway and Peak run a shorter 6-week course starting at Week 3. Individual response varies significantly.
Off weeks	Washout — Labs & Lifestyle Focus	No injections during this window — this is intentional. Your Week 12 labs are reviewed during this window, and it's a good time to focus on nutrition, exercise, sleep, and other lifestyle habits before your next course.

BPC-157 data is preclinical only. Your provider will assess whether therapy is appropriate to continue based on your response, tolerability, and goals.

Side Effects and What to Watch For

What You May Notice — Common	What to Do
Injection site redness or irritation	Common and usually mild. Rotate injection sites. Tell KORB provider if persistent.
Nausea or GI discomfort	Occasionally reported. Usually mild and self-limiting. Tell KORB provider if persistent.
Fatigue	Uncommon; tell KORB provider if persistent.

Worth Monitoring — Tell KORB Provider at Your Next Visit

Unusual pain or swelling at injection site	Could indicate infection. Contact KORB promptly if redness spreads or fever develops.
Systemic rash or hives	Tell KORB provider at next visit or sooner if spreading or severe.
Any unexplained systemic symptom	Tell KORB provider. Do not continue without guidance if you have a significant new symptom.

■ Stop injecting and seek emergency care immediately if you experience: trouble breathing, swelling of the face, lips, tongue, or throat, severe rash or hives, chest pain, or any symptom that feels severe or unsafe.

Lab Monitoring

Lab monitoring is a required part of your program — not optional. Your provider uses labs to assess your response and safety.

- Standard longevity panel (CBC, CMP, lipid panel, HbA1c, fasting glucose, fasting insulin, IGF-1, TSH, free T4, free T3, hs-CRP) — baseline and every 16-week renewal.
- PSA — added for men aged 45 and older at every draw.

Your Week 12 lab draw is scheduled approximately 12 weeks into your active cycle. You will receive an email from KORB Operations with instructions on when and where to complete your blood draw at Quest Diagnostics. Please complete your labs at least one week before your scheduled 16-week renewal visit, not the day before — some results, like IGF-1, can take about a week to come back. Unless your lab order says otherwise, plan to fast overnight and have your blood drawn first thing in the morning.

Safety Reminders

- Use BPC-157 only as prescribed. Do not change your dose, course length, or schedule without provider direction.
- BPC-157 is not appropriate during active malignancy or pregnancy.
- Do not restart the course early after your off weeks without provider direction.
- Tell KORB provider if you develop any new medical condition or become pregnant.
- Your prescription label is the authoritative source for your dose and instructions.
- Use a new syringe for every injection — do not reuse syringes.
- Do not combine two products in the same syringe. This is not recommended and may cause adverse effects.
- Follow your state or local regulations for sharps disposal — use an FDA-cleared sharps container; do not place used needles or syringes in household trash unless your local guidelines specifically allow it.

When to Contact KORB and When to Seek Emergency Care

Contact KORB Operations	Patient Portal — Message Your Provider	Seek Emergency Care Immediately
• Questions about timing, storage, or administration Email: info@korbhealth.com Phone: +1 (888) 959-7299 Hours: Mon–Fri, 9 AM–6 PM CT	• Injection-site irritation or redness that does not resolve • Mild but persistent side effects • Questions about whether to continue therapy	• Trouble breathing or throat / face swelling • Severe rash or widespread hives • Chest pain or fainting • Severe or rapidly worsening symptoms • Confusion or severe weakness • Any symptom that feels urgent or unsafe

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once in the morning and once in the afternoon on their clinic days. For anything that doesn't need your provider's direct medical judgment, contact KORB Operations by phone or email instead.

Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.

Key Reminders

Foundation: BPC-157 is an 8-week course (Weeks 1–8 of your cycle).

Gateway / Peak: BPC-157 is a shorter 6-week course starting at Week 3 (Weeks 3–8).

Inject daily at any consistent time — no food restriction.

Write the open date on your vial — discard 28 days after first use.

Preclinical data only — your provider will assess benefit at renewal.

Contact KORB Operations by phone or email for non-urgent questions, or your provider via the Patient Portal for medical questions. Seek emergency care for severe symptoms.