

Sermorelin

Patient Education Handout

Functional Health & Longevity Program · KORB Health Group · For Patient Reference

This handout is for educational reference only. It is not medical advice. Always follow your prescription label, your pharmacy-specific instructions, and your KORB provider's guidance. If you have questions, contact KORB by phone or email (see contact information below).

What Is Sermorelin

Sermorelin is a synthetic peptide that stimulates your pituitary gland to release growth hormone (GH) naturally. It belongs to a class of compounds called growth hormone releasing hormone (GHRH) analogs.

The key distinction: Sermorelin does not replace growth hormone. It encourages your own pituitary gland to produce and release GH through its normal pathways. Because it works with your body's existing system, it is considered a more physiologic approach than direct GH replacement.

Growth hormone release triggered by Sermorelin also stimulates your liver to produce IGF-1 (insulin-like growth factor 1). Your IGF-1 level is one of the key markers your provider will monitor throughout your program.

What Sermorelin May Support

This medication may be considered as an adjunctive support option for patients with goals related to:

- Sleep quality — particularly deep, restorative sleep
- Recovery — exercise recovery and general recovery physiology
- Body composition — lean mass preservation and fat metabolism as part of a lifestyle program
- Energy and wellness — general sense of vitality
- GH-axis support as part of a structured longevity program

Important: Benefits are not guaranteed.

Response to Sermorelin varies significantly between patients. Some patients notice changes within the first few weeks; others notice little or nothing over an entire cycle. This is normal. Sermorelin is an adjunctive support option — not a cure, not a guaranteed treatment, and not a replacement for lifestyle foundations.

What Sermorelin Is Not

Please do not think of it as any of the following:

- Growth hormone replacement — it stimulates your own GH, it does not replace it
- A guaranteed body composition or fat-loss treatment
- An anti-aging cure or age-reversal therapy
- A substitute for nutrition, resistance training, sleep, or recovery habits
- A therapy you can adjust or stop and restart on your own without provider guidance
- Appropriate if you have active malignancy, uncontrolled thyroid disease, or other contraindications

How to Use Sermorelin

Your prescription label — or the instructions given to you by your KORB clinical provider — is the authoritative source for your dose and directions. (Some pharmacies label vials “as directed by provider” rather than printing exact instructions on the label.) What follows is a general reference. **Do not adjust anything without speaking to your provider first.**

How to inject	When to inject	Schedule	Active weeks	Off weeks
Subcutaneous (SQ) injection — fatty tissue under the skin	At bedtime — on an empty stomach, at least 2 hours after eating	6 days ON, 1 day OFF — Monday–Saturday; Sunday off	Weeks 1–12 of your cycle	Weeks 13–16 — washout period before your renewal visit

Timing Matters

Sermorelin should be injected at bedtime, on an empty stomach. GH is released naturally in pulses during sleep — particularly during the first few hours of deep sleep. Injecting at bedtime aligns with this natural rhythm. Food — particularly carbohydrates — can blunt GH release. Wait at least 2 hours after your last meal before injecting.

The 6-On / 1-Off Schedule

Your schedule is Monday through Saturday — 6 days on, Sunday off. The rest day is intentional. Do not inject on Sundays. Do not make up missed doses on your day off.

Missed a Dose?

- If you miss a dose on a scheduled day: skip it — do not double up the next day
- If you are regularly missing doses: contact KORB by phone or email

Traveling with Your Medication

Traveling doesn't have to interrupt your program. Keep medication in its original labeled vial or packaging, and pack it in an insulated bag with an ice pack if it needs refrigeration — don't let it freeze or sit in a hot car or checked bag. If flying, medication and syringes are permitted in carry-on luggage; TSA recommends keeping the original pharmacy label visible. Traveling across time zones can shift your injection time by a few hours without a problem — if you'll be off your usual schedule for several days, contact KORB for guidance.

Storage and Handling



Refrigerate

36°F – 46°F (2°C – 8°C)



Do not freeze

Freezing damages medication



Protect from light

Keep away from direct sunlight



Discard at 28 days

Write open date on vial when opened

Before first use: refrigerate, protect from light, do not freeze or shake. After opening: write the open date on your vial and discard 28 days after first use — even if medication remains. Do not use if the solution appears cloudy, discolored, or contains particles.

What to Expect — Realistic Timeline

Timeline	Phase	What to Expect
Weeks 1–4	Starting out	Some patients notice early changes in sleep quality or morning energy. Many notice nothing yet. This is normal.
Weeks 4–8	Early response	Sleep quality, recovery, and energy may begin to improve. Do not adjust your schedule.
Weeks 8–12	Continued support	Ongoing GH/IGF-1 axis support. Your provider will review IGF-1 and labs at your 16-week renewal visit.
Weeks 13–16	Off Cycle — Labs & Lifestyle Focus	No injections — this is intentional. Your Week 12 labs are reviewed during this window, and this is a good stretch to focus on nutrition, exercise, sleep, and other lifestyle habits before your next cycle. Do not restart early without provider direction.

Do not compare your response to others. Response is highly individual. Age, baseline GH levels, body composition, sleep quality, and lifestyle habits all influence response.

Side Effects and What to Watch For

What You May Notice — Common

What to Do

Injection site redness or irritation

Common, usually mild. Rotate injection sites. Tell KORB provider if persistent.

Flushing or warmth after injection

Typically brief. Usually resolves within minutes.

Headache

May occur early in therapy. Usually mild and short-lived.

Fatigue

Uncommon; tell KORB provider if persistent.

Worth Monitoring — Tell KORB Provider at Your Next Visit

Water retention or puffiness

Can occur with GH stimulation. Tell KORB provider if noticeable.

Joint discomfort or tingling

May indicate dose adjustment is needed. Tell KORB provider.

Sleep changes

Some patients report improved sleep; rarely disrupted sleep early on.

■ Stop injecting and seek emergency care immediately if you experience: trouble breathing, swelling of the face, lips, tongue, or throat, severe rash or hives, chest pain, or any symptom that feels severe or unsafe.

Lab Monitoring

Lab monitoring is a required part of your program — not optional. Your provider uses labs to assess your response and safety.

- IGF-1 — your primary response marker. Target physiologic range; too high is not better.
- HbA1c and fasting glucose — GH stimulation can affect glucose metabolism.
- Thyroid (TSH, free T4, free T3) — Sermorelin requires a functioning thyroid axis.
- CBC, CMP, lipid panel, hs-CRP — standard panel at baseline and every 16-week renewal.
- PSA — added for men aged 45 and older at every draw.

Your Week 12 lab draw is scheduled approximately 12 weeks into your active cycle. You will receive an email from KORB Operations with instructions on when and where to complete your blood draw at Quest Diagnostics. Please complete your labs at least one week before your scheduled 16-week renewal visit, not the day before — some results, like IGF-1, can take about a week to come back. Unless your lab order says otherwise, plan to fast overnight and have your blood drawn first thing in the morning.

Safety Reminders

- Use Sermorelin only as prescribed. Do not change your dose, schedule, or route without provider direction.
- Do not combine with other GH-axis therapies unless specifically instructed by KORB.
- Do not restart after your off weeks early — the washout period is intentional.
- Tell KORB provider if you develop any new medical condition, start a new medication, or become pregnant.
- Your prescription label is the authoritative source for your dose and instructions.
- Use a new syringe for every injection — do not reuse syringes.
- Do not combine two products in the same syringe. This is not recommended and may cause adverse effects.
- Follow your state or local regulations for sharps disposal — use an FDA-cleared sharps container; do not place used needles or syringes in household trash unless your local guidelines specifically allow it.

When to Contact KORB and When to Seek Emergency Care

Contact KORB Operations	Patient Portal — Message Your Provider	Seek Emergency Care Immediately
• Questions about timing, storage, or administration Email: info@korbhealth.com Phone: +1 (888) 959-7299 Hours: Mon–Fri, 9 AM–6 PM CT	• Injection-site irritation or redness that does not resolve • Mild but persistent side effects • Questions about whether to continue therapy	• Trouble breathing or throat / face swelling • Severe rash or widespread hives • Chest pain or fainting • Severe or rapidly worsening symptoms • Confusion or severe weakness • Any symptom that feels urgent or unsafe

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once in the morning and once in the afternoon on their clinic days. For anything that doesn't need your provider's direct medical judgment, contact KORB Operations by phone or email instead.

Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.



Key Reminders

Sermorelin stimulates your own GH — it is not GH replacement.

Inject at bedtime, on an empty stomach, Monday through Saturday. Sunday off.

Write the open date on your vial — discard 28 days after first use.

Your off weeks (Weeks 13–16) are intentional. Do not restart early.

Labs at Week 12 are required before your 16-week renewal visit.

Response varies. More is not better — do not adjust your dose without provider direction.

Contact KORB Operations by phone or email for non-urgent questions, or your provider via the Patient Portal for medical questions. Seek emergency care for severe symptoms.