



Tesamorelin

Patient Education Handout

Functional Health & Longevity Program · KORB Health Group · For Patient Reference

This handout is for educational reference only. It is not medical advice. Always follow your prescription label, your pharmacy-specific instructions, and your KORB provider's guidance. If you have questions, contact KORB by phone or email (see contact information below).

What Is Tesamorelin

Tesamorelin is a synthetic analog of growth hormone releasing factor (GRF). It stimulates the pituitary gland to release growth hormone (GH) naturally — it does not replace GH, it encourages your own body to produce it.

Tesamorelin has the strongest human clinical evidence base among the peptides in the KORB Longevity Program. It is FDA-approved for HIV-associated lipodystrophy and is used in the KORB program off-label for body composition support, specifically targeting visceral adiposity reduction.

Your provider selects your dose based on your goals, labs, and clinical profile. Your IGF-1 and glucose response will be reviewed at every 16-week renewal visit.

What Tesamorelin May Support

This medication may be considered as an adjunctive support option for patients with goals related to:

- Visceral and central adiposity reduction (belly fat around the organs)
- Waist circumference improvement over sustained cycles
- Body composition and metabolic health
- IGF-1 axis support as part of a structured longevity program

Important: Benefits are not guaranteed.

Response varies between patients. Clinical studies show visceral fat reduction over sustained cycles, but individual results depend on baseline body composition, lifestyle, diet, exercise, and other factors. Your provider will assess your response at each renewal visit.

What Tesamorelin Is Not

Please do not think of it as any of the following:

- A weight-loss drug or obesity treatment — it targets visceral adiposity, not general weight loss
- A substitute for nutrition, resistance training, sleep, or recovery habits
- Growth hormone replacement — it stimulates your own GH, it does not replace it
- A guaranteed body composition treatment
- Appropriate in uncontrolled diabetes without careful provider oversight
- A therapy you can adjust, stop, or restart on your own without provider guidance

How to Use Tesamorelin

Your prescription label — or the instructions given to you by your KORB clinical provider — is the authoritative source for your dose and directions. (Some pharmacies label vials “as directed by provider” rather than printing exact instructions on the label.) What follows is a general reference. **Do not adjust anything without speaking to your provider first.**

How to inject	When to inject	Schedule	Active weeks	Off weeks
Subcutaneous (SQ) injection — fatty tissue under the skin	Evening	6 days ON, 1 day OFF — Monday–Saturday; Sunday off	Weeks 1–12 of your cycle	Weeks 13–16 — washout period before your renewal visit

Your specific dose and strength are decisions your KORB provider makes based on your goals, labs, and clinical profile — they are not something to choose or change yourself. Your dose is compounded specifically for you — always follow the exact units on your prescription label, or as directed by your KORB provider if your label doesn't list exact units. Sunday is your rest day every week; do not make up missed doses on your day off.

Missed a Dose?

- If you miss a dose on a scheduled day: skip it — do not double up the next day
- If you are regularly missing doses: contact KORB by phone or email so your provider can help adjust your plan

Traveling with Your Medication

Traveling doesn't have to interrupt your program. Keep medication in its original labeled vial or packaging, and pack it in an insulated bag with an ice pack if it needs refrigeration — don't let it freeze or sit in a hot car or checked bag. If flying, medication and syringes are permitted in carry-on luggage; TSA recommends keeping the original pharmacy label visible. Traveling across time zones can shift your injection time by a few hours without a problem — if you'll be off your usual schedule for several days, contact KORB for guidance.

Storage and Handling



Refrigerate

36°F – 46°F (2°C – 8°C)



Do not freeze

Freezing damages medication



Protect from light

Keep away from direct sunlight



Discard at 28 days

Write open date on vial when opened

Before first use: refrigerate, protect from light, do not freeze or shake. After opening: write the open date on your vial and discard 28 days after first use — even if medication remains. Do not use if the solution appears cloudy, discolored, or contains particles.

What to Expect — Realistic Timeline

Timeline	Phase	What to Expect
Weeks 1–4	Starting out	Most patients notice little initially. Energy or sleep changes are occasionally reported early.
Weeks 4–8	Active course	GH/IGF-1 axis support continues. Changes, if any, begin emerging over time.
Weeks 8–12	Continued support	Ongoing support. Your provider will review your IGF-1 and glucose at renewal.
Weeks 13–16	Off Cycle — Labs & Lifestyle Focus	No injections — this is intentional. Your Week 12 labs are reviewed during this window, and this is a good stretch to focus on nutrition, exercise, sleep, and other lifestyle habits before your next cycle. Do not restart early without provider direction.

Tesamorelin has the strongest human evidence base in this program for visceral fat reduction. Results still vary. Dose may be assessed and adjusted at your 16-week renewal visit based on your IGF-1 response, as tolerated, and as indicated or directed by your KORB provider.



Side Effects and What to Watch For

What You May Notice — Common	What to Do
Injection site redness or irritation	Common and usually mild. Rotate injection sites. Tell KORB provider if persistent.
Flushing or warmth after injection	Typically brief. Usually resolves within minutes.
Headache	May occur early in therapy. Usually mild and short-lived.
Nausea	Occasionally reported. Usually mild and self-limiting.

Worth Monitoring — Tell KORB Provider at Your Next Visit

Water retention or puffiness	Can occur with GH stimulation. Tell KORB provider if noticeable.
Joint discomfort or tingling	May indicate dose adjustment is needed. Tell KORB provider.
Glucose or HbA1c changes	Tesamorelin is glycemically neutral but glucose monitoring is required. Tell KORB provider if you notice changes.

■ **Stop injecting and seek emergency care immediately if you experience: trouble breathing, swelling of the face, lips, tongue, or throat, severe rash or hives, chest pain, or any symptom that feels severe or unsafe.**

Lab Monitoring

Lab monitoring is a required part of your program — not optional. Your provider uses labs to assess your response and safety.

- IGF-1 — primary response marker. Target physiologic range; too high is not better.
- HbA1c and fasting glucose — glycemically neutral but glucose monitoring remains required throughout.
- CBC, CMP, lipid panel, hs-CRP, TSH, free T4, free T3 — standard panel at baseline and every renewal.
- PSA — added for men aged 45 and older at every draw.

Your Week 12 lab draw is scheduled approximately 12 weeks into your active cycle. You will receive an email from KORB Operations with instructions on when and where to complete your blood draw at Quest Diagnostics. Please complete your labs at least one week before your scheduled 16-week renewal visit, not the day before — some results, like IGF-1, can take about a week to come back. Unless your lab order says otherwise, plan to fast overnight and have your blood drawn first thing in the morning.

Safety Reminders

- Use Tesamorelin only as prescribed. Do not change your dose, schedule, or route without provider direction.
- Do not combine with other GH-axis therapies unless specifically instructed by KORB.
- Do not restart after your off weeks early — the washout period is intentional.
- Tell KORB provider if you develop any new medical condition, start a new medication, or become pregnant.
- Your prescription label is the authoritative source for your dose and instructions.
- Use a new syringe for every injection — do not reuse syringes.
- Do not combine two products in the same syringe. This is not recommended and may cause adverse effects.
- Follow your state or local regulations for sharps disposal — use an FDA-cleared sharps container; do not place used needles or syringes in household trash unless your local guidelines specifically allow it.

When to Contact KORB and When to Seek Emergency Care

Contact KORB Operations	Patient Portal — Message Your Provider	Seek Emergency Care Immediately
• Questions about timing, storage, or administration Email: info@korbhealth.com Phone: +1 (888) 959-7299 Hours: Mon–Fri, 9 AM–6 PM CT	• Injection-site irritation or redness that does not resolve • Mild but persistent side effects • Questions about whether to continue therapy	• Trouble breathing or throat / face swelling • Severe rash or widespread hives • Chest pain or fainting • Severe or rapidly worsening symptoms • Confusion or severe weakness • Any symptom that feels urgent or unsafe

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once in the morning and once in the afternoon on their clinic days. For anything that doesn't need your provider's direct medical judgment, contact KORB Operations by phone or email instead.

Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.



Key Reminders

Tesamorelin stimulates your own GH — it is not GH replacement.

Inject every evening, Monday through Saturday. Sunday off.

Always follow your prescription label for your exact dose and units.

Write the open date on your vial — discard 28 days after first use.

Labs at Week 12 are required before your 16-week renewal visit.

Contact KORB Operations by phone or email for non-urgent questions, or your provider via the Patient Portal for medical questions. Seek emergency care for severe symptoms.