

GHK-Cu

Patient Education Handout

Functional Health & Longevity Program · KORB Health Group · For Patient Reference

This handout is for educational reference only. It is not medical advice. Always follow your prescription label, your pharmacy-specific instructions, and your KORB provider's guidance. If you have questions, contact KORB by phone or email (see contact information below).

What Is GHK-Cu

GHK-Cu is a copper-binding peptide that occurs naturally in the human body. It is found in blood plasma, saliva, and urine, and its levels decline with age. GHK-Cu is being studied for its potential role in tissue health, collagen support, wound healing, and anti-inflammatory processes.

GHK-Cu contains copper as part of its structure. Because of this, **lab monitoring of copper and zinc levels is a required part of your program.** Your provider will order the appropriate monitoring labs at baseline and at renewal.

What GHK-Cu May Support

This medication may be considered as an adjunctive support option for patients with goals related to:

- Skin quality, collagen support, and tissue healing
- Anti-inflammatory support in the context of a structured program
- Wound healing and cellular tissue wellness
- Recovery physiology as an adjunctive support option

Important: Benefits are not guaranteed.

Response to GHK-Cu varies significantly between patients. It is not a cosmetic treatment or a guaranteed skin improvement product. Your provider will assess whether therapy is appropriate to continue at your renewal visit.

What GHK-Cu Is Not

Please do not think of it as any of the following:

- A cosmetic product or skin treatment — it is a compounded injectable peptide
- A guaranteed anti-aging or wrinkle-reduction treatment
- Appropriate for patients with Wilson's disease — absolute contraindication
- A replacement for dermatological evaluation, wound care, or specialist care
- A therapy you can extend or restart on your own without provider guidance

How to Use GHK-Cu

Your prescription label — or the instructions given to you by your KORB clinical provider — is the authoritative source for your dose and directions. (Some pharmacies label vials “as directed by provider” rather than printing exact instructions on the label.) What follows is a general reference. **Do not adjust anything without speaking to your provider first.**

How to inject	Schedule	Timing	Gateway / Peak (add-on)
Subcutaneous (SQ) injection — fatty tissue under the skin	3 times per week — evenings (e.g. Mon / Wed / Fri)	Evening — choose consistent days	Weeks 5–8 of your cycle (starts 2 weeks after BPC-157)

Copper / Zinc Lab Monitoring — Required

GHK-Cu requires mandatory copper and zinc monitoring. Your provider will order serum copper, ceruloplasmin, and zinc at baseline and at each renewal while GHK-Cu is active. Therapy may be discontinued if copper or ceruloplasmin levels trend upward.

Choosing Your 3 Days

Choose 3 consistent days each week — for example, Monday, Wednesday, and Friday. Try to keep the same days each week to build a reliable routine.

Missed a Dose?

- If you miss a dose on a scheduled day: skip it — do not double up the next day
- If you are regularly missing doses: contact KORB by phone or email so your provider can help adjust your plan

Traveling with Your Medication

Traveling doesn't have to interrupt your program. Keep medication in its original labeled vial or packaging, and pack it in an insulated bag with an ice pack if it needs refrigeration — don't let it freeze or sit in a hot car or checked bag. If flying, medication and syringes are permitted in carry-on luggage; TSA recommends keeping the original pharmacy label visible. Traveling across time zones can shift your injection time by a few hours without a problem — if you'll be off your usual schedule for several days, contact KORB for guidance.

Storage and Handling

 Refrigerate 36°F – 46°F (2°C – 8°C)	 Do not freeze Freezing damages medication	 Protect from light Keep away from direct sunlight	 Discard at 28 days Write open date on vial when opened
---	---	---	--

Before first use: refrigerate, protect from light, do not freeze or shake. After opening: write the open date on your vial and discard 28 days after first use — even if medication remains. Do not use if the solution appears cloudy, discolored, or contains particles.

What to Expect — Realistic Timeline

Timeline	Phase	What to Expect
Weeks 5–6 (add-on)	Active course begins	Most patients notice nothing initially. This is normal. The therapy works gradually.
Weeks 7–8 (add-on)	Active course continues	Skin quality or tissue changes may begin to emerge. Individual response varies significantly.
After active course	Washout — Labs & Lifestyle Focus	No GHK-Cu injections. Copper/zinc labs will be reviewed at your 16-week renewal visit, and this is a good window to focus on nutrition, exercise, sleep, and other lifestyle habits before your next course.

GHK-Cu requires copper/zinc lab monitoring. Your provider will review these values at baseline and at your 16-week renewal. Therapy may be discontinued if copper or ceruloplasmin levels rise.

Side Effects and What to Watch For

What You May Notice — Common	What to Do
Injection site redness or irritation	Common and usually mild. Rotate injection sites. Tell KORB provider if persistent.
Mild skin changes at injection site	Occasional mild discoloration or sensitivity. Usually self-limiting.
Fatigue	Uncommon; tell KORB provider if persistent.

Worth Monitoring — Tell KORB Provider at Your Next Visit

Copper or ceruloplasmin trending up on labs	Your provider will review at renewal. Therapy may be discontinued if levels rise.
Unusual rash, hives, or skin reaction	Tell KORB provider promptly — copper-containing compounds can occasionally cause hypersensitivity.
Any systemic symptom	Tell KORB provider. Do not continue without guidance if you have a new unexplained systemic symptom.

■ Stop injecting and seek emergency care immediately if you experience: trouble breathing, swelling of the face, lips, tongue, or throat, severe rash or hives, chest pain, or any symptom that feels severe or unsafe.

Lab Monitoring

Lab monitoring is a required part of your program — not optional. Your provider uses labs to assess your response and safety.

- Standard longevity panel (CBC, CMP, lipid panel, HbA1c, fasting glucose, fasting insulin, IGF-1, TSH, free T4, free T3) — baseline and every renewal.
- GHK-Cu copper monitoring: Serum copper, ceruloplasmin, zinc — required at baseline and each renewal while GHK-Cu is active.
- PSA — added for men aged 45 and older at every draw.

Your Week 12 lab draw is scheduled approximately 12 weeks into your active cycle. You will receive an email from KORB Operations with instructions on when and where to complete your blood draw at Quest Diagnostics. Please complete your labs at least one week before your scheduled 16-week renewal visit, not the day before — some results, like IGF-1, can take about a week to come back. Unless your lab order says otherwise, plan to fast overnight and have your blood drawn first thing in the morning.

Safety Reminders

- Use GHK-Cu only as prescribed. Do not change your dose, schedule, or course length without provider direction.
- GHK-Cu is absolutely contraindicated in patients with Wilson's disease (copper metabolism disorder).
- Copper and zinc monitoring labs are required — do not skip your Week 12 lab draw.
- Tell KORB provider if you develop any new medical condition or become pregnant.
- Your prescription label is the authoritative source for your dose and instructions.
- Use a new syringe for every injection — do not reuse syringes.
- Do not combine two products in the same syringe. This is not recommended and may cause adverse effects.
- Follow your state or local regulations for sharps disposal — use an FDA-cleared sharps container; do not place used needles or syringes in household trash unless your local guidelines specifically allow it.

When to Contact KORB and When to Seek Emergency Care

Contact KORB Operations	Patient Portal — Message Your Provider	Seek Emergency Care Immediately
• Questions about timing, storage, or administration Email: info@korbhealth.com Phone: +1 (888) 959-7299 Hours: Mon–Fri, 9 AM–6 PM CT	• Injection-site irritation or redness that does not resolve • Mild but persistent side effects • Questions about whether to continue therapy	• Trouble breathing or throat / face swelling • Severe rash or widespread hives • Chest pain or fainting • Severe or rapidly worsening symptoms • Confusion or severe weakness • Any symptom that feels urgent or unsafe

The Patient Portal is your only HIPAA-compliant way to message your provider directly — providers check portal messages once in the morning and once in the afternoon on their clinic days. For anything that doesn't need your provider's direct medical judgment, contact KORB Operations by phone or email instead.

Phone and email are not appropriate for emergencies. When in doubt, go to urgent care or the emergency room.

Key Reminders

GHK-Cu requires copper/zinc lab monitoring — this is not optional.

Inject 3 times weekly in the evening. Choose consistent days (e.g. Mon / Wed / Fri).

Absolute contraindication: Wilson's disease.

Write the open date on your vial — discard 28 days after first use.

Short-course therapy — 4 weeks, offered as a Gateway / Peak add-on (Weeks 5–8 of your cycle).

Contact KORB Operations by phone or email for non-urgent questions, or your provider via the Patient Portal for medical questions. Seek emergency care for severe symptoms.