


## Abstract

Using a mathematical model of a honeybee colony infested with mites and virus, sensitivity analysis is conducted to determine what parameters the model is most sensitive to. The model consists of a system of five first-order ordinary differential equations and was developed by Dr. Vardayani Ratti. The sensitivity analysis suggests that the model is most sensitive to variations in: max eclosion rate, natural death rates of healthy bees, brood-maintenance coefficient, rate of forager mortality due to homing failure, and recruitment rate of hive bees to forager bees. Also, honeybees are highly dependent on temperature. The model's current parameter values are based on seasonal averages. Extending the model's parameters to match patterns in climate data is in-progress.

## Background

### Western honeybees (*Apis Mellifera*):

- Essential both agriculturally and economically [1]
- Two types of division of labor
  - Reproduction (queen, drones, brood, workers)
  - Colony growth/development (workers)
- Worker division of labor based on tasks performed
  - Hive bees: younger adults who work within hive
  - Forager bees: older adults are recruited to be foragers
- Since early 2000s, colonies have suffered large population losses
- Possible factors:
  - Parasitic mites
  - Viruses
  - Pesticides
  - Climate change



### Mites

- Ectoparasitic mites (*Varroa destructor*)
- Feed on honeybees during development/adult stages
- Carry and transmit deadly viruses inside the colony
- Believed to be one of the primary cause of colony losses [1]
- Dependent on the availability of brood in hive
  - Female mites enter uncapped brood cells to lay eggs
  - Once cell is capped, mite lays eggs and feeds on developing bee
  - Feeding can lead to infection and death of developing bee
  - Once cell is uncapped, mites leave on newly-developed host
- Able to coexist with honeybees in absence of virus
- Beekkeepers apply Varroacide treatment to try to control the mites



### Virus

- More than 20 honeybee viruses are known
- Twelve are carried and transmitted by mites
- Acute Bee Paralysis Virus (ABPV) is common cause of colony failure when transmitted by *Varroa*
- Symptoms of ABPV: paralysis, trembling, inability to fly, hair loss, rapid death

## Research Objectives

- Perform sensitivity analysis to determine what parameters the model is most sensitive to
- Implement refined weather data for more accurate parameter values (in-progress)
  - Physical parameters are highly dependent on temperature
- Reproduce results from 2017 research paper by Ratti et al. [2]

## Mathematical Model


$$\begin{aligned}
 \dot{x}_h &= \underbrace{\mu g(x_h + x_f) h(m)}_{\text{Bee eclosion (birth)}} - \underbrace{\beta_1 m \frac{x_h}{x_h + y + x_f}}_{\text{Bee virus infection}} - \underbrace{(d_1 + \delta_1) x_h}_{\text{Hive bee death}} - \underbrace{\gamma_1 (m + n) x_h}_{\text{Mite-induced death}} \\
 &\quad - \underbrace{x_h R(x_h, x_f)}_{\text{Forager recruitment}} \\
 \dot{x}_f &= \underbrace{x_h R(x_h, x_f)}_{\text{Forager recruitment}} - \underbrace{\beta_1 m \frac{x_f}{x_h + y + x_f}}_{\text{Bee virus infection}} - \underbrace{(p + d_2 + \delta_2) x_f}_{\text{Forager bee death}} - \underbrace{\gamma_2 (m + n) x_f}_{\text{Mite-induced death}} \\
 \dot{y} &= \underbrace{\beta_1 m \frac{x_h + x_f}{x_h + y + x_f}}_{\text{Bee virus infection}} - \underbrace{(d_3 + \delta_3) y}_{\text{Infected bee death}} - \underbrace{\gamma_3 (m + n) y}_{\text{Mite-induced death}} \\
 \dot{m} &= \underbrace{rm \left(1 - \frac{m + n}{\alpha(x_h + y + x_f)}\right)}_{\text{Logistic mite model}} + \underbrace{\beta_2 n \frac{y}{x_h + y + x_f}}_{\text{Mites acquire virus}} - \underbrace{\beta_3 m \frac{x_h + x_f}{x_h + y + x_f}}_{\text{Mites lose virus}} - \underbrace{\delta_4 m}_{\text{Treatment}} \\
 \dot{n} &= \underbrace{rm \left(1 - \frac{m + n}{\alpha(x_h + y + x_f)}\right)}_{\text{Logistic mite model}} - \underbrace{\beta_2 n \frac{y}{x_h + y + x_f}}_{\text{Mites acquire virus}} + \underbrace{\beta_3 m \frac{x_h + x_f}{x_h + y + x_f}}_{\text{Mites lose virus}} - \underbrace{\delta_5 n}_{\text{Treatment}}
 \end{aligned}
 \tag{1}$$

- Parameter values based on seasonal averages
- Division of labor among workers modeled by equations 1 and 2
- $g(x_h + x_f) = \frac{(x_h + x_f)^i}{R^i + (x_h + x_f)^i}$ , where the integer exponent  $i > 1$ ; represents brood-maintenance in colony
- $h(m) = e^{-mk}$ , where  $k > 0$ ; represents how birthrate of bees is affected by presence of mites
- $R(x_h, x_f) = \sigma_1 - \sigma_2 \left(\frac{x_h}{x_h + x_f}\right)$ ; describes effect of social inhibition on forager recruitment


## Sensitivity Analysis Methods

- Calculated two outputs:
  - Percent change in average total population
  - Colony lifespan
- Individually varied parameters +/- 10% of their seasonal avgs.
- Simulated colony for 10 years ( $t = 3650$  days)

## Results



**Figure 1:** Reproduced simulation figures. **Top-left:** bee-only model. **Top-right:** bee-mite model in absence of virus shows coexistence between bees and mites. **Bottom-left:** full bee-mite-virus model with low levels of varroacide treatment; colony collapses after start of 7th year. **Bottom-right:** full bee-mite-virus model with high levels of varroacide treatment; mites and virus are eradicated.




**Figure 2:** Sensitivity analysis results. **Top-left:** percent-change in average total population; low rate of forager mortality due to homing failure ( $p$ ). **Top-right:** percent-change in average total population; critical  $p$ -value. **Bottom-left:** colony lifespan; critical  $p$ -value. **Bottom-right:** colony lifespan; high  $p$ -value.

- In presence of mites and virus, colony depends on level of varroacide treatment (Figure 2, bottom)
- Max eclosion rate  $\mu$  has greatest effect on population size (Figure 2, top)
- Colony's long-term survival depends on variation of specific parameters (Figure 2, bottom)

## Conclusions and Future Work

- Successfully reproduced figures in 2017 research paper
- According to sensitivity analysis, most sensitive parameters:
  - $\mu$ : Max eclosion (birth) rate
  - $d_{1,2}$ : Natural death rates of hive and forager bees
  - $p$ : Rate of forager mortality due to homing failure
  - $\sigma_1$ : Max recruitment rate of hive bees to foragers
- Sensitivity analysis results agree with prior literature [2,3]
- Future work: extend model parameter to match patterns in climate data
  - Realistically, parameter values vary during season
  - Few models found using refined weather data [4]
- Implementation of climate change could provide more realistic scenario (Figure 3)



**Figure 3:** **Top:** plot of max eclosion rate  $\mu$  over two years. **Bottom:** daily climate data from Malahat, Canada (2006) [5].

## References and Acknowledgements

- [1] Jun Chen, Gloria DeGrandi-Hoffman, Vardayani Ratti, Yun Kang. Review on mathematical modeling of honeybee population dynamics. Mathematical Biosciences and Engineering, 2021, 18(6): 9606-9650.
- [2] Vardayani Ratti, Peter G. Kevan, Hermann J. Eberl. A Mathematical Model of Forager Loss in Honeybee Colonies Infested with Varroa destructor and the Acute Bee Paralysis Virus. Bulletin of Mathematical Biology, 2017 June, 79(6):1218-1253.
- [3] Komi Messan, Marisabel Rodriguez Messan, Jun Chen, Gloria DeGrandi-Hoffman, Yun Kang. Population dynamics of Varroa mite and honeybee: Effects of parasitism with age structure and seasonality, Ecological Modelling, vol. 440, 2021.
- [4] Becher, M.A., Grimm, V., Thorbek, P., Horn, J., Kennedy, P.J. and Osborne, J.L. (2014), BEEHAVE: a systems model of honeybee colony dynamics and foraging to explore multifactorial causes of colony failure. J Appl Ecol, 51: 470-482. <https://doi.org/10.1111/1365-2664.12222>
- [5] Daily weather data: <https://open.canada.ca/data/en/dataset/2ac87744-928b-4fd7-a8e-48-225f408b67>

• Thank you to Dr. Vardayani Ratti and the CSC<sup>2</sup> faculty