

A Practitioner's Guide to Estimating School Effects in Centralized Assignment Systems^{*}

Phi Adajar

Massachusetts Institute of Technology

October 6, 2026

[Click here for the latest version](#)

Centralized school assignment with lottery tie-breaking generates quasi-experiments that identify school effects, but existing methods require applicants' rank-order lists and lottery numbers, data that is not always available. I characterize which data from a match identifies a propensity score under which offers are ignorable. When each school draws its own lottery, offers at a school are randomly assigned among applicants rejected by every school they rank above it, so comparing applicants offered and rejected there requires only waitlists. Under a single lottery, this comparison is biased, and the order of each applicant's rejections is also needed. Rank-order lists identify the score without lottery numbers, while the assignment and published aggregates do not in general. I show how to compute each score in finite markets, how schools' waitlist orders reveal which of them share a lottery, and how to handle priorities and screened schools. In simulations, all valid designs recover the true effect, and richer data yields more precise estimates.

^{*}Contact: phiadajar@mit.edu. I thank Parag Pathak and Joshua Angrist for comments on earlier versions of this work. I acknowledge support from the National Science Foundation, the NBER PhD Fellowship for Global Math Talent, and the Jerry A. Hausman Graduate Dissertation Fellowship.

1 Introduction

A core problem in education research is estimating school effectiveness while accounting for the selection of students into schools. Many districts assign seats through deferred acceptance (DA) with lottery tie-breaking. [Abdulkadiroğlu et al. \(2017, 2022\)](#) show that match data from these districts identifies causal effects of attending a school: the match can be used to compute propensity scores, the probability that each applicant is offered a seat at each school. Following [Rosenbaum and Rubin \(1983\)](#), controlling for these scores compares applicants with the same chance of an offer, whose offers differ only because of the lottery.

Doing so requires rich data: rank-order lists, lottery numbers, capacities, and, at screened schools (which admit by test scores or other criteria), the scores used for screening. But this data is often inaccessible due to confidentiality requirements, record-keeping policies, or data degradation. For example, in other research with Chicago Public Schools, I have access to rank-order lists without lottery numbers. A district may instead release only the set of schools that rejected each applicant: the waitlists she is on.

Under what sets of information can a researcher recover propensity scores and then estimate school effects? I characterize which combinations of match data identify a propensity score for which offers are *ignorable* given the score — independent of potential outcomes given the score. The score need not condition on applicants' lists, only on information settled before an applicant reaches the school, which she does once every school she ranks above it has rejected her.

Using a continuum market model, I show that the information required to estimate causal effects depends on whether lottery tie-breakers are reused across schools. When each school draws its own lottery, as under multiple tie-breaking, offers among applicants who reach a school are randomly assigned, so causal identification requires only waitlists. When schools share a lottery, as under single tie-breaking, comparing applicants offered and rejected at a school is biased, and the order in which each applicant was rejected is also required. When a school uses its own lottery, lottery numbers alone yield an alternative design for estimating causal effects. Rank-order lists identify the score without lottery numbers whether or not schools share a lottery, while the assignment and published aggregates do not in general. Finally, waitlist order (each school's ordering of its applicants) alone reveals which schools share a lottery, so the researcher need not know this in advance. [Table 2](#) summarizes these results.

[Section 4](#) turns from identification to practice and computes each score in finite markets, and [Section 5](#) extends the methods to priority groups (such as preference for neighborhood applicants or siblings) and to screened schools. More data buys precision. In a market of 4,000 applicants with multiple tie-breaking, the root mean squared error of the estimated effect is 0.17 with lottery numbers alone, 0.09 with waitlists, and 0.06 with rank-order lists. Relative to estimates with rank-order lists, using lottery numbers alone requires about eight times as many applicants to achieve the same precision. [Section 7](#) turns these results into a step-by-step procedure.

Relative to [Abdulkadiroğlu et al. \(2017, 2022\)](#), who take lists and lottery numbers as given, I ask which subsets of that data suffice. Like [Borusyak and Hull \(2023\)](#), I exploit a known assignment process. The results also bear on the choice between single and multiple tie-breaking. While multiple tie-breaking can decrease student welfare ([Abdulkadiroğlu, Pathak and Roth, 2009](#); [Ashlagi](#)

and Nikzad, 2020), it also allows for simpler estimation of school effects.

2 Setup

Researchers and policymakers are often interested in the causal effect of attending a school s on an outcome, such as test scores, graduation, or behavioral measures. Formally, let D_i indicate that applicant i attends s , and let Y_{1i} and Y_{0i} be her potential outcomes if she does and does not attend, so that $Y_i = Y_{0i}(1 - D_i) + Y_{1i}D_i$. In a constant-effects model,

$$Y_i = \beta_s D_i + \eta_i, \quad (1)$$

where $\beta_s = Y_{1i} - Y_{0i}$ is the effect of attending s and $\eta_i = Y_{0i}$. Regressing Y_i on D_i does not recover β_s due to omitted variable bias. Applicants who attend s differ from those who do not in ways that are correlated with their outcomes, such as through prior achievement or neighborhood location.

Data from centralized assignment systems can account for this selection bias. When a district assigns seats through lottery tie-breaking, applicants with the same preferences are offered seats at random. An offer of a seat at s is then a valid instrument for enrollment at s , after controlling for each applicant's probability of receiving it (Abdulkadiroğlu et al., 2017). Write D_{1i} and D_{0i} for applicant i 's potential enrollment: whether she would attend s if offered a seat there and if not.

2.1 School choice with lottery tie-breaking

Computing that probability requires a model of how the district assigns seats. Let I be a set of n applicants and let $s \in \{0, 1, \dots, S\}$ index schools, with $s = 0$ denoting the outside option. School s has q_s seats, and the outside option has $q_0 > n$, so that all applicants can choose the outside option. Each applicant submits a strict preference ordering over schools, $\succ_i$, where $a \succ_i b$ means that i prefers school a to school b . These preferences define an applicant's *type*, θ_i .¹ Write $B_{\theta_s} = \{b \mid b \succ_{\theta} s\}$ for the set of schools that type θ prefers to s .

Schools choose between applicants using *tie-breakers*.² There are V tie-breakers, indexed by v ; school s uses tie-breaker $v(s)$, and S_v is the set of schools that use tie-breaker v . Applicant i 's value of tie-breaker v is R_{iv} , and schools prefer applicants with smaller values. Write $B_{\theta_s}^v = \{b \in S_v \mid b \succ_{\theta} s\}$ for the schools that type θ prefers to s and that share tie-breaker v .

Assumption 2.1.1 (Lottery tie-breaking). Each R_{iv} is a lottery number, drawn from $U[0, 1]$ independently across applicants and tie-breakers, and independently of types, potential enrollment, and potential outcomes.

Two designs are common. Under *single tie-breaking* (STB), $V = 1$ and all schools use the same lottery number. Under *multiple tie-breaking* (MTB), $V = S$ and $v(s) = s$, so each school draws a separate, independent lottery.

Seats are assigned through student-proposing deferred acceptance (DA), introduced by Gale and Shapley (1962). In the first round of the algorithm, each applicant applies to her most-preferred

¹In practice, schools may also have *priority groups* for certain sets of applicants, including siblings and those living nearby. Section 5.1 extends the results to priorities.

²Section 5.2 discusses non-lottery tie-breakers, such as test scores and interview evaluations.

school, and each school tentatively admits the applicants with the smallest values of its tie-breaker up to its capacity, and rejects the rest. In each later round, rejected applicants apply to their next-most-preferred school, and each school again tentatively admits up to capacity from the set of new applicants and those tentatively held. The algorithm ends when no applicant is rejected, resulting in a match μ . Write μ_i for applicant i 's assignment and $Z_i(s) \equiv \mathbb{1}\{\mu_i = s\}$ for an indicator that i is offered a seat at school s . Applicant i *reaches* s if she applies to it in some round, that is, if she ranks s and every school she prefers to s rejects her.

2.2 Causal effects and identification

The causal effect of attendance on outcomes can be recovered using an instrument. Offers alone are not a valid instrument for enrollment: the probability of an offer depends on where an applicant applies, which may be correlated with her potential outcomes. Suppose, however, that the researcher can find a control set X_i such that potential enrollment and potential outcomes are independent of the offer given these controls. If so, $Z_i(s)$ is a valid instrument for enrollment at s after controlling for X_i .³ A two-stage least squares (2SLS) regression of Y_i on D_i , instrumenting with $Z_i(s)$ and controlling for X_i , recovers β_s .

The control set X_i is often unwieldy, in which case propensity scores are a more practical control. In school choice, the natural control is each applicant's type, $X_i = \theta_i$, because given type, offers depend only on the lottery (Abdulkadiroğlu et al., 2017, Proposition 1). However, nearly every applicant submits a different list, so controlling for type runs into the curse of dimensionality (Hahn, 1998). To address this, Rosenbaum and Rubin (1983) show that instead of using X_i directly, it suffices to control for the *propensity score* $p_s(x) \equiv \mathbb{P}(Z_i(s) = 1 \mid X_i = x)$, the probability that applicants with $X_i = x$ receive an offer at s .

Definition 2.2.1. Offers at s are *ignorable*⁴ given X_i if $Z_i(s) \perp\!\!\!\perp (D_{0i}, D_{1i}, Y_{0i}, Y_{1i}) \mid X_i$.

Lemma 2.2.2 (Rosenbaum and Rubin, 1983, Theorem 3). If offers at s are ignorable given X_i , they are ignorable given $p_s(X_i)$.

By Lemma 2.2.2, it suffices to control for $p_s(X_i)$ rather than X_i , bypassing the curse of dimensionality. The second stage is

$$Y_i = \beta_s D_i + \sum_x \alpha_2(x) \mathbb{1}\{p_s(X_i) = x\} + \eta_i, \quad (2)$$

and the first stage is

$$D_i = \gamma_s Z_i(s) + \sum_x \alpha_1(x) \mathbb{1}\{p_s(X_i) = x\} + \nu_i, \quad (3)$$

where both sums run over the values of $p_s(X_i)$ strictly between 0 and 1. As argued above, because offers are ignorable given $p_s(X_i)$, 2SLS estimation of Equations (2) and (3) recovers β_s . If there are heterogeneous treatment effects, the estimate corresponds to a weighted average of effects for compliers with non-degenerate scores (Imbens and Angrist, 1994).

³Validity also requires relevance, that offers shift enrollment at s ; the exclusion restriction, that an offer affects outcomes only through enrollment at s ; and, with heterogeneous effects, monotonicity, that no applicant enrolls at s only when not offered a seat there. Random assignment delivers none of these by itself.

⁴This is the independence half of what Rosenbaum and Rubin (1983, §1.3) call *strongly ignorable* assignment. "Strongly ignorable" also requires overlap, where each applicant has a non-degenerate propensity score.

The same propensity score can be used to test for balance. Conditional on the score, characteristics fixed before assignment — such as baseline test scores or gender — should not differ between applicants who are and are not offered a seat at s .

2.3 Cutoffs and the large-market score

By [Lemma 2.2.2](#), estimating the effect of s requires only each applicant's probability of an offer at s . [Abdulkadiroğlu et al. \(2017\)](#) show how to compute this score from the data used to run DA: preferences, lottery numbers, and capacities.

The match computed by DA can be summarized using only each school's cutoff, the lottery number of the marginal admitted applicant ([Azevedo and Leshno, 2016](#)). Let c_s be the largest tie-breaker value among applicants offered a seat at s if s is full, and $c_s = 1$ otherwise. The DA assignment is *stable*: no applicant would rather attend a school that has a vacant seat or that seats someone with a larger value of its tie-breaker than hers. Hence applicant i is offered a seat at a school s she ranks if and only if

$$R_{iv(s)} \leq c_s \quad \text{and} \quad R_{iv(b)} > c_b \quad \text{for all } b \in B_{\theta, s}. \quad (4)$$

Each applicant is therefore assigned to her favorite school among those whose cutoffs she clears. In a finite market, cutoffs depend on the realized lottery draw, so $p_s(\theta)$ has no closed form.

Following [Abdulkadiroğlu et al. \(2017\)](#), I characterize the score in a *continuum economy*, a model of large markets: the set of applicants is $I = [0, 1]$, types are distributed according to F (statements about types refer to types in its support), and q_s is the proportion of I that can be seated at school s . DA is defined in the continuum through the same cutoffs, which are now constants rather than random.

If i is rejected from a set of schools during DA, those rejections are informative about her lottery numbers: a rejection at b means that her value of tie-breaker $v(b)$ is larger than c_b . The *most informative disqualification* (MID) of tie-breaker v for type θ at school s is the most lenient cutoff among the schools that θ prefers to s and that use tie-breaker v :

$$MID_{\theta s}^v = \max \{c_b \mid b \in B_{\theta s}^v\}, \quad (5)$$

with $MID_{\theta s}^v = 0$ when $B_{\theta s}^v$ is empty ([Abdulkadiroğlu et al., 2017](#)). If she reaches s , her value of tie-breaker v must be larger than $MID_{\theta s}^v$.

Theorem 2.3.1 ([Abdulkadiroğlu et al., 2017, Theorem 1 and Appendix A.9](#); [Abdulkadiroğlu et al., 2022, Appendix C.1](#)). In a continuum economy satisfying [Assumption 2.1.1](#), for all types θ that rank s ,

$$p_s(\theta) = \prod_v (1 - MID_{\theta s}^v) \times \max \left(\frac{c_s - MID_{\theta s}^{v(s)}}{1 - MID_{\theta s}^{v(s)}}, 0 \right), \quad (6)$$

with $p_s(\theta) = 0$ if $MID_{\theta s}^v = 1$ for some v , and $p_s(\theta) = 0$ for types that do not rank s .

The first factor in Equation (6) is the probability that a type- θ applicant is rejected by every school she prefers to s . The second is the probability that she is offered a seat at s , given those rejections. If a school she prefers to s has room for everyone ($MID_{\theta s}^v = 1$), she never reaches s . Canceling $1 - MID_{\theta s}^{v(s)}$ gives an equivalent form that is continuous in the cutoffs:

$$p_s(\theta) = \prod_{v \neq v(s)} (1 - MID_{\theta s}^v) \times \max(c_s - MID_{\theta s}^{v(s)}, 0). \quad (7)$$

By Assumption 2.1.1, the right side of Equation (7) is the probability of the event in Equation (4) at any cutoffs.

In the special cases of STB and MTB, the general formula for $p_s(\theta)$ simplifies. Under STB, write $MID_{\theta s} \equiv MID_{\theta s}^1 = \max\{c_b \mid b \in B_{\theta s}\}$; then Equation (6) reduces to

$$p_s(\theta) = (1 - MID_{\theta s}) \times \max\left(\frac{c_s - MID_{\theta s}}{1 - MID_{\theta s}}, 0\right) = \max(c_s - MID_{\theta s}, 0). \quad (8)$$

Under MTB, $MID_{\theta s}^{v(s)} = 0$ and $MID_{\theta s}^{v(b)} = c_b$ for $b \in B_{\theta s}$, so

$$p_s(\theta) = \prod_{b \in B_{\theta s}} (1 - c_b) \times c_s. \quad (9)$$

The identification results below rely on applicants who rank a school first, because their score there is the school's cutoff. I therefore assume that every school is someone's first choice.

Assumption 2.3.2 (Rich support). In the continuum economy, every school is ranked first by a positive proportion of applicants.

3 Data requirements to identify a propensity score

Theorem 2.3.1 shows how to compute propensity scores for s when cutoffs (derived from lottery numbers) and preferences are both known. This data is often unavailable; in this section, I show which data identifies a propensity score for s , and which does not. To do so, I use the continuum economy framework of Section 2.3. Suppose throughout this section that s is oversubscribed, so $0 < c_s < 1$.

A *data regime* Ω is the collection of variables the researcher observes; Ω_i denotes those pertaining to applicant i and P_Ω their joint distribution. Unless otherwise stated, the researcher always observes capacities q , the assignment μ , and the tie-breaker map $v(\cdot)$. The researcher may also observe the variables in Table 1.

The researcher may also observe characteristics G_i fixed before the match, such as baseline test scores, which are independent of lottery numbers. Whether i *qualifies* at school b is the indicator $\mathbb{1}\{R_{iv(b)} \leq c_b\}$ that her lottery number clears its cutoff. Let $\mathcal{I}_s \equiv (\theta_i, G_i, (\mathbb{1}\{R_{iv(b)} \leq c_b\})_{b \in B_{\theta_i}})$ collect i 's type, her characteristics, and whether she qualifies at each school she prefers to s .⁵ These are the variables settled before i can apply to s ; they do not include her lottery number at s . Let $A_i(s) \equiv \mathbb{1}\{\mu_i = s \text{ or } s \in W_i\}$ indicate that i reaches s ; by Equation (4), $A_i(s)$ is a function of $\mathcal{I}_s$.

⁵For a type that does not rank s , let $B_{\theta s}$ be the set of all schools it ranks.

Table 1: Match data beyond capacities, the assignment, and the tie-breaker map

Symbol	Name	Description
$\mathcal{A}$	aggregates	published cutoffs and applicant counts
r_i	assigned rank	position of μ_i on i 's list; missing if i is unassigned
W_i	waitlists	schools that rejected i : those she ranked above μ_i (all ranked schools if she is unassigned), or the waitlists she is on
W_i^o	ordered rejections	W_i , ordered by i 's preferences
O_s	waitlist order	school s 's ordering of the applicants who applied to it
θ_i	type	i 's rank-order list
R_i	lottery numbers	i 's value of each tie-breaker

Formally, $W_i \equiv B_{\theta_i \mu_i}$. Variables without a subscript i , such as W or R , denote the variable for all applicants.

Definition 3.0.1. Let $\mathcal{M}$ be a class of continuum markets satisfying [Assumptions 2.1.1](#) and [2.3.2](#) in each of which s is oversubscribed, $0 < c_s < 1$. A data regime Ω *identifies a propensity score for s over $\mathcal{M}$* if and only if there are functions g , k , and h such that, in every market in $\mathcal{M}$:

- (i) $K_i = k(X_i) \in \{0, 1\}$ selects a subsample of applicants;
- (ii) the controls are $X_i = g(\Omega_i; P_\Omega)$, where $g(\Omega_i; P_\Omega)$ takes the same value for any two applicants with the same J_{is} ,⁶ and, among applicants with $K_i = 1$, offers at s are ignorable given X_i ;
- (iii) $p_s(X_i) = h(\Omega_i; P_\Omega)$, so that h computes the propensity score from the data; and
- (iv) a positive proportion of applicants have $K_i = 1$ and $0 < p_s(X_i) < 1$.

Applicants with $K_i = 1$ and $0 < p_s(X_i) < 1$ are offered a seat at s under some lottery draws and not others. They are *randomized at s* , and form the estimation sample for [Equations \(2\) and \(3\)](#).

[Table 2](#) summarizes the results of this section, in the order of the subsections that follow.

Table 2: Data regimes and the propensity scores they identify for school s

Also observed	X_i	$p_s(X_i)$	Randomized at s	STB	MTB
θ, R	θ_i	Equation (6)	$0 < p_s(\theta_i) < 1$	Yes	Yes
θ	θ_i	Equation (6)	$0 < p_s(\theta_i) < 1$	Yes	Yes
W^o	$A_i(s), A_i(s) MID_{\theta_i s}^{v(s)}$	Equation (11)	$A_i(s) = 1, MID_{\theta_i s}^{v(s)} < c_s$	Yes	Yes
W	$A_i(s)$	$c_s A_i(s)$	$A_i(s) = 1$	No	Yes
$\mathcal{A}, r, R$	—	—	—	No	No

Data is in addition to (q, μ, v) . Entries report whether the regime identifies a propensity score for the offer at s ([Definition 3.0.1](#)), and the estimation sample; rows follow [Theorem 3.1.1](#) and [Propositions 3.1.3](#), [3.2.2](#), [3.3.1](#), [3.3.4](#) and [3.4.1](#). By [Corollary 3.5.2](#), the first three rows also hold with waitlist order O in place of v . Under STB, waitlists fail even with waitlist order and lottery numbers added ([Proposition 3.3.4](#)); if some schools share a lottery and others do not, the W row holds at s under the condition of [Proposition 3.3.1](#). With lottery numbers alone, qualification at a school with its own lottery gives an instrument with the compliers of the W row ([Proposition 3.4.2](#)).

3.1 Types

With types and lottery numbers, the researcher observes everything DA uses, which is the setting of [Abdulkadiroğlu et al. \(2017\)](#).

⁶This rules out conditioning on the match result μ itself.

Theorem 3.1.1 ([Abdulkadiroğlu et al., 2017](#), Theorem 1 and Appendix A.9; [Abdulkadiroğlu et al., 2022](#), Appendix C.1). The data (q, μ, v, θ, R) identifies a propensity score for s over any class of markets, with $X_i = \theta_i$ and $p_s(X_i)$ given by [Equation \(6\)](#). In that calculation, the cutoff c_b is the largest $R_{iv(b)}$ among applicants offered a seat at b if b is full, and 1 otherwise.

Lottery numbers and the assignment reveal each cutoff, [Theorem 2.3.1](#) gives the score, and offers are ignorable given θ_i by [Assumption 2.1.1](#). The rest of this section asks how much of this data can be dropped.

Some districts release rank-order lists but not lottery numbers, as in the Chicago example of [Section 1](#). Without lottery numbers, the cutoffs cannot be read off the assignment as the largest admitted lottery number, but they can be generated by simulating DA with new lottery draws.

An equivalent way to do so is to solve for cutoffs using lists and capacities.⁷ Let $p_s(\theta; c)$ denote the score in [Equation \(6\)](#) evaluated at cutoffs c . In the continuum, cutoffs describe a stable assignment if and only if they clear the market ([Azevedo and Leshno, 2016](#)):

$$\int p_s(\theta; c) dF(\theta) \leq q_s, \quad \text{with equality if } c_s < 1, \quad s = 1, \dots, S. \quad (10)$$

Lemma 3.1.2. In a continuum economy satisfying [Assumptions 2.1.1](#) and [2.3.2](#), [Equation \(10\)](#) has exactly one solution $c \in [0, 1]^S$.

Proof. See [Appendix A](#). □

Types and capacities therefore pin down the cutoffs, whichever schools share a lottery.⁸

Proposition 3.1.3. The data (q, μ, v, θ) identifies a propensity score for s over any class of markets, with $X_i = \theta_i$ and $p_s(X_i) = p_s(\theta_i; c)$, the score of [Equation \(6\)](#) at the unique solution c of [Equation \(10\)](#). The assignment μ is not needed.

Proof. See [Appendix A](#). □

3.2 Ordered rejections

A district may release only the schools that rejected each applicant, in the order she ranked them, but neither preference lists nor lottery numbers. The standard lottery design compares applicants who win and lose a school's lottery ([Abdulkadiroğlu et al., 2011](#)). Within a DA match, the analog compares applicants offered a seat at s with those rejected there.

Lemma 3.2.1. In a continuum economy satisfying [Assumption 2.1.1](#), for every type θ that ranks s and reaches it with positive probability,

$$\mathbb{P}(Z_i(s) = 1 \mid A_i(s) = 1, \theta_i = \theta) = \max \left(\frac{c_s - MID_{\theta_s}^{v(s)}}{1 - MID_{\theta_s}^{v(s)}}, 0 \right). \quad (11)$$

⁷When the assignment is observed, each cutoff is also the offer rate among applicants who rank the school first. Solving for cutoffs does not need the assignment.

⁸[Azevedo and Leshno \(2016, Theorem 1\)](#) obtain uniqueness either from full support of applicant types, which requires every combination of lottery numbers across schools and so fails whenever schools share a lottery, or from a regularity condition that gives uniqueness only for almost every vector of capacities. The proof of [Lemma 3.1.2](#) uses instead that lottery numbers are independent of types and across tie-breakers, together with rich support.

Further, the offer $Z_i(s)$ is independent of $(D_{0i}, D_{1i}, Y_{0i}, Y_{1i})$ conditional on $A_i(s) = 1$ and $\theta_i = \theta$.

Proof. See [Appendix A](#). □

Rejections at more-preferred schools truncate the range of possible lottery numbers: reaching s truncates $R_{iv(s)}$ from below at $MID_{\theta, s}^{v(s)}$ and leaves it otherwise uniform. Under MTB, no school ranked above s shares its lottery, so there is no truncation.

Proposition 3.2.2. The data (q, μ, v, W^v) identifies a propensity score for s over any class of markets, with $X_i = (A_i(s), A_i(s)MID_{\theta, s}^{v(s)})$ and $p_s(X_i)$ given by [Equation \(11\)](#) if $A_i(s) = 1$ and 0 otherwise. Each cutoff is identified as the offer rate among applicants who rank the school first, $c_b = \mathbb{P}(Z_i(b) = 1 \mid A_i(b) = 1, B_{\theta, b} = \emptyset)$.

Proof. See [Appendix A](#). □

Ordered rejections reveal which rejections preceded s , and so which schools enter $MID_{\theta, s}^{v(s)}$.

3.3 Waitlists

Without the order of rejections, the researcher knows that i reached s but not which of her rejections came before it. The truncation in [Lemma 3.2.1](#) is then unknown, unless no school ranked above s shares its lottery.

Proposition 3.3.1. If no school other than s uses $v(s)$, then (q, μ, v, W) identifies a propensity score for s over any class of markets in which this condition holds, with $X_i = A_i(s)$ and $p_s(X_i) = c_s A_i(s)$.

Proof. See [Appendix A](#). □

Under this condition, offers among applicants who are offered a seat at s or rejected there are randomly assigned with probability c_s . The proof uses only a weaker condition, that no type that ranks s ranks a school with tie-breaker $v(s)$ above s , but checking it requires applicants' lists.

Corollary 3.3.2. Over the class of MTB markets, (q, μ, v, W) identifies a propensity score for every school, as in [Proposition 3.3.1](#).

Proof. See [Appendix A](#). □

Under STB, every school shares one lottery, so the condition fails whenever there is more than one school. The weaker condition then holds at s if and only if no type that ranks s ranks another school above it. When it fails, waitlists alone need not suffice. The following example, used for the rest of the section, shows why.

Example 3.3.3. Schools a and b have capacities 0.2 and 0.3. Types θ_1 ($a \succ b$), θ_2 ($b \succ a$), and θ_3 (only a) each have mass $1/3$, and applicants have no other characteristics. Under STB, $(c_a, c_b) = (0.3, 0.6)$. By [Lemma 3.2.1](#), among applicants who reach b , a type- θ_1 applicant is offered a seat there with probability $(0.6 - 0.3)/(1 - 0.3) = 3/7$,

and a type- θ_2 applicant with probability 0.6. Yet an applicant of either type who is rejected at both schools has $W_i = \{a, b\}$; only the order of her rejections, (a, b) or (b, a) , tells the two apart. Under MTB, $(c_a, c_b) \approx (0.24, 0.51)$, and applicants of both types who reach b are offered a seat there with probability c_b .

Restricting to applicants whose waitlists reveal their score at b does not help, because which scores are revealed depends on the offer.⁹

Indeed, under STB no controls computed from waitlists make offers at b ignorable, even with more data added.

Proposition 3.3.4. The data $(q, \mu, v, \mathcal{A}, r, R, W, O)$ does not identify a propensity score for s over the class of STB markets.

Proof. See [Appendix A](#). □

The proof uses the market of [Example 3.3.3](#) with $s = b$: a method that worked over the class of STB markets would have to work in this market, and none does. Unassigned applicants of types θ_1 and θ_2 with the same lottery number have the same waitlists, $\{a, b\}$, and the same position in each school's order, so the controls must treat them alike. The proof shows that this puts every applicant offered a seat at b in one cell, although among applicants who reach b , type θ_1 is offered a seat with probability $3/7$ and type θ_2 with probability 0.6.¹⁰ Adding aggregates, assigned ranks, and lottery numbers does not help, because the controls can use lottery numbers only through J_{ib} .

Remark 3.3.5. The designs randomize different applicants. The waitlist design randomizes only applicants offered a seat at s or rejected there. Controlling for types also randomizes applicants who rank s but may be seated at a school they prefer. With heterogeneous effects, the designs therefore estimate effects for different compliers, and richer data, by randomizing more applicants, gives more precise estimates ([Section 6](#)).

3.4 Assignments, aggregates, and lottery numbers

The sparsest data shows where applicants were placed, and possibly their lottery numbers, but not where they applied. It does not identify a propensity score, under any tie-breaking design.

Proposition 3.4.1. The data $(q, \mu, v, \mathcal{A}, r, R)$ does not identify a propensity score for s over the class of MTB markets, or over the class of STB markets.

Proof. See [Appendix A](#). □

Under STB, this follows from [Proposition 3.3.4](#), because data that fails to identify a score still fails with variables removed. Under MTB, the proof again uses the market of [Example 3.3.3](#) with $s = b$. The data cannot separate applicants who lost the lottery at b from applicants who never applied

⁹An offered applicant reveals $B_{\theta,b} = W_i$, while a rejected one does so only if $W_i = \{b\}$, so the selection depends on the offer.

¹⁰If the researcher knew that no applicant ranks a above b , so that there were no type- θ_1 applicants, waitlists would identify the score at b , because the weaker condition after [Proposition 3.3.1](#) would hold. Knowing this usually requires applicants' lists.

there. An unassigned type- θ_3 applicant, who does not rank b , has the same data as unassigned applicants of types θ_1 and θ_2 with the same lottery numbers. Any control the data allows therefore puts unassigned type- θ_3 applicants, who are never offered a seat at b , in the same cell as every applicant who is offered one. As under STB, lottery numbers do not help.

At a school with its own lottery, however, lottery numbers alone support a different design, which instruments with qualification at s itself, $\mathbb{1}\{R_{iv(s)} \leq c_s\}$, instead of the offer. Qualification does not depend on whether i ranks s , and it is computed from (q, μ, v, R) , since c_s is the largest $R_{iv(s)}$ among applicants offered a seat at s .

Proposition 3.4.2. If no school other than s uses $v(s)$, then qualification at s is independent of $(\theta_i, A_i(s), D_{0i}, D_{1i}, Y_{0i}, Y_{1i})$, it changes an applicant's assignment only by moving her to s from a school she ranks below s , and its compliers are applicants who reach s and enroll there if and only if offered a seat. If another school uses $v(s)$, offers at that school need not be independent of qualification at s .

Proof. See [Appendix A](#). □

At a school with its own lottery, qualification therefore has the same propensity score, c_s , for every applicant. It is then a valid instrument for enrollment at s on the full sample, with no controls, whenever the offer satisfies the exclusion restriction. [Table 2](#) still records lottery numbers alone as identifying no score, because c_s is the propensity score of qualification, not of the offer. Qualification's first stage is weaker than the offer's, because most applicants never reach s . When the lottery is shared, qualification satisfies the exclusion restriction only if the other schools on that lottery do not affect outcomes.

3.5 Tie-breakers

Suppose v is unknown. Schools s and s' *share applicants* if a positive proportion of applicants apply to both. Let $\kappa_{ss'}$ be the probability that O_s and $O_{s'}$ order the same way two randomly drawn common applicants, those who apply to both schools.

Proposition 3.5.1. For s and s' that share applicants, $\kappa_{ss'} = 1$ if $v(s) = v(s')$ and $\kappa_{ss'} < 1$ otherwise. Since $\kappa_{ss'}$ is a function of the distribution of waitlist order, the data (q, μ, O) identifies whether two such schools share a tie-breaker.

Proof. See [Appendix A](#). □

The scores need only pairs of schools that share applicants, because a school b ranked above s enters $p_s(\theta)$ only if applicants of type θ can reach s , and those who do apply to both schools. Because only those pairs enter the scores, waitlist order can stand in for the tie-breaker map in each design that uses applicants' lists or the order of their rejections.

Corollary 3.5.2. Without the tie-breaker map, each of the following identifies a propensity score for s over any class of markets:

- (a) (q, μ, O, θ, R) , with the controls and score of [Theorem 3.1.1](#);
- (b) (q, μ, O, θ) , with the controls and score of [Proposition 3.1.3](#); and

(c) (q, μ, O, W^o) , with the controls and score of [Proposition 3.2.2](#).

Proof. See [Appendix A](#). □

In finite markets, agreement on a few common applicants is weak evidence, which [Section 4.3](#) accounts for.

4 Implementation in finite markets

[Section 3](#) shows which data identifies a propensity score in the continuum, where cutoffs are constants. A district's match is finite: its cutoffs depend on the realized lottery draw, and the district may not report which schools share a lottery. This section turns the results of [Section 3](#) into estimators for such a match.

[Section 4.1](#) estimates each score from the data at hand, [Section 4.2](#) shows that, in large markets, the estimates approximate the propensity scores of the finite match, and [Section 4.3](#) recovers which schools share a lottery from waitlist order. The designs that use the offer's score all end the same way: 2SLS of [Equations \(2\) and \(3\)](#) on applicants whose estimated score is strictly between 0 and 1, instrumenting enrollment with the offer and controlling for a dummy for each value of the score.

4.1 Generating propensity scores

The scores of [Section 3](#) depend on the continuum cutoffs, which no finite match reveals. Each recipe below is the finite-market counterpart of the [Section 3](#) result named in its label, with the cutoffs replaced by estimates from the data at hand: realized cutoffs, solved cutoffs, or offer rates. Recipes (a)–(d) give the score of the offer and recipe (e) the score of qualification.

- (a) *Types and lottery numbers* ([Theorem 3.1.1](#)). Evaluate [Equation \(6\)](#) at the realized cutoffs $\hat{c}_n$, where $\hat{c}_{ns}$ is the largest tie-breaker value among applicants offered a seat at s if s is full, and 1 otherwise.
- (b) *Types* ([Proposition 3.1.3](#)). Evaluate [Equation \(6\)](#) at the solved cutoffs $\check{c}_n$, the largest cutoffs that clear the market at the empirical distribution of types. With q_s counting seats, they solve

$$\sum_i p_s(\theta_i; c) \leq q_s, \quad \text{with equality if } c_s < 1, \quad s = 1, \dots, S. \quad (12)$$

By [Lemma 3.1.2](#), applied to the empirical distribution of types, they are the only solution whenever every school is some applicant's first choice. Simulating DA with new lottery draws also gives cutoffs, but each draw adds noise, which solving avoids.¹¹ The solved cutoffs can be computed one school at a time: given the other cutoffs, demand at s , $\sum_i p_s(\theta_i; c)$, is continuous and piecewise linear in c_s , so each school's cutoff has a closed form, and a nonlinear Gauss–Seidel iteration computes $\check{c}_n$.

¹¹In an economy where two applicants compete for one seat and four for two seats, both cells have a score of 1/2. Solved scores pool such cells exactly; scores simulated from repeated lottery draws do not.

Lemma 4.1.1. Start from $c = (1, \dots, 1)$ and update the schools in turn, setting c_s to the largest value in $[0, 1]$ at which $\sum_i p_s(\theta_i; c) \leq q_s$, given the other cut-offs. The cutoffs weakly decrease at every step and converge to $\check{c}_n$, the largest solution of Equation (12).

Proof. See Appendix A. □

- (c) *Ordered rejections (Proposition 3.2.2).* Evaluate Equation (11) at offer rates: estimate each c_b , including c_s , by the offer rate $\bar{c}_{nb}$ among applicants who reach b and rank no school above it with tie-breaker $v(b)$, and compute $\widehat{MID}_{\theta, s}^{v(s)}$ from W_i^o and these estimates. Restrict the sample to applicants with $A_i(s) = 1$ and $\widehat{MID}_{\theta, s}^{v(s)} < \bar{c}_{ns}$. In small markets, these offer rates rest on few applicants, and the ratio in Equation (11) amplifies their errors.
- (d) *Waitlists (Proposition 3.3.1).* If no other school uses $v(s)$, the score is c_s for every applicant who reaches s , so estimate by 2SLS on applicants with $A_i(s) = 1$, with no further controls. Even in a finite market, whether i reaches s then does not depend on her lottery number at s (Lemma A.2).
- (e) *Lottery numbers (Proposition 3.4.2).* Qualification at the realized cutoff has the same score, $\hat{c}_{ns}$, for every applicant. At a school with its own lottery, estimate by 2SLS on the full sample, with no controls,

$$Y_i = \beta_s D_i + \alpha + \eta_i, \quad D_i = \gamma_s \mathbf{1}\{R_{iv(s)} \leq \hat{c}_{ns}\} + \alpha' + v_i, \quad (13)$$

instrumenting enrollment with qualification.¹²

4.2 Large markets

Each recipe uses a continuum score, at estimated cutoffs, in place of the propensity score of the finite match. The two agree in large markets. To make “large markets” precise, I follow Abdulkadiroğlu et al. (2017, 2022) and embed a finite market in an asymptotic sequence. The market of size n consists of the first n applicants of an i.i.d. sequence $(\theta_i, G_i, R_i, D_{0i}, D_{1i}, Y_{0i}, Y_{1i})_{i \geq 1}$ drawn from the continuum economy, with capacities per applicant converging to q , and “almost surely” refers to this sequence. Let $\mathbb{P}_n$ denote probabilities over lottery draws in the market of size n , given applicants’ types, and let $p_{ns}(\theta) = \mathbb{P}_n(Z_i(s) = 1 \mid \theta_i = \theta)$ be the finite-market score of the offer.

Abdulkadiroğlu et al. (2017, Theorem 2) show that under single tie-breaking the score at the realized cutoffs approximates p_{ns} , and Abdulkadiroğlu et al. (2022, Theorem 2) show this in probability under general tie-breaking when all full schools have distinct cutoffs. The following proposition covers all five recipes of Section 4.1, with the score of the offer in parts (a)–(d) and the score of qualification in part (e).

Proposition 4.2.1. Suppose types have finite support and Assumptions 2.1.1 and 2.3.2 hold, and let c solve Equation (10). Along the asymptotic sequence, the following hold almost surely.

¹²Restricting to applicants known to have ranked s , such as those offered a seat there, would select on the lottery.

(a) *Types and lottery numbers.* The realized cutoffs converge, $\hat{c}_n \rightarrow c$, and

$$\sup_{\theta} |p_s(\theta; \hat{c}_n) - p_{ns}(\theta)| \rightarrow 0.$$

(b) *Types.* The solved cutoffs converge, $\check{c}_n \rightarrow c$, and

$$\sup_{\theta} |p_s(\theta; \check{c}_n) - p_{ns}(\theta)| \rightarrow 0.$$

(c) *Ordered rejections.* The offer rates converge, $\bar{c}_{nb} \rightarrow c_b$ for every school b , and for every type θ that reaches s with positive probability in the continuum,

$$\mathbb{P}_n(Z_i(s) = 1 \mid A_i(s) = 1, \theta_i = \theta) \rightarrow \max \left(\frac{c_s - MID_{\theta s}^{v(s)}}{1 - MID_{\theta s}^{v(s)}}, 0 \right).$$

(d) *Waitlists.* If no type that ranks s ranks a school with tie-breaker $v(s)$ above s , then $\mathbb{P}_n(Z_i(s) = 1 \mid A_i(s) = 1, \theta_i = \theta) \rightarrow c_s$ for every type θ that reaches s with positive probability in the continuum.

(e) *Lottery numbers.* Qualification at the realized cutoff satisfies $\mathbb{P}_n(R_{iv(s)} \leq \hat{c}_{ns} \mid \theta_i, D_{0i}, D_{1i}, Y_{0i}, Y_{1i}) \rightarrow c_s$.

Proof. See [Appendix A](#). □

Each part follows from the convergence of the realized cutoffs to the continuum cutoffs, which carries over to the solved cutoffs and the offer rates; part (e) is the finite-market counterpart of [Proposition 3.4.2](#). Given types, simulated lottery numbers have the same distribution as the real ones, so part (a) also holds, in probability, for the score at simulated cutoffs.

As in [Abdulkadiroğlu et al. \(2017\)](#), standard errors treat the estimated score as known. With saturated controls, 2SLS depends on the score only through the partition of applicants into cells with a common score. Because the cutoffs converge, the estimated partition equals the limiting one in large markets, unless the limiting market has an exact tie, such as two equal cutoffs on one lottery or two equal scores. [Section 6](#) reports coverage in finite markets.

4.3 Recovering tie-breakers

When a district reports waitlist order but not the tie-breaker map, [Corollary 3.5.2](#) lets the order stand in for the map. In a finite market, compare the waitlist orders of each pair of schools s and s' on their common applicants, those who applied to both. If the two orders rank any two common applicants differently, the schools use different lotteries. If the orders agree on every common applicant, they may still agree by chance.

To bound that chance, group the common applicants by the schools that rejected them before they applied to the second of s and s' , which their lists or the order of their rejections reveal. Hold cutoffs fixed. If s and s' use different lotteries, the applicants in a group were rejected at the same cutoffs, so by [Assumption 2.1.1](#) their lottery numbers at each of the two schools are i.i.d., independently across the schools. The two orders then agree within every group with probability $\pi_{ss'} = \prod_g 1/m_g!$, where m_g is the size of group g . Join s and s' if $w_{ss'} = -\log \pi_{ss'}$ is at least

$\bar{w} = \log(L/\varepsilon)$, where L is the number of pairs compared, chosen without using the match (for example, all pairs). The probability that any pair with different lotteries is joined by chance is then at most ε . Finally, check with a topological sort that, within each connected component of joined schools, the union of the schools' orderings is acyclic; if it is not, remove joins in increasing order of weight until it is.

Without waitlist order, the match still places applicants offered a seat at each school above those rejected there, which can refute but not confirm a shared lottery.

5 Generalizations

The results so far assume that each school ranks all of its applicants by a lottery. In practice, schools may give priority to some applicants, such as siblings or those living nearby, and screened schools admit applicants by test scores or other criteria. Both extensions follow [Abdulkadiroğlu et al. \(2017, 2022\)](#).

5.1 Priorities

Let $\rho_{\theta s} \in \{0, 1, \dots, J, \infty\}$ be type θ 's priority at s , with smaller values better and ∞ if θ is ineligible; types now include priorities. For now, I assume each applicant is eligible for at most one priority group. School s admits applicants in increasing order of $\rho_{\theta s} + R_{iv(s)}$, so its cutoff c_s lies in $[0, J + 1]$. Its *marginal priority* is $\rho_s = \lfloor c_s \rfloor$ and its *lottery cutoff* is $\tau_s = c_s - \rho_s$. An applicant who reaches s is always seated if $\rho_{\theta s} < \rho_s$ ($t_{\theta s} = a$, always seated), seated if and only if $R_{iv(s)} \leq \tau_s$ if $\rho_{\theta s} = \rho_s$ ($t_{\theta s} = c$, conditionally seated), and never seated if $\rho_{\theta s} > \rho_s$ ($t_{\theta s} = n$, never seated). Following [Abdulkadiroğlu et al. \(2017, 2022\)](#),

$$MID_{\theta s}^v = \begin{cases} 0 & \text{if } B_{\theta s}^v = \emptyset \text{ or } \rho_{\theta b} > \rho_b \text{ for all } b \in B_{\theta s}^v, \\ 1 & \text{if } \rho_{\theta b} < \rho_b \text{ for some } b \in B_{\theta s}^v, \\ \max\{\tau_b \mid b \in B_{\theta s}^v, \rho_{\theta b} = \rho_b\} & \text{otherwise,} \end{cases} \quad (14)$$

and

$$p_s(\theta) = \prod_v (1 - MID_{\theta s}^v) \times \begin{cases} 0 & \text{if } t_{\theta s} = n, \\ 1 & \text{if } t_{\theta s} = a, \\ \max\left(\frac{\tau_s - MID_{\theta s}^{v(s)}}{1 - MID_{\theta s}^{v(s)}}, 0\right) & \text{if } t_{\theta s} = c, \end{cases} \quad (15)$$

with $p_s(\theta) = 0$ if $MID_{\theta s}^v = 1$ for some v .

Transforming the data into a “pseudo-school” market accommodates applicants eligible for multiple priority groups. To do so, split each school into pseudo-schools $s^0, \dots, s^J$, one per priority group, all using $v(s)$, and let each applicant rank the pseudo-schools of her groups in place of s . In the continuum, pseudo-schools of groups better than ρ_s have cutoff 1, that of group ρ_s has cutoff τ_s , and those of worse groups have cutoff 0, so [Equations \(14\) and \(15\)](#) are [Equations \(5\) and \(6\)](#) in the augmented economy.

The scores of [Section 3](#) then carry over, with three changes. Each data regime must record

applicants' priority groups: at s for the designs that use waitlists or ordered rejections, and also at the schools ranked above s for ordered rejections. Cutoffs are identified within priority groups: τ_b is the offer rate in b 's marginal group, and [Assumption 2.3.2](#) must hold within each group, as in [Abdulkadiroğlu et al. \(2017, Assumption 1\)](#). And because [Lemma 3.1.2](#) covers only markets without priorities, the design with types but no lottery numbers reads its cutoffs from these offer rates instead of solving [Equation \(10\)](#).

5.2 Screened schools

Screened schools break ties with non-lottery tie-breakers, such as test scores, which may be correlated with potential outcomes. [Abdulkadiroğlu et al. \(2022\)](#) replace lottery randomization with local randomization near the cutoff. Given a bandwidth δ , an applicant in the marginal priority group at a screened school is never seated there if her tie-breaker exceeds the cutoff by more than δ , always seated if it falls more than δ below, and conditionally seated otherwise. If the tie-breaker has a continuously differentiable distribution, conditionally seated applicants clear the cutoff with probability approaching $1/2$ as $\delta \rightarrow 0$, independently of their potential outcomes. The classification parallels priorities: near its cutoff, each screened school behaves like a lottery school with cutoff $1/2$, and the resulting local DA propensity score replaces each relevant screened school's factor in [Equation \(6\)](#) with $1/2$ ([Abdulkadiroğlu et al., 2022, Theorem 1](#)). That result needs distinct cutoffs at all full schools ([Abdulkadiroğlu et al., 2022, Assumption 2](#)). Estimation adds controls for the screened tie-breakers within a bandwidth δ_n with $\delta_n \rightarrow 0$ and $n\delta_n \rightarrow \infty$ ([Abdulkadiroğlu et al., 2022, Theorem 2](#)).

The local score requires screened tie-breaker values, which no data regime in [Table 2](#) reveals. They are needed for offers at screened schools, and for the type-level score $p_s(\theta)$ of applicants who rank a screened school above s . They are not needed for designs that condition on reaching s . Screened tie-breakers are independent of $R_{iv(s)}$, so [Equation \(11\)](#) holds as stated, and the rejection designs of [Propositions 3.2.2](#) and [3.3.1](#) and the qualification design at a school with its own lottery apply unchanged.

6 Simulations

I simulate two economies to show that each data regime of [Table 2](#) recovers school effects in finite markets, and to compare their precision. The first is Example 2 of [Abdulkadiroğlu et al. \(2017\)](#), replicated m times: schools a , b , and c each have m seats, and m applicants of each of four types rank $c; c \succ b \succ a; b \succ a$; and a . The target school is a ; under STB, its large-market scores there are $0, 1/12, 1/12$, and $5/6$. The second, the *random-preference economy*, has six schools and logit preferences, in which an unobserved ability raises outcomes and the utility of both the target school, which is of middling selectivity, and the most popular school. In both, the effect of enrolling at the target school is 1 , 85 percent of applicants offered a seat enroll, and no other school affects outcomes. Untreated outcomes differ by type in the first economy and with ability in the second, so comparisons that ignore selection are biased. For each economy, tie-breaking design, and data regime, I draw the market and its lotteries 1,000 times and estimate the effect by 2SLS

as in [Section 4. Appendix B](#) gives the exact specification.¹³

Table 3: Effect estimates by data regime

Data	STB			MTB		
	Mean	SD	Coverage	Mean	SD	Coverage
<i>AANP (2017) Example 2, 250 applicants per type</i>						
Offer, no controls	2.96	0.12	0.00	2.66	0.11	0.00
Waitlists	1.88	0.17	0.00	1.00	0.16	0.94
Ordered rejections	0.99	0.16	0.96	1.00	0.16	0.94
Types and lottery numbers	0.99	0.15	0.96	1.00	0.11	0.94
Types, solved cutoffs	0.99	0.15	0.96	1.00	0.11	0.94
Lottery numbers (qualification)	0.98	0.49	0.96	1.00	0.32	0.95
<i>Random preferences, 2,000 applicants</i>						
Offer, no controls	1.79	0.11	0.00	1.66	0.11	0.00
Waitlists	1.18	0.14	0.70	1.00	0.13	0.94
Ordered rejections	1.00	0.15	0.95	1.00	0.13	0.94
Types and lottery numbers	1.00	0.14	0.95	1.00	0.12	0.94
Types, solved cutoffs	1.00	0.14	0.95	1.00	0.12	0.94
Lottery numbers (qualification)	1.01	0.40	0.94	1.00	0.23	0.95

Mean and standard deviation of 2SLS estimates of the effect of enrolling at the target school across 1,000 draws, and the share of draws whose 95 percent confidence interval, from robust standard errors, contains the true effect of 1. Example 2 of [Abdulkadiroğlu et al. \(2017\)](#) has 1,000 applicants and 250 seats per school; the random-preference economy has 2,000 applicants and 1,400 seats at six schools. “Offer, no controls” instruments enrollment with the offer on all applicants. The other rows follow [Section 4.1](#).

Ignoring selection is costly: in [Table 3](#), instrumenting with the offer without controls overstates the effect by 66 to 196 percent.

The order of rejections identifies effects under STB and MTB. Waitlists alone do so only under MTB; under STB they overstate the effect by 88 percent in Example 2 and 18 percent in the random-preference economy. With observable rank-order lists, 2SLS recovers the effect both with lottery numbers and with cutoffs solved from the lists alone, and the two give identical or nearly identical estimates. With saturated controls, only the partition of applicants into score cells matters, and it depends only on the order of the cutoffs. Qualification recovers the effect under both designs here only because no other school affects outcomes ([Proposition 3.4.2](#)).

While all valid designs are centered on the true effect, richer data yields more precise estimates because it randomizes more applicants. Qualification uses every applicant, although most provide no variation; waitlists use the applicants who reach the target school; and lists add applicants who may be seated at schools they prefer ([Remark 3.3.5](#)).

[Figure 1](#) plots the root mean squared error of each valid design against market size. In Example 2 under MTB, with 4,000 applicants, it is 0.17 with lottery numbers, 0.09 with waitlists, and 0.06 with lists. Since the error falls with the square root of market size, matching the precision of lists with lottery numbers alone takes about eight times as many applicants. In the random-preference economy, lists add fewer applicants, and the gains over waitlists are smaller.


[Table B.1](#) in [Appendix B](#) repeats [Table 3](#) with 60 and 300 applicants. Estimates from the valid designs remain centered near 1. In the 60-applicant market, where score cells contain few appli-

¹³Code for all simulations is available from the author.

cants, coverage falls to 0.82–0.83 for ordered rejections and types under STB and to 0.90 for types under MTB, while waitlists and ordered rejections under MTB and qualification under either design stay at 0.93–0.96.

To test the procedure of [Section 4.3](#), I let schools 1–3 of the random-preference economy share one lottery, schools 4 and 5 share another, and school 6 have its own. With $\varepsilon = 0.05$, the procedure recovers this map in all 1,000 draws with 2,000 applicants and in 992 of 1,000 draws with 300, and it never joins schools that use different lotteries. Using the recovered map instead of the true one changes the mean and standard deviation of every estimate that uses the map by less than 0.001.

Figure 1: Precision by data regime and market size



Notes: Root mean squared error of 2SLS estimates of the effect of enrolling at the target school across 300 draws per market size, for the designs that are valid under each tie-breaking design. Under STB, waitlists and lottery numbers alone do not identify a propensity score, so they are omitted. Under MTB, ordered rejections give the same estimates as waitlists, so only waitlists are shown. “Lottery numbers” is the qualification design. Both axes are logarithmic.

7 Practical recommendations

The results reduce to a procedure that estimates the effect of school s from whatever match data a district releases. Steps 1 and 2 prepare the data. Steps 3 and 4 choose a design that instruments with the offer, and step 5 estimates effects with it. Step 6 covers the one design that instruments with qualification instead.

1. *Remove priorities and classify applicants at screened schools.* If schools admit by priority group, split each school into one pseudo-school per group, all using the school's lottery (Section 5.1). At each screened school, classify each applicant as always, never, or conditionally seated from her screened tie-breaker and a bandwidth δ ; conditionally seated applicants clear the cutoff with probability $1/2$ (Section 5.2). The remaining steps then treat each pseudo-school as a school and each screened school as a lottery school with cutoff $1/2$ for its conditionally seated applicants, and every regime must record applicants' priority groups. This classification requires screened tie-breakers; without them, no step identifies effects at a screened school.
2. *Find the tie-breakers.* If it is not known which schools share a lottery, recover it from waitlist order and applicants' lists or ordered rejections (Section 4.3). With unordered waitlists, waitlist order still shows that two schools use different lotteries whenever it ranks two of their common applicants differently.
3. *Do you have rank-order lists?* If so, calculate propensity scores at s using Equation (6): at the realized cutoffs if lottery numbers are observed, and at the solved cutoffs otherwise (Section 4.1). With priorities and no lottery numbers, estimate the cutoffs by offer rates within priority groups instead of solving for them (Section 5.1). Then go to step 5. If not, go to step 4.
4. *Do you have waitlist data?* If so, restrict to applicants who reach s , $A_i(s) = 1$.¹⁴
 - *Do you know the order of each applicant's rejections?* If so, the score is Equation (11), with cutoffs estimated by offer rates (Section 4.1). This works whether or not s shares its lottery.
 - *If not, does s have its own lottery?* If so, as under MTB, the score is the offer rate c_s (Section 4.1). If s shares its lottery, as under STB, waitlists do not identify a propensity score at s in general (Proposition 3.3.4).

If either case gives a score, go to step 5. Without waitlist data, go to step 6.

5. *Estimate effects.* Restrict the sample to applicants randomized at s . Estimate Equations (2) and (3) by 2SLS, instrumenting enrollment at s with the offer and controlling for a dummy for each value of the score and, with screened schools, for the screened tie-breakers within the bandwidth. Test balance by regressing baseline characteristics on the offer with the same controls.
6. *Do you have lottery numbers?* If so, and if s has its own lottery, estimate Equation (13) by 2SLS on the full sample of applicants (Section 4.1). In large markets every applicant qualifies with probability c_s , so no controls are needed. Otherwise, the data does not identify a propensity score for the offer at s in general (Proposition 3.4.1).

¹⁴At a lottery school, this step needs no screened tie-breakers, even when applicants rank screened schools above s : conditioning on reaching s absorbs them (Section 5.2).

Propensity scores from different steps randomize different applicants, so the resulting estimates average effects over different compliers ([Remark 3.3.5](#)). For districts, releasing the propensity score itself always enables estimation of school effects without releasing preferences or lottery numbers. Under MTB, waitlists alone suffice for causal identification, though more information increases precision.

Prior work has shown that MTB can decrease student welfare relative to STB ([Abdulkadiroğlu, Pathak and Roth, 2009](#); [Ashlagi and Nikzad, 2020](#)). However, MTB has a minor benefit for researchers: it identifies school effects from waitlists alone.

References

- Abdulkadiroğlu, Atila, Joshua D. Angrist, Susan M. Dynarski, Thomas J. Kane, and Parag A. Pathak.** 2011. “Accountability and Flexibility in Public Schools: Evidence from Boston’s Charters and Pilots.” *Quarterly Journal of Economics*, 126(2): 699–748.
- Abdulkadiroğlu, Atila, Joshua D. Angrist, Yusuke Narita, and Parag A. Pathak.** 2017. “Research Design Meets Market Design: Using Centralized Assignment for Impact Evaluation.” *Econometrica*, 85(5): 1373–1432.
- Abdulkadiroğlu, Atila, Joshua D. Angrist, Yusuke Narita, and Parag A. Pathak.** 2022. “Breaking Ties: Regression Discontinuity Design Meets Market Design.” *Econometrica*, 90(1): 117–151.
- Abdulkadiroğlu, Atila, Parag A. Pathak, and Alvin E. Roth.** 2009. “Strategy-Proofness versus Efficiency in Matching with Indifferences: Redesigning the NYC High School Match.” *American Economic Review*, 99(5): 1954–1978.
- Ashlagi, Itai, and Afshin Nikzad.** 2020. “What Matters in School Choice Tie-Breaking? How Competition Guides Design.” *Journal of Economic Theory*, 190: 105120.
- Azevedo, Eduardo M., and Jacob D. Leshno.** 2016. “A Supply and Demand Framework for Two-Sided Matching Markets.” *Journal of Political Economy*, 124(5): 1235–1268.
- Borusyak, Kirill, and Peter Hull.** 2023. “Nonrandom Exposure to Exogenous Shocks.” *Econometrica*, 91(6): 2155–2185.
- Gale, David, and Lloyd S. Shapley.** 1962. “College Admissions and the Stability of Marriage.” *American Mathematical Monthly*, 69(1): 9–15.
- Hahn, Jinyong.** 1998. “On the Role of the Propensity Score in Efficient Semiparametric Estimation of Average Treatment Effects.” *Econometrica*, 66(2): 315–331.
- Imbens, Guido W., and Joshua D. Angrist.** 1994. “Identification and Estimation of Local Average Treatment Effects.” *Econometrica*, 62(2): 467–475.
- Rosenbaum, Paul R., and Donald B. Rubin.** 1983. “The Central Role of the Propensity Score in Observational Studies for Causal Effects.” *Biometrika*, 70(1): 41–55.

A Proofs

Lemma A.1. Fix any continuum economy, and write $d_s(c) \equiv \int p_s(\theta; c) dF(\theta)$ for demand at s . Start from $c = (1, \dots, 1)$ and update the schools in turn, setting c_s to the largest value in $[0, 1]$ at which $d_s(c) \leq q_s$, given the other cutoffs. The cutoffs weakly decrease at every step and converge to the largest solution of [Equation \(10\)](#).

Proof. I show that the iterates decrease, stay above every solution, and converge to a solution. By [Equation \(7\)](#), d_s is continuous, nondecreasing in c_s , nonincreasing in c_b for $b \neq s$, and zero at $c_s = 0$, so the update $\varphi_s(c_{-s}) = \max\{x \in [0, 1] : d_s(x, c_{-s}) \leq q_s\}$ exists and is nondecreasing in c_{-s} .

The iterates decrease. The first update of each school starts from 1, so it weakly lowers the cutoff. Suppose every update before a later update of s has weakly lowered the cutoff it changed. Then the other cutoffs are weakly below their values at the previous update of s , and since φ_s is nondecreasing, the new value of c_s is weakly below the old one.

The iterates stay above every solution. Any solution c' satisfies $c'_s \leq \varphi_s(c'_{-s})$. So if $c' \leq c$ before an update of s , as holds at the start, then $c'_s \leq \varphi_s(c'_{-s}) \leq \varphi_s(c_{-s})$ after it, and $c' \leq c$ still holds.

The limit is the largest solution. The cutoffs decrease to a limit c^* , and $c^* \geq c'$ for every solution c' . Demand at s is at most q_s just after each update of s , so $d_s(c^*) \leq q_s$ by continuity. If $c_s^* < 1$, then for each $x \in (c_s^*, 1]$, late updates of s set $c_s < x$, so $d_s(x, c_{-s}) > q_s$ there, and $d_s(x, c_{-s}^*) > q_s$ because $c_{-s} \geq c_{-s}^*$; letting $x \downarrow c_s^*$ gives $d_s(c^*) = q_s$. So c^* solves Equation (10) and is the largest solution. $\square$

Proof of Lemma 3.1.2. A solution exists by Lemma A.1. I show that any two solutions c and c' satisfy $c \geq c'$, so that, by symmetry, they are equal. By Assumption 2.1.1, Equation (7) is the probability of the event in Equation (4) at any cutoffs, so d_s evaluated at any cutoffs is the mass of applicants assigned to s when each applicant is assigned to her favorite school among those whose cutoffs she clears. Suppose that $T = \{t : c_t < c'_t\}$ is nonempty, and let M and M' be the sets of applicants assigned to schools in T at c and at c' . Each $t \in T$ has $c_t < c'_t \leq 1$, so it is full at c , and no school exceeds its capacity at a solution. Hence M has mass $\sum_{t \in T} q_t$, and M' has mass at most $\sum_{t \in T} q_t$.

Next, $M \subseteq M'$. An applicant assigned to $t \in T$ at c clears t at c' , because $R_{iv(t)} \leq c_t < c'_t$. She fails at c every school b she prefers to t , and if $b \notin T$ then $c'_b \leq c_b$, so she fails b at c' as well. So at c' she is assigned to t or to a school in T that she prefers to t . With the masses above, $M' \setminus M$ is null.

Finally, let t^* maximize c_t over T . Applicants who rank t^* first, have $R_{iv(t^*)} \in (c_{t^*}, c'_{t^*}]$, and have $R_{iv} > c_{t^*}$ for every other tie-breaker v have positive mass, by Assumptions 2.1.1 and 2.3.2 and because $c_{t^*} < c'_{t^*} \leq 1$. They are assigned to their first choice t^* at c' , and they fail every school in T at c , because $c_t \leq c_{t^*}$ for each $t \in T$. So they lie in $M' \setminus M$, a contradiction. Hence T is empty, and $c \geq c'$. $\square$

Proof of Proposition 3.1.3. DA is stable, so its cutoffs solve Equation (10) (Azevedo and Leshno, 2016, Lemma 1), and by Lemma 3.1.2 they are its only solution, so the researcher can compute them from q , v , and the distribution of types, without μ . By Theorem 2.3.1, $p_s(\theta_i; c)$ is the offer probability of type θ_i , and offers are ignorable given θ_i by Assumption 2.1.1. Take $K_i = 1$ for all i . Applicants who rank s first, a positive mass by Assumption 2.3.2, have score $c_s \in (0, 1)$. $\square$

Proof of Lemma 3.2.1. An applicant reaches s exactly when she is offered no seat at a school she prefers to s . Applying Equation (4) to these schools in order of her preferences, $A_i(s) = 1$ if and only if $R_{iv(b)} > c_b$ for all $b \in B_{\theta_s}$, that is, $R_{iv} > MID_{\theta_s}^v$ for every tie-breaker v ; she is then offered a seat at s if and only if $R_{iv(s)} \leq c_s$. Write $U_i = (D_{0i}, D_{1i}, Y_{0i}, Y_{1i})$. By Assumption 2.1.1, conditional on $\theta_i = \theta$ and U_i , the lottery numbers are independent uniform draws, so conditional also on $A_i(s) = 1$, $R_{iv(s)}$ is uniform on $(MID_{\theta_s}^{v(s)}, 1]$, an interval of positive length because θ reaches s with positive probability. This gives Equation (11), conditional on U_i as well as without it. Since $Z_i(s)$ is binary, it is therefore independent of U_i given $A_i(s) = 1$ and $\theta_i = \theta$. $\square$

Proof of Proposition 3.2.2. Take $K_i = A_i(s)$. By Lemma 3.2.1, applicants who rank b first are offered a seat at b with probability c_b . They have positive mass by Assumption 2.3.2, and μ_i and W_i^0 reveal them, so each cutoff is identified. For an applicant who reaches s , the data reveals $B_{\theta_i s}$: it is W_i if $\mu_i = s$, and the schools listed before s in W_i^0 otherwise. So the data reveals X_i and, by Equation (11), its score. Since $A_i(s)$ is a function of $\mathcal{J}_{is}$, and $MID_{\theta_i s}^{v(s)}$ is a function of θ_i and the cutoffs, X_i is the same for applicants with the same $\mathcal{J}_{is}$. By Lemma 3.2.1, given $A_i(s) = 1$, the offer probability depends on the type only through $MID_{\theta_i s}^{v(s)}$ and does not depend on $(D_{0i}, D_{1i}, Y_{0i}, Y_{1i})$; given $A_i(s) = 0$, there is no offer. So offers are ignorable given X_i . Applicants who rank s first, a positive mass by Assumption 2.3.2, reach s and have $MID_{\theta_i s}^{v(s)} = 0$, so their score is $c_s \in (0, 1)$. $\square$

Proof of Proposition 3.3.1. Since no other school uses $v(s)$, no type that ranks s ranks a school with tie-breaker $v(s)$ above it, and $MID_{\theta_s}^{v(s)} = 0$ for every type θ that ranks s . By Lemma 3.2.1, the offer rate among applicants who reach s is then c_s for every type, and offers are independent of potential outcomes given $A_i(s) = 1$ and type. Offers are therefore ignorable given $X_i = A_i(s)$, a function of $\mathcal{J}_{is}$ and of (μ_i, W_i) ; take $K_i = A_i(s)$. The cutoff $c_s = \mathbb{P}(Z_i(s) = 1 \mid A_i(s) = 1)$ is identified. By Assumption 2.3.2, a positive mass ranks s first and so reaches it, and their score c_s lies in $(0, 1)$ because s is oversubscribed. $\square$

Proof of Corollary 3.3.2. Under MTB, no two schools share a tie-breaker, so the condition of Proposition 3.3.1 holds at every school. $\square$

Both negative results use the market of Example 3.3.3 with $s = b$. It satisfies Assumptions 2.1.1 and 2.3.2 under both designs, and b is oversubscribed, so a data regime that identifies a propensity score over the class of STB or MTB markets must satisfy Definition 3.0.1 in this market under that design. Set both potential outcomes Y_{0i} and Y_{1i} to 0 for type θ_1 , 1 for type θ_2 , and 2 for type θ_3 , and call this value the applicant's outcome.

Applicants have no other characteristics, so $\mathcal{J}_{ib}$ consists of the type and, for types θ_1 and θ_3 , whether i clears a . By condition (ii), $X_i = f(\mathcal{J}_{ib})$ for some function f , and applicants with the same data have the same X_i . Applicants offered a seat at b are of type θ_2 , or of type θ_1 and fail to clear a . Each proof below shows that f takes one value, x , at these two values of $\mathcal{J}_{ib}$. Then every applicant offered a seat at b has $X_i = x$, so every other value of X_i has score 0, and condition (iv) requires $K_i = 1$ when $X_i = x$. All type- θ_2 applicants, who are the applicants with outcome 1, have $X_i = x$, and a share c_b of them is offered a seat at b . Each proof then finds another outcome whose applicants with $X_i = x$ are offered a seat at b at a lower rate, so offers are not ignorable given X_i among applicants with $K_i = 1$, and condition (ii) fails.

Proof of Proposition 3.3.4. Under STB, an unassigned applicant of type θ_1 or θ_2 has $W_i = \{a, b\}$, r_i missing, and a position in each school's order that depends only on R_i , and both types are unassigned when $R_i > c_b = 0.6$. So unassigned applicants of the two types with the same R_i have the same data, and f takes the same value at type θ_2 and at type θ_1 failing to clear a . The applicants with outcome 0 are of type

θ_1 . Those with $X_i = x$ include every type- θ_1 applicant who fails to clear a , and only these can be offered a seat at b , so among outcome-0 applicants with $X_i = x$, the share offered a seat at b is at most $(0.6 - 0.3)/(1 - 0.3) = 3/7 < c_b$. $\square$

Proof of Proposition 3.4.1. Under STB, functions g , k , and h that satisfied Definition 3.0.1 for $(q, \mu, v, \mathcal{A}, r, R)$ would satisfy it for the larger data of Proposition 3.3.4, which is impossible. Under MTB, the data is (μ_i, r_i, R_i) up to constants. An applicant with $R_{ia} > c_a$ and $R_{ib} > c_b$ is unassigned whatever her type, so applicants of all three types with the same such lottery numbers have the same data, and f takes one value, x , at type θ_2 and at types θ_1 and θ_3 failing to clear a . Type- θ_3 applicants who fail to clear a have outcome 2 and positive mass, and none of them is offered a seat at b . $\square$

Proof of Proposition 3.4.2. By Equation (4), $Z_i(s) = A_i(s)\mathbf{1}\{R_{iv(s)} \leq c_s\}$. If only s uses $v(s)$, then $A_i(s)$, and i 's assignment when $Z_i(s) = 0$, depend only on her type and the other tie-breakers, while qualification at s is a function of $R_{iv(s)}$. By Assumption 2.1.1, qualification is therefore independent of $(\theta_i, A_i(s), D_{0i}, D_{1i}, Y_{0i}, Y_{1i})$. If $A_i(s) = 0$, qualification leaves i 's assignment unchanged; if $A_i(s) = 1$, it moves her to s from the school she is assigned to when rejected there, which she ranks below s . So its compliers are applicants who reach s and enroll there if and only if offered a seat. For the second claim, take STB in Example 3.3.3 with $s = b$. A type- θ_1 applicant qualifies at b if and only if $R_i \leq 0.6$ and is offered a seat at a if and only if $R_i \leq 0.3$, so her probability of an offer at a is $1/2$ if she qualifies at b and 0 if she does not. $\square$

Proof of Proposition 3.5.1. If $v(s) = v(s')$, both schools order common applicants by the same lottery number, so $\kappa_{ss'} = 1$. Otherwise, fix the types of two common applicants i and j . The event that both apply to s and s' is an intersection of events that each involve a single tie-breaker, so, conditional on it, tie-breakers $v(s)$ and $v(s')$ stay independent, and so do the orderings of i and j at s and s' . Each ordering is nondegenerate, since, conditional on applying, each lottery number is uniform on an interval $(\underline{r}, 1]$ with $\underline{r} < 1$. So i precedes j at s with probability $\lambda \in (0, 1)$ and at s' with probability $\lambda' \in (0, 1)$. They are ordered the same way with probability $\lambda\lambda' + (1 - \lambda)(1 - \lambda') < 1$. Averaging over types, $\kappa_{ss'} < 1$. $\square$

Proof of Corollary 3.5.2. By Proposition 3.5.1, waitlist order reveals whether two schools share a tie-breaker whenever they share applicants. It therefore identifies a map that agrees with v on every pair of schools that share applicants: give two schools the same tie-breaker if a chain of such pairs with $\kappa = 1$ connects them, since each link in the chain shares a tie-breaker. Each cutoff c_b is the offer rate at b among applicants who rank b first, a positive mass by Assumption 2.3.2, and each of the three data sets reveals these applicants; so the cutoffs are identified without v . Given the cutoffs, the controls and score in each part depend on v only for applicants who reach s with positive probability, and only through whether the schools in $B_{\theta_i s} \cup \{s\}$ share tie-breakers; every other applicant has score 0 under any map.

An applicant whose type reaches s with positive probability applies to all of these schools with that probability, so, except on a null set of applicants, every pair of them shares applicants. Computing the controls and score with any map that agrees with v on pairs that share applicants therefore gives the result of [Theorem 3.1.1](#), [Proposition 3.1.3](#), or [Proposition 3.2.2](#). $\square$

Proof of [Lemma 4.1.1](#). Dividing demand and capacity by n does not change the update, and it turns [Equation \(12\)](#) into [Equation \(10\)](#) for the continuum economy whose type distribution is the empirical distribution of types, with capacities q_s/n . The iteration is therefore the one in [Lemma A.1](#) for that economy, and [Lemma A.1](#) gives the result. $\square$

[Section 4.1](#) uses the following lemma for the waitlist design.

Lemma A.2. In any finite market, if no school that applicant i ranks above s uses $v(s)$, then, holding fixed all types and every tie-breaker value other than $R_{iv(s)}$, $A_i(s)$ does not depend on $R_{iv(s)}$.

Proof. Until i proposes to s , DA does not use $R_{iv(s)}$: it compares i with other applicants only at schools she has proposed to, and by assumption none of them uses $v(s)$. So DA runs identically under any value of $R_{iv(s)}$ until i proposes to s , and whether she ever proposes to s , which is $A_i(s)$, does not depend on $R_{iv(s)}$. $\square$

The proof of [Proposition 4.2.1](#) rests on one lemma. Along the asymptotic sequence of [Section 4.2](#), let $\hat{c}_n$ be the realized cutoffs.

Lemma A.3. Suppose types have finite support and [Assumptions 2.1.1](#) and [2.3.2](#) hold, and let c be the solution of [Equation \(10\)](#), which is unique by [Lemma 3.1.2](#). Then $\hat{c}_n \rightarrow c$ almost surely; for every school s and every type θ that ranks s ,

$$\mathbb{P}_n(A_i(s) = 1 \mid \theta_i = \theta) \rightarrow \prod_v (1 - MID_{\theta s}^v);$$

and for every type θ , $p_{ns}(\theta) \rightarrow p_s(\theta; c)$, all almost surely.

Proof. By [Lemma 3.1.2](#) and [Azevedo and Leshno \(2016, Lemma 1\)](#), the continuum economy has a unique stable matching, with cutoffs c . By Varadarajan's theorem, the empirical distribution of applicants' types and lottery numbers converges weakly to that of the continuum economy almost surely, and capacity per applicant converges to q , so the markets converge to the continuum economy in the sense of [Azevedo and Leshno \(2016\)](#); their strict-preference assumption holds because lottery numbers have no atoms. By [Azevedo and Leshno \(2016, Theorem 2\)](#), $\hat{c}_n \rightarrow c$ almost surely. Fix a type θ that ranks s and an applicant i of that type; for other types, $A_i(s) = Z_i(s) = 0$ and $p_s(\theta; c) = 0$. By [Equation \(4\)](#) in the market of size n , i reaches s if and only if $R_{iv(b)} > \hat{c}_{nb}$ for all $b \in B_{\theta s}$, and is offered a seat there if, in addition, $R_{iv(s)} \leq \hat{c}_{ns}$. Since $\hat{c}_n \rightarrow c$ and each R_{iv} differs from each c_b almost surely, these indicators converge almost surely to the same indicators at c . Conditioning on the first

n types, as $\mathbb{P}_n$ does, is the same as conditioning on all types, because the indicators depend only on the first n applicants, who are independent of the rest. So by the dominated convergence theorem for conditional expectations, the conditional probabilities converge almost surely to the probabilities of the events at c , which are $\prod_v(1 - MID_{\theta s}^v)$ and, as computed in the proof of [Lemma 3.1.2](#), $p_s(\theta; c)$. $\square$

Proof of Proposition 4.2.1. (a) By [Lemma A.3](#), $\hat{c}_n \rightarrow c$ and $p_{ns}(\theta) \rightarrow p_s(\theta; c)$ almost surely. Since [Equation \(7\)](#) is continuous in the cutoffs, $p_s(\theta; \hat{c}_n) \rightarrow p_s(\theta; c)$ almost surely as well. By the triangle inequality, $p_s(\theta; \hat{c}_n) - p_{ns}(\theta) \rightarrow 0$ almost surely, uniformly in θ because there are finitely many types.

(b) By the strong law of large numbers, the empirical distribution F_n of types satisfies $F_n(\theta) \rightarrow F(\theta)$ almost surely for each of the finitely many types, so demand under F_n , $\sum_{\theta} F_n(\theta) p_s(\theta; c)$, converges to demand under F , $d_s(c) = \sum_{\theta} F(\theta) p_s(\theta; c)$, uniformly in c , because $0 \leq p_s \leq 1$. Dividing [Equation \(12\)](#) by n , demand under F_n at $\check{c}_n$ is at most capacity per applicant, which converges to q_s , with equality if $\check{c}_n < 1$. Let $\bar{c}$ be a limit point of $\check{c}_n$. Along a subsequence converging to $\bar{c}$, demand at $\check{c}_n$ converges to $d_s(\bar{c})$, because d_s is continuous by [Equation \(7\)](#). So $d_s(\bar{c}) \leq q_s$, with equality if $\bar{c} < 1$, because then $\check{c}_n < 1$ eventually along the subsequence. So $\bar{c}$ solves [Equation \(10\)](#), and $\bar{c} = c$ by [Lemma 3.1.2](#). Since $[0, 1]^S$ is compact, $\check{c}_n \rightarrow c$ almost surely. The second claim follows as in (a), with $\check{c}_n$ in place of $\hat{c}_n$.

(c) I first show that the offer rates converge and then find the limit of the offer probability. Fix a school b , and let E_b be the set of types that rank b and rank no school above it with tie-breaker $v(b)$, so that $\bar{c}_{nb}$ is the ratio of $\sum_{i \leq n} \mathbb{1}\{\theta_i \in E_b\} Z_i(b)$ to $\sum_{i \leq n} \mathbb{1}\{\theta_i \in E_b\} A_i(b)$. Fix $\delta > 0$. By [Lemma A.3](#), almost surely every realized cutoff is within δ of its limit once n is large. Then, by [Equation \(4\)](#) in the market of size n , each of $Z_i(b)$ and $A_i(b)$ lies between the indicators of the same event with every cutoff moved by δ , once in the direction that shrinks the event and once in the direction that enlarges it. These bounds are functions of (θ_i, R_i) alone, so by the strong law of large numbers, the averages over $i \leq n$ of these bounds times $\mathbb{1}\{\theta_i \in E_b\}$ converge almost surely to their expectations. As $\delta \rightarrow 0$, these expectations converge to the probabilities of the events at c , because lottery numbers have no atoms. Taking $\delta = 1/k$ for every k , the numerator and denominator of $\bar{c}_{nb}$, divided by n , converge almost surely to the continuum probabilities $\mathbb{P}(\theta_i \in E_b, Z_i(b) = 1)$ and $\mathbb{P}(\theta_i \in E_b, A_i(b) = 1)$. The second is positive by [Assumption 2.3.2](#), because types that rank b first are in E_b and reach b . Since $MID_{\theta b}^{v(b)} = 0$ for every $\theta \in E_b$, [Lemma 3.2.1](#) shows that the ratio of the two is c_b , so $\bar{c}_{nb} \rightarrow c_b$ almost surely.

For the offer probability, by [Lemma A.3](#), $\mathbb{P}_n(Z_i(s) = 1 \mid \theta_i = \theta) \rightarrow p_s(\theta; c)$ and $\mathbb{P}_n(A_i(s) = 1 \mid \theta_i = \theta) \rightarrow \prod_v(1 - MID_{\theta s}^v)$, which is positive because θ reaches s with positive probability. Since $Z_i(s) = 1$ implies $A_i(s) = 1$, the conditional probability is the ratio of the two, which converges to $p_s(\theta; c) / \prod_v(1 - MID_{\theta s}^v)$. By [Equation \(7\)](#), this limit is the right side of the display in (c).

(d) By assumption, $MID_{\theta s}^{v(s)} = 0$ for every type θ that ranks s , so the limit in (c) is c_s .

(e) As in the proof of [Lemma A.3](#), with the event $R_{iv(s)} \leq \hat{c}_{ns}$ in place of the offer, $\mathbb{P}_n(R_{iv(s)} \leq \hat{c}_{ns} \mid \theta_i) \rightarrow \mathbb{P}(R_{iv(s)} \leq c_s) = c_s$ almost surely, by [Assumption 2.1.1](#). The event depends only on types and lottery numbers, and lottery numbers are independent of types and potential outcomes, so conditioning also on $(D_{0i}, D_{1i}, Y_{0i}, Y_{1i})$ leaves its probability unchanged. $\square$

B Additional simulation results

Each draw redraws the market as well as its lotteries. In Example 2 of [Abdulkadiroğlu et al. \(2017\)](#), schools a , b , and c each have m seats, and m applicants have each of the four types of [Section 6](#). An applicant of type $k \in \{1, 2, 3, 4\}$, numbered in the order listed there, has untreated outcome $Y_{0i} = k - 1 + \xi_i$. In the random-preference economy, applicant i has ability $a_i \sim N(0, 1)$ and utility $u_{ij} = \delta_j + a_i \mathbb{1}\{j \in \{1, 4\}\} + e_{ij}$ for schools $j = 1, \dots, 6$, where δ_j falls linearly from 0.5 to -0.5 and each e_{ij} is standard Gumbel. The outside option has utility $e_{i0} - 0.5$, and i ranks the schools she prefers to it, in order of utility. School 4 is the target and school 1 the most popular. Untreated outcomes are $Y_{0i} = a_i + \xi_i$. Capacities are (150, 150, 250, 250, 300, 300) with 2,000 applicants and (20, 20, 35, 35, 45, 45) with 300. In both economies, $\xi_i \sim N(0, 1)$; a random 85 percent of applicants enroll at the target school if and only if offered a seat there, and the rest never enroll there; and outcomes are $Y_i = Y_{0i} + D_i$.

For the procedure of [Section 4.3](#), schools 1–3 of the random-preference economy share one lottery, schools 4 and 5 share another, and school 6 has its own. Each school's waitlist order lists the applicants who applied to it by their lottery numbers there. The procedure examines all $L = 15$ pairs of schools, groups common applicants using their lists, and joins pairs with $w_{ss'} \geq \log(15/0.05)$. Each draw then estimates the effect at school 4 with types and lottery numbers, types alone, and ordered rejections, once with the true map and once with the recovered one.

Table B.1: Effect estimates by data regime in small markets

Data	STB			MTB		
	Mean	SD	Coverage	Mean	SD	Coverage
<i>AANP (2017) Example 2, 15 applicants per type</i>						
Offer, no controls	2.98	0.47	0.01	2.68	0.45	0.04
Waitlists	1.91	0.72	0.68	1.02	0.64	0.94
Ordered rejections	1.01	0.79	0.82	1.02	0.64	0.94
Types and lottery numbers	1.00	0.69	0.83	1.01	0.46	0.90
Types, solved cutoffs	1.00	0.69	0.83	1.01	0.46	0.90
Lottery numbers (qualification)	1.06	2.12	0.93	1.12	1.26	0.96
<i>Random preferences, 300 applicants</i>						
Offer, no controls	1.79	0.29	0.22	1.66	0.29	0.36
Waitlists	1.19	0.34	0.93	1.02	0.33	0.94
Ordered rejections	1.00	0.39	0.95	1.02	0.33	0.94
Types and lottery numbers	1.00	0.37	0.95	1.02	0.32	0.94
Types, solved cutoffs	1.00	0.37	0.95	1.02	0.32	0.94
Lottery numbers (qualification)	0.99	0.96	0.96	1.04	0.58	0.96

As in [Table 3](#), with 15 applicants of each type and 15 seats per school in Example 2 of [Abdulkadiroğlu et al. \(2017\)](#), and 300 applicants and 200 seats in the random-preference economy. In the 60-applicant market under STB, 46 of the 1,000 draws yield no estimate for ordered rejections or the designs that use types; statistics for those rows use the remaining 954 draws.